

Supplementary Figure S1. Western blot analysis confirmed the expression of ACE2 and TMPRSS2 in 293T-ACE2-TMPRSS2 cells.

Plasmids encoding ACE2 and TMPRSS2 were co-transfected into 293T cells to generate a stable cell line (293T-ACE2-TMPRSS2 cells). β -actin was used as the internal control.

Supplementary Figure S2. The gating strategy of Flow cytometry.

Representative gating of CD19⁺ B cells, CD3⁺ T cells, CD4⁺ T cells, CD8⁺ T cells and CD16⁺CD56⁺ NK cells from donor PBMCs is shown. Briefly, live cells are gated as Zombie UV-. Cells were then gated as CD19⁺, CD3⁺ or CD19⁻CD3⁻ cells. T cells were further subdivided into either CD8⁺ or CD4⁺ populations. CD3⁻CD19⁻ cells were then divided into CD56⁺CD16⁺ NK cells. For each assay, 10,000 events were sampled after the exclusion of debris, doublets, and dead cells.

Supplementary Figure S3. Representative ELISpot results.

Representatively positive ELISpot results from different healthy donors. Approximately 1×10^5 PBMCs per well were incubated in the presence of Omicron spike RBD protein (10 μ g/mL) (experimental wells), phytohemagglutinin (PHA, 5 μ g/mL) (positive controls), or DMSO (negative controls) for 36h at 37°C and 5% CO₂. The spot-forming units (SFU) were counted using an automatic ELISpot Reader and the numbers of SFU were shown at the top right of the figure.