

**Supplementary Table S1. GMT values with 95%CI in different subgroups**

	Delta convalescents (n = 36)									Healthy donors (n = 49)				
	All participants (n = 85)	All convalescents	<18y		18-60y (n = 18)			>60y (n=10)		All healthy donors (n = 49)	SARS-CoV-1 convalescents (n = 10)	Other Healthy donors (n = 39)		
			unvaccinated (n = 8)	All 18-60y (n = 18)	Vaccinated (n = 10)	Unvaccinated (n=8)	All > 60y (n = 10)	Vaccinated (n = 5)	Unvaccinated (n = 5)			All	2 <sup>nd</sup> dose (n = 21)	3 <sup>rd</sup> dose (n = 18)
WT	506 (355.8-719.7)	1697.6 (1383-2083)	706.8 (373.8-1337)	2208.2 (2168-2250)	2240.1 (2231-2250)	2168.8 (2077-2265)	2186 (2110-2264)	2243.4 (2240-2246)	2141.1 (2001-2291)	197.4 (93.9-263.8)	488 (283.7-839.5)	157.4 (93.9-263.8)	49.7 (34.3-72)	498.5 (275.3-902.8)
Alpha	313.5 (195.5-502.7)	1183.7 (759.2-1846)	139 (12.6-1533)	1663.7 (1287-2151)	1744.5 (1118-2722)	1567.9 (1143-2150)	1969.8 (1743-2226)	2048.8 (1823-2302)	1908.8 (1472-2475)	76 (36.1-118.8)	114.5 (48.7-269.3)	65.5 (36.1-118.8)	24.5 (15.9-37.9)	103.5 (48.2-222.5)
Beta	188.4 (115.5-307.1)	530.5 (324.2-868.1)	43.9 (1.1-1678)	501.9 (268.4-938.2)	653.1 (262.9-1622)	361.1 (127.9-1020)	1031.4 (625.1-1702)	1251 (670.6-2334)	883.8 (320.6-2436)	38.8 (26.2-61.8)	34.9 (23-53.1)	40.2 (26.2-61.8)	23.2 (15.5-34.7)	54.6 (30.9-96.7)
Delta	232.6 (147.6-366.7)	1100.1 (782.6-1546)	465.5 (125.1-1732)	1350.3 (984.6-1852)	1808.3 (1353-2417)	937.3 (520.8-1687)	1568.5 (1130-2177)	1664.1 (1022-2709)	1496 (769.3-2909)	42.5 (29.6-57.6)	47.4 (22.9-53.1)	41.3 (29.6-57.6)	27 (19.1-38.1)	54.7 (33.9-88.5)
Omicron	130.7 (88.4-193.3)	289.5 (180.9-463.3)	46.4 (16-134.7)	502.6 (283.3-891.7)	801.3 (374.9-1713)	280.5 (117.7-668.8)	216.7 (113.2-414.9)	233.2 (80.8-672.6)	204.3 (56-745.2)	42.6 (31.3-59)	41.6 (17.2-100.8)	42.9 (31.3-59)	37.5 (26.2-53.8)	50 (26.6-94.1)

**Supplementary Table S2. Proportions of lymphocyte subsets in PBMCs from vaccines before and after stimulation.**

	Un-stimulated					Stimulated				
	Total B	Total T	CD4 <sup>+</sup> T	CD8 <sup>+</sup> T	NK	Total B	Total T	CD4 <sup>+</sup> T	CD8 <sup>+</sup> T	NK
	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)
Healthy donors (n = 35)	10.5 ± 3	72±6.5	45.2± 5.9	26.1± 6.3	7.1 ± 4.2	10.4 ± 3.1	71.6 ± 6	45 ± 5.9	26.1 ± 5.9	7.5 ± 4.4
SARS-CoV-1 convalescents (n = 8)	7.6 ± 2.5	74 ± 7.6	48 ± 7.6	25.6 ± 7.1	7.2 ± 3.9	8.1 ± 2.6	73 ± 6.5	48.2 ± 6.6	24.5 ± 7	7.5 ± 3.8
Other healthy donors (n = 27)	11.4 ± 2.6	71.4 ± 6.1	44.4 ± 5.1	26.3 ± 6.2	7.1 ± 4.4	11.1 ± 2.9	71.2 ± 5.9	44.1 ± 5.4	26.6 ± 5.5	7.5 ± 4.6
2 <sup>nd</sup> dose (n = 18)	10.5 ± 3.3	73.4 ± 5.9	46.5 ± 5	26.5 ± 4.9	7.4 ± 4.6	10.6 ± 3.2	72.3 ± 5.9	45.5 ± 5.5	26.3 ± 4.7	7.6 ± 5
3 <sup>rd</sup> dose (n = 16)	10.4 ± 3.2	71.1 ± 6.6	43.7 ± 6.8	26.5 ± 7.1	6.7 ± 4	10.2 ± 3.1	71.5 ± 5.8	44.3 ± 6.5	26.8 ± 6.5	7.4 ± 3.8

Data were presented as mean ± SD. Only one of the SARS-CoV-1 convalescents remained unvaccinated.

**Supplementary Table S3. Descriptive statistics of SFUs per 10<sup>6</sup> PBMCs in healthy donors in ELISpot experiment.**

	Un-stimulated	Stimulated
Healthy donors (n = 32)	125.9 ± 105.9	188.1 ± 147.9
SARS-CoV-1 convalescents (n = 10)	177 ± 124.8	256 ± 119.2
Other healthy donors (n = 22)	102.7 ± 89.9	157.3 ± 151.7
2 <sup>nd</sup> dose (n = 15)	124.7 ± 98	198 ± 166
3 <sup>rd</sup> dose (n = 16)	123.8 ± 118.4	177.5 ± 138.6

Data were presented as mean ± SD.