Prediction Model for Detection of Sporadic Pancreatic Cancer (PRO-TECT) in a Population-Based Cohort Using Machine Learning and Further Validation in a Prospective Study

Authors:

Wansu Chen, PhD¹, Yichen Zhou, MS¹, Fagen Xie, PhD¹, Rebecca K. Butler, ScM¹, Christie Y. Jeon, ScD², Tiffany Q. Luong, MPH¹, Yu-Chen Lin, MPH², Eva Lustigova, MPH¹, Joseph R. Pisegna, MD³, Sungjin Kim, MS², Bechien U. Wu, MD, MPH⁴.

Affiliations:

¹ Kaiser Permanente Southern California Research and Evaluation, Pasadena, CA

² Cedars-Sinai Medical Center, Los Angeles, CA

³ Division of Gastroenterology and Hepatology, VA Greater Los Angeles Healthcare System, Los Angeles, CA and Departments of Medicine and Human Genetics David Geffen School of Medicine at UCLA

⁴ Center for Pancreatic Care, Department of Gastroenterology, Los Angeles Medical Center, Southern California Permanente Medical Group, Los Angeles, CA

Address where the work was conducted:

100 S Los Robles, 2nd Floor, Pasadena, CA 91101

Correspondence:

Wansu Chen, Ph.D.

Department of Research and Evaluation

Kaiser Permanente Southern California

100 S Los Robles, 2nd Floor, Pasadena, CA 91101

Email: Wansu.Chen@KP.org

Abbreviations

ALT = Alanine transaminase

AUC = area under the curve

CI = confidence interval

DoD = Department of Defense

EHR = electronic health record

GND = Greenwood-Nam-D'Agostino

HbA1C = glycated hemoglobin

ICD-9-CM = Ninth Revision of International Classification of Diseases, Clinical Modification

ICD-10-CM = Tenth Revision of International Classification of Diseases, Clinical Modification

IR = Incidence rate

KPSC = Kaiser Permanente Southern California

NDI = National Death Index

NOD = new onset diabetes

PC = pancreatic cancer

PDAC = pancreatic ductal adenocarcinoma

PPV = positive predictive value

RSF = Random Survival Forest

SEER = Surveillance, Epidemiology, and End Results

VA = Veterans Affairs

Author Conflict of Interest / Study Support

- Guarantor of the article:** Dr. Wansu Chen accepts full responsibility for the conduct of the study. She has had access to the data and has control of the decision of the decision to publish.
- Specific author contributions:** Wansu Chen: Conceptualization, Methodology, Software, Validation, Investigation, Resources, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision; Yichen Zhou: Methodology, Software, Validation, Formal analysis, Investigation, Data Curation, Writing - Review & Editing, Visualization; Fagen Xie: Methodology, Software, Formal analysis, Investigation, Data Curation, Writing - Review & Editing; Rebecca Butler: Methodology, Software, Formal analysis, Investigation, Data Curation, Writing - Review & Editing; Christie Jeon: Conceptualization, Validation, Investigation, Writing - Review & Editing, Supervision; Tiffany Luong: Writing - Review & Editing, Project administration; Yu-Chen Lin: Software, Validation, Investigation, Writing - Review & Editing; Eva Lustigova: Resources, Writing - Review & Editing, Supervision, Project administration; Joseph Pisegna: Writing - Review & Editing; Sungjin Kim: Software, Validation, Investigation, Writing - Review & Editing;

Bechien Wu: Conceptualization, Methodology, Validation, Resources, Writing - Review & Editing, Supervision, Funding acquisition. All authors have approved the final submitted draft.

- **Financial support:** Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health under Award Number R01CA230442. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.
- **Potential competing interests:** The authors declare they have no conflict of interest for this study.

Manuscript Type: Original Investigation

Word count: Abstract 250; Text 3,116

Study Highlights

What Is Known

- Patients with pancreatic cancer are often diagnosed at late stages.
- Early detection is needed to impact the natural history of disease progression and improve patient survival.

What Is New Here

- Machine-learning was used to develop a population-based model for early detection of pancreatic cancer. The model was internally and externally validated in cohorts of 1.8 million and 2.6 million individuals, respectively.
- Calibration was excellent in prospective pilot testing for detection of pancreatic malignancy.

ABSTRACT

OBJECTIVES: There is currently no widely accepted approach to screening for pancreatic cancer (PC). We aimed to develop and validate a risk prediction model for PC across two health systems using electronic health records (EHR).

METHODS: This retrospective cohort study consisted of patients 50-84 years of age meeting utilization criteria in 2008-2017 at Kaiser Permanente Southern California (KPSC, model training, internal validation) and the Veterans Affairs (VA, external validation). ‘Random survival forests’ models were built to identify the most relevant predictors from >500 variables and to predict PC within 18 months of cohort entry. A prospective study was then conducted in KPSC to assess feasibility of the model for real-time implementation.

RESULTS: The KPSC cohort consisted of 1.8 million patients (mean age 61.6) with 1,792 PC cases. The estimated 18-month incidence rate of PC was 0.77 (95% CI 0.73-0.80)/1,000 person-years. The three models containing age, abdominal pain, weight change and two laboratory biomarkers (ALT change/HgA1c, rate of ALT change/HgA1c, or rate of ALT change/rate of HgA1c change) had comparable discrimination and calibration measures (c-index: mean=0.77, SD=0.01-0.02; calibration test: p-value 0.2-0.4, SD 0.2-0.3). The VA validation cohort consisted of 2.6 million patients (mean age 66.1) with an 18-month incidence rate of 1.27 (1.23-1.30). A total of 606 patients were screened in the prospective pilot study at KPSC with 9 patients (1.5%) diagnosed with a pancreatic or biliary cancer.

CONCLUSIONS: Using widely available parameters in EHR, we developed a population-based parsimonious model for early detection of sporadic PC suitable for real-time application.

Keywords: risk prediction; pancreatic cancer; machine learning; general population; glycated hemoglobin; alanine transaminase; weight loss

INTRODUCTION

Pancreatic cancer is the third leading cause of cancer deaths with 48,220 estimated deaths in 2021 in the US.¹ Because of the lack of an early detection strategy, majority of patients (50-55%) have metastases at distant sites at the time of diagnosis.^{2,3} Once diagnosed, the average 5-year survival is only 10.8%.¹ Accounting for 90% of all pancreatic cancer cases, pancreatic ductal adenocarcinoma (PDAC) is by far the most common form of pancreatic cancer, and also the most lethal.

Due to the low incidence of pancreatic cancer in the general population (13.2 per 100,000 person-years),¹ widespread population-based screening is not currently recommended by the United States Preventative Services Task Force.⁴ Therefore, alternative approaches to early detection are needed in order to substantially impact the natural history of this disease and improve survival for patients.

The emergence of comprehensive EHR and maturation of machine learning offers an opportunity to enhance efforts in early detection in pancreatic cancer. To date, efforts to develop clinical prediction models in pancreatic cancer have focused on specific populations such as those with new-onset diabetes⁵⁻⁷ or within the confines of a case-control study.^{8,9} Efforts to target the high-risk patients in the general population are sparse.¹⁰ There is a critical need for novel risk stratification tools which are both sensitive and specific for rapid identification of patients at increased risk of developing pancreatic cancer.

The aim of the present study was to develop and validate a clinical prediction model for risk of PDAC across several large health systems. Specifically, we sought to apply machine-learning combined with a comprehensive approach to data in EHR to predict the risk of sporadic PDAC.

METHODS

Study Design and Setting

We conducted a retrospective cohort study utilizing multi-ethnic health plan enrollees of Kaiser Permanente Southern California (KPSC), a large integrated healthcare system that provides comprehensive healthcare services for >4.7 million enrollees across 15 medical centers and 235 medical offices. Model training and internal validation were conducted based on EHR data. The demographics and socioeconomic status of KPSC health plan enrollees are comparable to those of residents in the Southern California region.¹¹ The internally validated models were externally tested using EHR of Veterans Affairs (VA).¹² The study protocol was approved by the KPSC's institutional Review Board.

Study Population

Model training and internal validation: Patients 50-84 years of age and had ≥ 1 clinic-based visit (index visit) within a KPSC facility in 2008-2017 were identified. Patients who had history of pancreatic cancer, or not continuously enrolled in the KPSC health plan in the past 12 months (gaps 45 days or less were allowed) were excluded. The requirement of continuous enrollment allowed adequate data to define study variables. For patients with multiple qualifying index visits, we selected one randomly as the index visit. The corresponding visit date was referred to

as the index date (t_0). Follow-up started on t_0 and ended with the earliest of the following events: disenrollment from the health plan, end of the study (December 31, 2018), reached the maximum length of follow-up (18 months), non-PDAC related death, or PDAC diagnosis or death (outcome). A minimum of 30 days of follow-up is required.

Model testing: Veterans 50-84 years of age who had >1 outpatient visit (index visit) within a VA facility in 2008-2017 and another clinic-based visit within the 12 months prior to the index date were identified. Patients who had history of pancreatic cancer were excluded. The same follow-up rules mentioned above were applied to the VA cohort except for “disenrollment from the health plan”.

Outcome Identification

The study outcome was PDAC diagnosis or death with pancreatic cancer in the 18 months after the index date. For the KPSC cohort, PDAC was identified from the Cancer Registry by using the Tenth Revision of International Classification of Diseases, Clinical Modification (ICD-10-CM) code C25.x and histology codes (eTable 1). Pancreatic cancer deaths were derived from the linkage with the California State Death Master files and identified using ICD-10-CM codes C25.x.¹³ For the VA cohort, cases of PDAC were similarly identified through an internal VA Central Cancer Registry, and PDAC deaths identified through the VA Mortality Data Repository, which integrates vital status data from the National Death Index (NDI), VA, and DoD administrative records.

Patient Demographic and Clinical Features at Baseline

A complete list of extracted and derived features for the KPSC cohort is shown in eTable 2.

Except for demographic variables, values within each time interval (0-6 months, 7-12 months, 1-2 years and >2 years) were generated. Definitions of the derived variables were described in eTable3. Since the VA dataset was solely used for testing purposes, only limited number of features were extracted (Table 1).

Missing values were imputed¹⁴ if the frequency of missing was <60%. We used predictive mean matching method¹⁵ with k=5. Laboratory measures with ≥60% missingness or change/change rate measures with ≥80% missingness were not included in the model development process. Ten imputed datasets were generated.

Model Training, Validation and Testing

To overcome the limitations of regression-based models that are traditionally used for analysis of time-to-event data, we applied 'random survival forests' (RSF), a nonparametric machine learning method,¹⁶⁻¹⁸ to pre-select features and train/validate models. First, we iteratively preselected features based on the average minimum depth (eTable 4) and used those features to develop and validate risk prediction models based on 5-fold cross validation.¹⁹ Age was forced into the model. Preselected features were added incrementally to identify the feature that yielded the maximum improvement of c-index. This process continued until the c-index increased <0.005. Of the 50 models derived from the 50 training datasets (10 imputation

datasets x 5-fold cross validation), the three that appeared the most often were selected as the winning models.

Algorithms of the winning models were applied to the corresponding KPSC validation datasets. By design, the KPSC validation datasets did not include any observations of the KPSC training datasets from which the winning models were developed. One winning model was first directly applied to VA imputed datasets, and subsequently recalibrated to achieve better performance.

Performance Measures

The discriminative power for each of the winning models was evaluated by c-index, a concordance measure, averaged across all the relevant validation datasets for cohort members. Calibration was assessed by calibration plots with five risk groups (<50th, 50–74th, 75–89th, 90–94th, 95–100th percentiles).²⁰ Greenwood-Nam-D'Agostino (GND) calibration test was also performed to assess goodness-of-fit.

We estimated sensitivity, specificity, positive predictive value (PPV), and relative increase in risk in comparison to that of the entire cohort at various levels of risk thresholds. For this analysis we restricted the patients to those with complete follow up or developed PDAC in 18 months. The results were averaged across the validation datasets for each winning model.

Early Detection Model

To facilitate earlier detection of PDAC by ≥ 90 days, we also established a cohort which included patients identified in the main cohort who had ≥ 90 days of cancer-free follow up. The same model training and validation methods mentioned above were applied.

Prospective pilot study

We conducted a subsequent prospective study from February to December 2021 to evaluate feasibility of real-time implementation of the final prediction model. We aimed to assess the calibration of the model (frequency of observed vs. expected cancer). This study was approved as a separate protocol by the KPSC Institutional Review Board. We prospectively ran a final algorithm on a bi-weekly basis to identify patients aged 50-84 without a prior history of pancreatic cancer whose predicted risk of pancreatic cancer is $\geq 1\%$. All algorithm-identified patients were manually reviewed. Patients with findings suspicious for a pancreatic cancer on cross-sectional imaging obtained through routine clinical care up to 3 months prior to the risk identification date (the date when the algorithm was applied to EHR) were monitored. The diagnosis was based on either histology where available or presumptive clinical diagnosis as documented in patients' EHR.

Statistical Analysis

Analyses were performed using SAS (Version 9.4 for Unix; SAS Institute, Cary, NC) or R Version 3.6.0 (R Foundation, Vienna, Austria).

RESULTS

Characteristics of the study cohorts

1.8 million KPSC patients were eligible (eFigure 1), of which 53.3% were females, 45.3% were white, 29.8% were Hispanic, 9.5% were African American and 10.7% were Asian and Pacific Islanders (Table 1). The majority (60.7%) used commercial insurance, and slightly under one-third (31.8%) were on Medicare. On average, the KPSC patients were 61.6 years of age, with average membership length of 18.9 years. 35.7% of the patients were obese and additional 35.6% were overweight. Diabetes was common (about 20%), while acute and chronic pancreatitis were rare (<1%).

The 2.6 million eligible veterans were predominantly male (94.3%), white (67.9%) and African American (16.8%) and were older (66.1 years of age) than the KPSC cohort. Smoking, diabetes, acute and chronic pancreatitis were more prevalent in the VA cohort compared to those of the KPSC cohort. ALT and HbA1c at baseline appeared comparable between the two cohorts.

Incidence of PDAC

Table 2 displays the follow-up time in years, number, incidence rate (IR) of PDAC, and time to PDAC for all patients and for subgroups of patients defined by important features. 1,792 KPSC patients developed PDAC within 18 months of follow-up (IR=0.77, 95% CI 0.73-0.80/1,000 person-years (PY) (Table 2). A total of 4,582 patients in the VA cohort developed PDAC (IR 1.27 (1.23-1.30)). In the VA cohort, abdominal pain in the 6 months prior to t_0 increased the IR to 4.35 (4.01-4.70). Time to PDAC appeared to be longer for the VA cohort (median 233 days, IQR 116-370 days) compared to that of the KPSC cohort (median 205 days, IQR 91-358 days). The

distributions of cancer stage (I-IV) were comparable between the two cohorts (eTable 5), if the higher frequency of missingness in the VA cohort is ignored.

Model Training, Validation and Testing

The number and size of the KPSC training, KPSC validation and VA testing datasets are shown in eTable 6.

For the main cohort, the preselection process identified 29 potential predictors (eTable 4). Of the 50 training samples, the three winning models containing age, abdominal pain, weight change and two biomarkers (alanine transaminase (ALT) change/HbA1c, rate of ALT change/HbA1c, and rate of ALT change/rate of HbA1c change) appeared most often (Table 3). Internal validation based on KPSC validation datasets revealed comparable results among the top three models (c-index: mean 0.77 for all three models (M1-M3) and SD 0.01-0.02; calibration test: p-value 0.2-0.4 and SD 0.2-0.3). When M1 was directly applied to VA testing datasets, the mean c-index was 0.69 (SD 0.003) (data not shown); however, after the algorithm was recalibrated based on VA's datasets, the mean c-index based on 10 testing datasets was 0.71 (SD 0.002) (Table 3).

For the early detection cohort, the preselection process identified 32 potential predictors (eTable 4). The three best models (E1-E3, c-index: 0.74-0.77) contained the same features as those selected by M1-M3 except that abdominal pain was not chosen (Table 3). The calibration

test for model E3 was significant ($p=0.04$), indicating a lack of model fit. The recalibrated E1 model based on the VA testing datasets achieved a mean c-index of 0.68 (SD 0.003) (Table 3).

The hyperparameters used and the features selected by ≥ 5 out of 50 models for each cohort can be found in eTables 7 and 8, respectively.

Figure 1 displays the calibration plots for all six models (M1-M3, E1-E3). It appears that all fit well for the four out of five lower risk groups (i.e., risk < 95th percentile). However, for the highest risk group (risk ≥ 95 th percentile), M1-M3 properly estimated the risks at KPSC, while E1-E3 slightly overestimated the risks for KPSC patients, and M1, E1 slightly underestimated the risks for VA patients.

Sensitivity, specificity, PPV and fold increase in risk were comparable among the three winning models for both main cohort (M1-M3) and early detection cohort (E1-E3) (Table 4). The top 2.5% of the testing sample based on M1 experienced 1% risk of PDAC over 18 months, which was 7- to 8- fold higher than the baseline risk of PDAC in the KPSC cohort. Twenty % of the total PDAC cases occurring within 18 months were identified in this top 2.5% model-predicted high-risk group. Patients within the top 20% predicted risk of PDAC experienced 0.6% risk of PDAC over 18 months. This identified more than 50% of PDAC occurring within 18 months with a specificity of 80%. While sensitivities and fold increases in PDAC incidence rate were lower in the VA population, PPVs were higher for VA validation data compared to those of KPSC (Table 4).

Compared to the models developed based on the main cohort (M1-M3), the models developed based on early detection cohort (E1-E3) had slightly compromised sensitivity, PPV and fold increase in risk.

Model application/implementation

A total of 606 patients were identified by the model with predicted risk of $\geq 1\%$. Nine (1.5%) patients had abnormalities identified in the region of the pancreas with the following diagnoses/stages: PDAC stages 1b, II, III, IV; pancreatic lymphoma; pancreatic neuroendocrine tumor; ampullary carcinoma; cholangiocarcinoma (n=2).

To facilitate external application of the RSF-based prediction model (M1), we have developed a publicly available web-based tool (<https://pcriskdev.kp-scalresearch.org/>). A hypothetical 70-year-old male patient with hemoglobin A1c value of 7.5%, weight loss of 4 lbs. and ALT increase of 4 IU/L in one year has an estimated 18-month risk of PDAC 0.30%.

For demonstration, decision rules based on one of the trees built for M1 is displayed in eFigures 2 and 3 for the left and the right side of the decision tree, respectively.

DISCUSSION

We applied machine learning methods to EHR data to derive and validate clinical prediction models for sporadic pancreatic cancer across two large integrated healthcare systems. Despite

inclusion of >500 potential features in the candidate pool, the machine learning models incorporated traditional parameters including age, glycated hemoglobin, alanine aminotransferase, weight, and abdominal pain. The final models were both parsimonious (with only 4-5 predictors) and reasonably accurate in both internal and external validation. In addition, the model proved well-calibrated when applied prospectively for real-time identification of not only pancreatic but biliary cancers.

While there has been some progress in studying approaches to early detection in high-risk patients based on either family history or genetic susceptibility²¹ as well as those with specific conditions such as late-onset diabetes,⁶ limited data exist on identification of patients at risk for sporadic pancreatic cancer. This study presents a novel approach to risk stratification at the population-level based on dynamic parameters contained within structured data from EHR.

Although parameters included in the model are well-established parameters for PDAC,^{5,6,10} their selection using an unbiased, comprehensive data-driven approach helped ensure inclusion of the most relevant combination of parameters. A recently developed model to predict risk of pancreatic cancer among patients with late or new onset diabetes at age 50 or later similarly identified increasing age, weight loss and change in blood glucose as key parameters for determining risk of pancreatic cancer in this patient population.⁵ The current model extends the concept of model-based risk prediction to a much broader population while maintaining reasonably high levels of discriminative accuracy for prediction of pancreatic cancer.

External validation is key to assessing model performance. In a review of 127 prediction models, Siontis et. al found 32 (25%) had at least one external validation.²² AUC estimates significantly decreased during external validation vs. the derivation study with a median AUC reduction 0.05.²² In the current study, the c-index declined by 0.08 (or 10%) when model M1 was directly transported and about 0.06 (or 8%) and 0.09 (or 12%) after models M1 and E1 were recalibrated, respectively. Although the comparison should be interpreted cautiously, the larger reduction observed in the current study could be attributable to multiple factors. First, a higher frequency of PDAC cases in the VA cohort were pancreatic cancer deaths identified through mortality records compared to the KPSC cohort. Second, given the differences in age and sex between KPSC and VA populations, a higher incidence rate of PDAC in the VA dataset compared to that of KP's was observed as expected. This could have impacted model accuracy especially for the models without recalibration.

Strengths of the current study included a comprehensive, data-driven approach to model development, use of structured data elements and external validation in a separate healthcare system with distinct patient population. The present study also extended model development to assess feasibility of implementing the model through application in a prospective pilot study that demonstrated ability to identify pancreaticobiliary cancers in real-time.

The present study also had several limitations. First, several parameters identified in the prediction models (abdominal pain, abnormal ALT) are often associated with advanced stage pancreatic cancer. However, several early-stage pancreatic cancer cases were detected by the

algorithm in the prospective feasibility study. Nevertheless, an ongoing concern relates to the timing with respect to clinical diagnosis. The 30-day cancer-free period used in the present model is likely insufficient to provide a reasonable window of opportunity for intervention to impact the disease course. To address this concern, we also developed an early detection model that restricted the study population to patients with ≥ 90 days cancer-free follow-up from the index date. Further testing of this restricted model for early detection of pancreatic cancer is the subject of an ongoing single-arm prospective interventional study (NCT04883450). Despite the reasonably high performance in terms of discriminative ability, the absolute risk in the highest risk category (top 2.5%) approached 1% over 18-months. This level of risk is likely below the threshold for cost-effective screening based on currently available testing.²³ Second, of the 1479 and 4,582 events identified in the KPSC and VA cohorts, respectively, 300 and 2,564 events were captured by data sources other than Cancer Registry. An evaluation based on the KPSC Cancer Registry of the same time window showed that about 90% of pancreatic cancer cases were PDAC. Third, to estimate sensitivity, specificity, PPV and fold of risk increase, we relied on a subset of patients (~70% and ~80% of the total patients in the KPSC and VA cohorts, respectively) with complete follow up unless they died of pancreatic cancer. This restriction over-estimated the risk of PDAC, because the patients who were excluded from this analysis were at-risk for some periods of time. Finally, several cancers identified during prospective pilot testing of the algorithm were biliary in origin given similarities in anatomic location and clinical presentation.

In conclusion, we developed a parsimonious clinical risk prediction model for sporadic pancreatic cancer in a large, diverse integrated health system and subsequently applied the model in a separate health system. We also evaluated the model prospectively for real-time application. The model identified five key factors in determining risk of pancreatic cancer. Findings from the present study provide a potential framework for a systematic approach to targeted screening for pancreatic cancer based on automated analysis of data in EHR.

References

1. Cancer Stat Facts: Pancreatic Cancer. *SEER* <https://seer.cancer.gov/statfacts/html/pancreas.html>. Accessed July 13, 2021.
2. Stathis A, Moore MJ. Advanced pancreatic carcinoma: current treatment and future challenges. *Nat Rev Clin Oncol*. 2010;7(3):163-172.
3. Stokes JB, Nolan NJ, Stelow EB, et al. Preoperative capecitabine and concurrent radiation for borderline resectable pancreatic cancer. *Ann Surg Oncol*. 2011;18(3):619-627.
4. US Preventive Services Task Force, Owens DK, Davidson KW, et al. Screening for Pancreatic Cancer: US Preventive Services Task Force Reaffirmation Recommendation Statement. *JAMA*. 2019;322(5):438-444.
5. Sharma A, Kandlakunta H, Nagpal SJS, et al. Model to Determine Risk of Pancreatic Cancer in Patients With New-Onset Diabetes. *Gastroenterology*. 2018;155(3):730-739 e733.
6. Boursi B, Finkelman B, Giantonio BJ, et al. A Clinical Prediction Model to Assess Risk for Pancreatic Cancer Among Patients With New-Onset Diabetes. *Gastroenterology*. 2017;152(4):840-850 e843.
7. Chen W, Butler RK, Lustigova E, Chari ST, Wu BU. Validation of the Enriching New-Onset Diabetes for Pancreatic Cancer Model in a Diverse and Integrated Healthcare Setting. *Dig Dis Sci*. 2021;66(1):78-87.
8. Kim J, Yuan C, Babic A, et al. Genetic and Circulating Biomarker Data Improve Risk Prediction for Pancreatic Cancer in the General Population. *Cancer Epidemiol Biomarkers Prev*. 2020;29(5):999-1008.
9. Klein AP, Lindstrom S, Mendelsohn JB, et al. An absolute risk model to identify individuals at elevated risk for pancreatic cancer in the general population. *PLoS One*. 2013;8(9):e72311.
10. Yu A, Woo SM, Joo J, et al. Development and Validation of a Prediction Model to Estimate Individual Risk of Pancreatic Cancer. *PLoS One*. 2016;11(1):e0146473.
11. Koebnick C, Langer-Gould AM, Gould MK, et al. Sociodemographic characteristics of members of a large, integrated health care system: comparison with US Census Bureau data. *Perm J*. 2012;16(3):37-41.
12. Fihn SD, Francis J, Clancy C, et al. Insights from advanced analytics at the Veterans Health Administration. *Health Aff (Millwood)*. 2014;33(7):1203-1211.
13. Chen W, Yao J, Liang Z, et al. Temporal Trends in Mortality Rates among Kaiser Permanente Southern California Health Plan Enrollees, 2001-2016. *Perm J*. 2019;23.
14. Wright MN, Ziegler A. ranger: A Fast Implementation of Random Forests for High Dimensional Data in C++ and R. *J Stat Softw*. 2017;77(1).
15. Little RJA. Missing-Data Adjustments in Large Surveys. *J Bus Econ Stat*. 1988;6(3):287-296.
16. Ishwaran H, Kogalur UB, Blackstone EH, Lauer MS. Random survival forests. *Ann Appl Stat*. 2008(3):841-860.

17. Dietrich S, Floegel A, Troll M, et al. Random Survival Forest in practice: a method for modelling complex metabolomics data in time to event analysis. *Int J Epidemiol.* 2016;45(5):1406-1420.
18. Ishwaran H KU. randomForestSRC: Fast Unified Random Forests for Survival, Regression, and Classification (RF-SRC). <http://web.ccs.miami.edu/~hishwaran/>. Accessed July 8, 2021.
19. Stone M. Cross-Validatory Choice and Assessment of Statistical Predictions. *J R Stat Soc Series B Stat Methodol.* 1974;36(2):111-147.
20. Demler OV, Paynter NP, Cook NR. Tests of calibration and goodness-of-fit in the survival setting. *Stat Med.* 2015;34(10):1659-1680.
21. Goggins M, Overbeek KA, Brand R, et al. Management of patients with increased risk for familial pancreatic cancer: updated recommendations from the International Cancer of the Pancreas Screening (CAPS) Consortium. *Gut.* 2020;69(1):7-17.
22. Siontis GC, Tzoulaki I, Castaldi PJ, Ioannidis JP. External validation of new risk prediction models is infrequent and reveals worse prognostic discrimination. *J Clin Epidemiol.* 2015;68(1):25-34.
23. Schwartz NRM, Matrisian LM, Shrader EE, Feng Z, Chari S, Roth JA. Potential Cost-Effectiveness of Risk-Based Pancreatic Cancer Screening in Patients With New-Onset Diabetes. *J Natl Compr Canc Netw.* 2021:1-9.

Figure Legends

Figure 1 – Calibration plots of winning models. x-axis: predicted; y-axis: observed. The five clusters represent the five risk groups defined by the ranges of predicted risks: <50th, 50–74th, 75–89th, 90–94th, 95–100th percentiles. Within each cluster, there are multiple dots representing the pairs of predicted and observed risks, calculated based on the corresponding validation datasets.

eFigure 1 – Consort Diagram (KPSC and VA main cohorts)

eFigure 2 – One of the decision trees for KPSC M1 (left side)

eFigure 3 – One of the decision trees for KPSC M1 (right side)

Table 1. Characteristics of study subjects at baseline, n (%) unless otherwise stated.

Demographics and Lifestyle Characteristics	Kaiser Permanente Southern California N=1,801,931		Veterans Affairs N=2,690,895	
Age, mean (SD)	61.6 (9.4)		66.1 (9.1)	
Female	960266 (53.3)		153630 (5.7)	
Race/Ethnicity				
Non-Hispanic White	815773 (45.3)		1828095 (67.9)	
Non-Hispanic Black	171424 (9.5)		451523 (16.8)	
Hispanic	536079 (29.8)		148690 (5.5)	
Asian and Pacific Islander	192179 (10.7)		38536 (1.4)	
Multiple/Other/Unknown	86476 (4.8)		224051 (8.3)	
Medical Insurance (one or more)				
Commercial	1094182 (60.7)			
Medicare	572452 (31.8)			
Medi-CAL/Other State Programs	64150 (3.6)			
Private Pay	483042 (26.8)			
Years Since First enrollment, mean (SD)	18.9 (13.5)			
Family History of Pancreatic Cancer	25386 (1.4)			
Tobacco Use				
Ever	700429 (38.9)		1914180 (71.1)	
Never	1101502 (61.1)		776715 (28.9)	
Weight Defined by BMI (kg/m ²)				
Underweight (<18.5)	19095 (1.1)			
Normal Weight (18.5-24.9)	429010 (23.8)			
Overweight (25-29.9)	640644 (35.6)			
Obese (30+)	642905 (35.7)			
Unknown	70277 (3.9)			
Weight Change in 1 Year in lb., median (IQR) ^a	-0.2 (-4.8, 3.5)		-0.3 (-5.8, 4.8)	
Lab Tests	Value in Prior 6 Months		Change in 1 Year	
	N	Median (IQR)	N	Median (IQR)
ALT, IU/L	920993	22.0 (17.0, 30.0)	479881	0.0 (-5.0, 4.0)
HgA1c, %	744601	6.2 (5.8, 7.1)	409975	0.0 (-0.3, 0.3)
ALP, IU/L	309460	70.0 (57.0, 87.0)	96562	0.0 (-9.0, 10.0)
Total Bilirubin, mg/dL	303803	0.7 (0.5, 0.9)	94098	0.0 (-0.2, 0.2)
HGB for Males, g/dL	437416	14.5 (13.4, 15.4)	202538	-0.1 (-0.8, 0.5)
HGB for Females, g/dL	515110	13.2 (12.4, 14.0)	264229	-0.1 (-0.6, 0.5)
HCT, L/L	952548	41.0 (38.0, 43.7)	466785	-0.3 (-2.1, 1.5)

RBC, million/mm ³	934454	4.5 (4.2, 4.9)	449246	0.0 (-0.2, 0.2)				
Sodium, mEq/L	952253	139.0 (137.0, 141.0)	496334	0.0 (-2.0, 2.0)				
Total Cholesterol, mg/dL	964899	184.0 (156.0, 215.0)	506608	-4.0 (-22.0, 13.0)				
Platelets, count/L	934239	232.0 (194.0, 277.0)	449030	-3.0 (-25.0, 19.0)				
Medical Conditions	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>
Gallstone Disorders	20085 (1.1)	13184 (0.7)	20183 (1.1)	104529 (5.8)				
Acute Pancreatitis	3256 (0.2)	2065 (0.1)	3199 (0.2)	17975 (1.0)	9510 (0.4)	5646 (0.2)	8859 (0.3)	18209 (0.7)
Chronic Pancreatitis	1550 (0.1)	1182 (0.1)	1475 (0.1)	3738 (0.2)	7749 (0.3)	5125 (0.2)	6707 (0.2)	9719 (0.4)
Benign Pancreatic Disease	2019 (0.1)	1398 (0.1)	1711 (0.1)	2780 (0.2)				
Biliary Tract Disease	26199 (1.5)	20974 (1.2)	25512 (1.4)	38538 (2.1)				
Depression	189508 (10.5)	150514 (8.4)	190868 (10.6)	317686 (17.6)				
Diabetes	376022 (20.9)	325789 (18.1)	335965 (18.6)	327408 (18.2)	842,311 (31.3)	775,964 (28.8)	831,566 (30.9)	735,251 (27.3)
Medical Procedures	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>
Abdominal/Chest CT	121427 (6.7)	82841 (4.6)	119308 (6.6)	304085 (16.9)				
Abdominal/Chest MRI	1861 (0.1)	1366 (0.1)	2226 (0.1)	7057 (0.4)				
Abdominal/Chest Ultrasound	70721 (3.9)	55004 (3.1)	93179 (5.2)	333051 (18.5)				
Any Abdominal Surgery	11447 (0.6)	7360 (0.4)	11559 (0.6)	77283 (4.3)				
Surgical Procedures on esophagus	18492 (1.0)	13226 (0.7)	22164 (1.2)	125701 (7.0)				
Upper GI Endoscopy	37108 (2.1)	27232 (1.5)	43386 (2.4)	160899 (8.9)				
Colonoscopy	95156 (5.3)	74199 (4.1)	125372 (7.0)	417049 (23.1)				
Medications	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>
Pancreatic Enzyme	1246 (0.07)	1105 (0.06)	1314 (0.07)	2437 (0.1)				
Antidiabetic Medications – Insulin	88277 (4.9)	79163 (4.4)	78660 (4.4)	78323 (4.4)				
Antidiabetic Medications – Non-Insulin	158667 (8.8)	148703 (8.3)	155764 (8.6)	182396 (10.1)				
GI-Related Signs/Symptoms	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>	<i>0-6 m Prior</i>	<i>7-12 m Prior</i>	<i>13-24 m Prior</i>	<i>24+ m Prior</i>
Abdominal Pain	133380 (7.4)	92422 (5.1)	142470 (7.9)	483652 (26.8)	107586 (4.0)	68717 (2.6)	108170 (4.0)	237763 (8.8)
Chest Pain	114152 (6.3)	79399 (4.4)	128549 (7.1)	483953 (26.7)				
Constipation	60032 (3.3)	41935 (2.3)	64439 (3.6)	193331 (10.7)				
Diarrhea	44025 (2.4)	32187 (1.8)	52487 (2.9)	202067 (11.2)				
Itching	46314 (2.6)	33730 (1.9)	57062 (3.2)	207679 (11.5)				
Malaise or Fatigue	104298 (5.8)	71136 (3.9)	111646 (6.2)	362901 (20.1)				
Melena	11089 (0.6)	7433 (0.4)	13235 (0.7)	66321 (3.7)				
Nausea or Vomiting	51787 (2.9)	36348 (2.0)	57767 (3.2)	215982 (12.0)				

Abbreviations: ALP, alkaline phosphatase; ALT, alanine transaminase; BMI, body mass index; CT, computerized tomography; GI, gastrointestinal; HCT, hematocrit; HbA1c: hemoglobin A1c; HGB, hemoglobin; IQR, interquartile range; MRI, magnetic resonance imaging; RBC, red blood cell; SD, standard deviation.

^a 1 lb = 0.45 kg

Table 2. Total follow-up (f/u) time, number, and incidence rate of PDAC per 1,000 person-years (PY) and 95% CI.

	Kaiser Permanente Southern California (KPSC)				Veterans Affairs (VA)			
	Total f/u Time (years)	No. of PDAC	Incidence Rate of PDAC/ 1000 PY (95% CI)	Days to PDAC (median, IQR)	Total f/u Time (years)	No. of PDAC	Incidence Rate of PDAC/ 1000 PY (95% CI)	Days to PDAC (median, IQR)
All	2331767	1792	0.77 (0.73, 0.80)	205 (91, 358)	3614215	4582	1.27 (1.23, 1.30)	233 (116, 370)
Age Group in years								
50-59	1133128	350	0.31 (0.28, 0.34)	198 (84, 357)	989814	679	0.69 (0.64, 0.74)	212 (98, 352)
60-69	711410	624	0.88 (0.81, 0.95)	197 (91, 355)	1630532	2026	1.24 (1.19, 1.30)	239 (122, 374)
70-79	369643	604	1.63 (1.51, 1.77)	219 (91, 362)	677576	1304	1.92 (1.82, 2.03)	237 (113, 369)
80-84	117585	214	1.82 (1.59, 2.08)	213 (112, 353)	316293	573	1.81 (1.67, 1.96)	235 (116, 367)
Sex								
Female	1246235	864	0.69 (0.65, 0.74)	219 (97, 370)	213181	101	0.47 (0.39, 0.57)	248 (104, 398)
Male	1085520	928	0.85 (0.80, 0.91)	220 (88, 350)	3401034	4481	1.32 (1.28, 1.36)	233 (116, 369)
Race/ Ethnicity								
Non-Hispanic White	1065194	934	0.88 (0.82, 0.93)	206 (90, 362)	2456059	2986	1.22 (1.17, 1.26)	238 (119, 370)
Non-Hispanic Black	226546	262	1.16 (1.02, 1.30)	208 (100, 356)	612641	692	1.13 (1.05, 1.22)	220.5 (100.5, 377)
Asian/Pacific Islander	255168	147	0.58 (0.49, 0.67)	216 (114, 367)	52570	48	0.91 (0.68, 1.20)	192 (118.5, 373.5)
Hispanic	692443	410	0.59 (0.53, 0.65)	201 (87, 345)	202263	211	1.04 (0.91, 1.19)	239 (121, 369)
Unknown ^a					290682	645	2.22 (2.05, 2.40)	220 (115, 357)
ALT change in 1 year in IU/L								
≤ -5	156757	174	1.11 (0.95, 1.28)	216 (92, 360)	248360	359	1.45 (1.30, 1.60)	247 (123, 367)
(-5, 5]	340273	330	0.97 (0.87, 1.08)	232 (107, 392)	434050	523	1.20 (1.10, 1.31)	246.5 (132.5, 367.5)
> 5	123457	165	1.34 (1.14, 1.55)	149 (63, 303)	187824	322	1.71 (1.53, 1.91)	184 (86, 302)
Unknown	1711281	1123	0.66 (0.61, 0.70)	203 (94, 355)	2743981	3378	1.23 (1.19, 1.27)	234 (116, 373)
Rate of ALT Change in 1 Year								
≤ -0.01	180995	205	1.13 (0.99, 1.30)	210 (91, 358)	281222	402	1.43 (1.29, 1.57)	247 (126, 373)
(-0.01, 0.01]	272305	255	0.94 (0.83, 1.06)	234 (114, 384)	343020	420	1.22 (1.11, 1.35)	244.5 (132.5, 367.5)
> 0.01	167186	209	1.25 (1.09, 1.43)	166 (65, 330)	245991	382	1.55 (1.40, 1.71)	191 (86, 310)
Unknown	1711281	1123	0.66 (0.62, 0.70)	203 (94, 355)	2743981	3378	1.23 (1.19, 1.27)	234 (116, 373)
HgA1c Value Prior to Index date								
<6.5%	586772	326	0.55 (0.50, 0.62)	220 (90, 369)	1132107	1094	0.97 (0.91, 1.02)	241 (118, 373)
6.5-6.9%	124249	139	1.12 (0.94, 1.32)	190 (74, 360)	228430	366	1.60 (1.44, 1.77)	234 (118, 367)
7.0-7.4%	80113	122	1.52 (1.27, 1.81)	159 (88, 310)	157876	289	1.83 (1.63, 2.05)	221 (112, 366)
≥7.5%	179700	305	1.70 (1.51, 1.89)	200 (90, 341)	399057	969	2.43 (2.28, 2.58)	216 (102, 354)
Unknown	1360933	900	0.66 (0.62, 0.71)	212 (96, 365)	1696745	1864	1.10 (1.05, 1.15)	239 (121, 378)
HgA1c Change in 1 Year in %								
< -0.3	118111	143	1.21 (1.02, 1.42)	185 (88, 343)	287080	479	1.67 (1.52, 1.82)	227 (115.5, 371)

[-0.3, 0.3]	298587	204	0.68 (0.59, 0.78)	238 (100, 372)	625990	637	1.02 (0.94, 1.10)	251 (130, 388)
> 0.3	120115	211	1.76 (1.53, 2.01)	190 (90, 329)	291,676	660	2.26 (2.10, 2.44)	220 (103, 357)
Unknown	1794953	1234	0.69 (0.65, 0.73)	205 (90, 358)	2409469	2806	1.16 (1.12, 1.21)	233 (117, 370)
Rate of HgA1c Change in 1 Year								
< -0.0008	126922	146	1.15 (0.97, 1.35)	184 (88, 343)	322572	519	1.61 (1.47, 1.75)	227 (115, 371)
[-0.0008, 0.0008]	280155	181	0.64 (0.56, 0.75)	224 (98, 371)	554791	535	0.96 (0.89, 1.05)	249 (129, 378)
> 0.0008	129736	231	1.78 (1.56, 2.02)	200 (92, 343)	327480	722	2.21 (2.05, 2.37)	225.5 (105, 359)
Unknown	1794953	1234	0.69 (0.65, 0.73)	205 (90, 358)	2409469	2806	1.16 (1.12, 1.21)	233 (117, 370)
Rate of Weight Change in 1 Year								
< -0.02	503012	683	1.35 (1.26, 1.46)	160 (68, 322)	869902	2049	2.36 (2.26, 2.46)	185 (86, 313)
[-0.02, 0.02]	1033671	703	0.68 (0.63, 0.73)	238 (124, 387)	1437632	1403	0.98 (0.93, 1.03)	286 (156, 409)
> 0.02	393142	209	0.53 (0.46, 0.61)	238 (114, 358)	750864	621	0.83 (0.76, 0.89)	281 (145, 408)
Unknown	401941	197	0.49 (0.43, 0.56)	202 (90, 364)	556817	509	0.91 (0.84, 0.996)	244 (133, 387)
Abdominal Pain in Prior 6 mos ^b								
Yes	168579	322	1.91 (1.71, 2.13)	116 (54, 256)	139459	606	4.35 (4.01, 4.70)	136 (67, 252)
No	134525	139	1.03 (0.87, 1.22)	199 (110, 349)	3474756	3976	1.14 (1.11, 1.18)	250.5 (128, 382.5)
Unknown	2028662	1331	0.66 (0.62, 0.69)	230 (110, 377)				

Abbreviations: ALT, alanine transaminase; CI, confidence interval; F/U, follow-up; HgA1c, hemoglobin A1c; IQR, interquartile range; PDAC, pancreatic ductal adenocarcinoma; PY, person-years.

^a Not estimated for KPSC dataset due to the small number of events (39) in this group.

^b For VA data, "Unknown" was interpreted as "No" due to the inability to distinguish between the two.

Table 3. Predictors selected in top 3 models^a developed based on the main cohort and restricted cohort, and model performance during internal validation and external testing.

Model Name	Predictors							Training KPSC	Validation KPSC			Testing ^b VA
	Age	ALT		HgA1c		Abdomi- nal Pain	Weight Change in 1 yr	No. of times selected out of 50 training samples	c-index Mean (SD)	χ2 Mean (SD)	p-value Mean (SD)	c-index Mean (SD)
		Change	Rate of Change in 1 yr	HgA1c	Rate of Change in 1 yr							
Main Cohort												
M1	X	X		X		X	X	11	0.77 (0.02)	5.9 (4.1)	0.4 (0.3)	0.71 (0.002)
M2	X		X	X		X	X	11	0.77 (0.02)	8.5 (5.3)	0.2 (0.3)	NA
M3	X		X		X	X	X	9	0.77 (0.01)	6.7 (2.1)	0.2 (0.2)	NA
Early Detection Cohort												
E1	X	X		X		Not Selected	X	5	0.77 (0.04)	7.3 (4.9)	0.3 (0.3)	0.68 (0.003)
E2	X		X	X			X	6	0.74 (0.02)	7.4 (4.2)	0.2 (0.2)	NA
E3	X		X		X		X	5	0.76 (0.03)	10.3 (2.0)	0.04 (0.02)	NA

Abbreviations: ALT, alanine transaminase; C-index, concordance index; HbA1c, hemoglobin A1c; KPSC, Kaiser Permanente Southern California; NA, not applicable; SD, standard deviation; VA, Veterans Affairs

^a Top three models were selected based on the number of times each model was selected out of 50 training samples.

^b Models were recalibrated using the exact features selected by Model M1 and Model E1, respectively.

Table 4. Percent of patients^a whose risk was among the top 20%, 15%, 10%, 5%, and 2.5%, sensitivity, specificity, positive predictive value (PPV), and risk fold increase for each of the winning models based on KPSC validation datasets and one of the winning models based on VA validation dataset.

	KPSC															VA				
	High-Risk Patients					High-Risk Patients					High-Risk Patients					High-Risk Patients				
	Top 20%	Top 15%	Top 10%	Top 20%	Top 2.5%	Top 20%	Top 15%	Top 10%	Top 5%	Top 2.5%	Top 20%	Top 15%	Top 10%	Top 5%	Top 2.5%	Top 20%	Top 15%	Top 10%	Top 5%	Top 2.5%
Main Cohort	Model M1					Model M2					Model M3					Model M1				
N ^b	50952	37454	24959	424761	6249	49746	37512	24949	12572	6244	49934	37545	24850	12468	6238	424761	318408	212392	106118	53528
Sensitivity (%)	56.6	48.8	39.0	51.7	19.5	55.5	47.5	38.8	27.1	18.5	56.6	48.9	40.0	28.0	19.5	51.7	45.5	37.7	27.2	19.4
Specificity (%)	79.6	85.1	90.0	80.2	97.5	80.1	85.0	90.1	95.0	97.5	80.1	85.0	90.1	95.0	97.5	80.2	85.1	90.1	95.1	97.5
PPV (%)	0.4	0.5	0.6	0.6	1.1	0.4	0.4	0.5	0.8	1.0	0.4	0.5	0.6	0.8	1.1	0.6	0.7	0.8	1.2	1.7
Fold Increase in Risk ^c	2.8	3.3	3.9	2.6	7.9	2.8	3.2	3.9	5.4	7.4	2.8	3.3	4.0	5.6	7.9	2.6	3.0	3.7	5.4	7.7
Early Detection Cohort	Model E1					Model E2					Model E3					Model E1				
N ^b	49944	37462	24974	424521	6244	49966	37475	24984	12492	6246	49917	37439	24960	12480	6240	424521	318131	212172	106087	53047
Sensitivity (%)	55.1	44.1	35.1	49.6	14.5	50.7	42.2	33.2	20.3	12.6	52.3	44.2	33.6	22.3	13.1	49.6	42.8	34.5	23.6	15.5
Specificity (%)	80.0	85.0	90.0	80.1	97.5	80.9	80.0	85.0	90.0	95.0	80.0	85.0	90.0	95.0	97.5	80.1	85.1	89.8	95.0	97.5
PPV (%)	0.3	0.3	0.4	0.4	0.6	0.3	0.3	0.4	0.4	0.5	0.3	0.3	0.4	0.5	0.6	0.4	0.6	0.6	0.8	1.1
Fold Increase in Risk ^c	2.6	2.9	3.5	2.5	5.8	2.5	2.8	3.3	4.1	5.0	2.6	2.9	3.4	4.5	5.2	2.5	2.9	3.5	4.7	6.2

Abbreviations: KPSC, Kaiser Permanente Southern California; PPV, positive predictive value; VA, Veterans Affairs.

^a Estimated in patients with complete 18 months follow up or those who developed PDAC in 18 months.

^b Number of eligible patients whose risk was above each risk threshold.

^c Compared with the incidence rate in the entire cohort.

Model M1: Main cohort; age, weight change, abdominal pain, ALT change, HgA1c; estimated based on 11 validation samples for KPSC and 10 imputed testing datasets for VA.

Model M2: Main cohort; age, weight change, abdominal pain, ALT change rate, HgA1c; estimated based on 11 validation samples.

Model M3: Main cohort; age, weight change, abdominal pain, ALT change rate, HgA1c change rate; estimated based on 9 validation samples.

Model E1: Early detection cohort; age, weight change, ALT change, HgA1c; estimated based on 5 validation samples for KPSC and 10 imputed testing datasets for VA.

Model E2: Early detection cohort; age, weight change, ALT change rate, HgA1c, estimated based on 6 validation samples.

Model E3: Early detection cohort; age, weight change, ALT change rate, HgA1c change rate, estimated based on 5 validation samples.