Examining the Association between the Gastrointestinal Microbiota and Gulf War Illness: A Prospective Cohort Study

Ashley Kates^{1,2*}, Julie Keating¹, Kelsey Baubie¹, Nathan Putman-Buehler², Lauren Watson⁴, Jared Godfrey^{1,2}, Courtney L. Deblois^{1,5,6}, Garret Suen⁵, Dane B. Cook^{1,7}, David Rabago⁸, Ronald Gangnon^{9,10}, Nasia Safdar^{1,2}

¹Research, William S. Middleton Memorial Veterans Hospital, Madison, Wisconsin, USA

²Department of Medicine, School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin, USA

³Department of Biochemistry, College of Agriculture & Life Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

⁴SSM Health, St. Mary's Hospital, Madison, Wisconsin, USA

⁵Department of Bacteriology, College of Agriculture and Life Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

⁶Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, Wisconsin, USA

⁷Department of Kinesiology, University of Wisconsin-Madison, Madison, Wisconsin, USA

⁸Department of Family and Community Medicine, College of Medicine, Penn State University, Hershey, Pennsylvania, USA

⁹Department of Biostatistics and Medical Informatics, School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin, USA

¹⁰Department of Population Health Sciences, School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin, USA

* Corresponding author

email: akates@medicine.wisc.edu

Abstract

Gulf War Illness (GWI) affects 25-35% of the 1991 Gulf War Veteran population. Patients with GWI experience pain, fatigue, cognitive impairments, gastrointestinal dysfunction, skin disorders, and respiratory issues. In longitudinal studies, many patients with GWI have shown little to no improvement in symptoms since diagnosis. The gut microbiome and diet play an important role in human health and disease, and preliminary studies suggest it may play a role in GWI. To examine the relationship between the gut microbiota, diet, and GWI, we conducted an eight-week prospective cohort study collecting stool samples, medications, health history, and dietary data. Sixty-nine participants were enrolled into the study, 36 of which met the case definition for GWI. The gut microbiota of participants, determined by 16S rRNA sequencing of stool samples, was stable over the duration of the study and showed no within person (alpha diversity) differences. Between group analyses (beta diversity) identified statistically significant different between those with and without GWI. Several taxonomic lineages were identified as differentially abundant between those with and without GWI (n=9) including a greater abundance of Lachnospiraceae and Ruminococcaceae in those without GWI. Additionally, there were taxonomic differences between those with high and low HEI scores including a greater abundance of Ruminococcaceae in those with higher HEI scores. This longitudinal cohort study of GWVs found that participants with GWI had significantly different microbiomes from those without GWI. Further studies are needed to determine the role these differences may play in the development and treatment of GWI.

Introduction

Gulf War Illness (GWI) is a devastating chronic multi-symptom syndrome impacting 25-35% of the 700,000 coalition troops deployed during the 1990-91 Gulf War.(1)(2) Following their

return from the Persian Gulf, Veterans experienced a unique pattern of symptoms across multiple physiologic domains with no identifiable cause. Over the last 30 years, a number of theories as to the cause of GWI have arisen with consensus being that troops were exposed to environmental contaminants – most likely pyridostigmine bromide (PB) or chemical nerve agents – during their deployment during Operations Desert Shield and Desert Storm.(3) Individuals with GWI experience musculoskeletal pain, fatigue, neurological symptoms (e.g., memory and cognitive issues, difficulty sleeping, depression, dizziness, headaches), rashes and other skin disorders, respiratory symptoms (e.g., persistent cough and difficulty breathing), as well as gastrointestinal issues (e.g., diarrhea, nausea, vomiting, and Irritable Bowel Syndrome).(4) Few effective treatments have been found, and patients have experienced little to no improvement in their symptoms over time.(5) Additionally, these patients are aging faster and developing additional comorbidities such as heart disease, diabetes, and arthritis at younger ages than the general population.(6)

While several case definitions for GWI exist, the Kansas case definition is most frequently reported; it is also one of the recommended case definitions by the Institute of Medicine.(4)(7) According to the Kansas case definition, patients must have moderately severe symptoms in three or more of the six symptom domains (chronic fatigue, neurological symptoms, joint or muscle pain, gastrointestinal disturbance, respiratory symptoms, and skin problems) for at least 6 months that are not explained by any other medical or psychiatric condition. Using these criteria, roughly one-third of GW Veterans (GWVs) meet the definition of GWI.(4)

Research to date primarily focuses on neurologic symptoms of GWI; however, gut dysfunction is also a commonly reported symptom of GWI patients.(8) Veterans with GWI related gut dysfunction experience pain, nausea, vomiting, bloating, and diarrhea similar to Irritable Bowel Syndrome (IBS) and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). Disturbances to the gut microbiota may play a role in disease pathogenesis of both

ME/CFS and IBS.(9)(10) Given the similarities between these syndromes and GWI, researchers have begun to consider the role of the gut microbiome role in GWI; preliminary studies show significantly different gut microbiomes between healthy controls and those with GWI in both animals and humans. Mice treated with PB and corticosterone had over 100 different operational taxonomic units (OTUs) than controls ($p=0.005$).(11) Similarly, mice treated with PB and permethrin had reduced alpha diversity ($p < 0.001$) compared to controls; high fat diets further reduced alpha diversity while returns to normal feed provided partial recovery, indicating the potential role of diet in GWI symptom exacerbation.(12) A small pilot study of GWVs with GWI found those with GWI and gut symptoms had significantly different microbiomes compared to both those with GWI and no gut symptoms and healthy controls including higher abundances of Bacteroidetes, Actinobacteria, Erysipelotrichaceae, and Bifidobacteriaceae.(13)

We undertook a prospective cohort study to assess the relationship between GWI and the gut microbiome in GWVs with GWI and healthy controls. We hypothesized those with GWI would have less diverse microbiomes and higher Bacteroidetes abundance than healthy controls.

Methods

We conducted an eight-week prospective cohort study assessing the gut microbiomes of GWVs. A detailed description of the study methodology has previously been published.(14)

Study population, recruitment, and consent

Veterans meeting deployment criteria were identified from VA databases and were recruited via mailed invitation letters. Following invitation letters, GWVs interested in participating completed a phone screening to determine eligibility. Inclusion and exclusion criteria can be found in Table 1. Written informed consent was obtained at the enrollment visit

by a trained team member prior to initiating additional study procedures. Participants received \$50 for completing the enrollment visit and an additional \$50 for the close-out visit. This study and all associated protocols were approved by the University of Wisconsin Madison Health Sciences Institutional Review Board (ID No. 2017-1212) and the Madison VA Research and Development Committee.

Table 1. Study inclusion and exclusion criteria

Inclusion Criteria
Age 43-75 years
Deployed to the Persian Gulf as part of Operation Desert Shield and/or Operation Desert Storm during the 1990-1991 Gulf War
Exclusion Criteria
Diagnosed with a neurologic or musculoskeletal condition (lupus, multiple sclerosis, Parkinson's disease, or amyotrophic lateral sclerosis)
Diagnosed with inflammatory bowel disease
One or more of the GWI Kansas case definition exclusionary conditions: schizophrenia, active cancer treatment, or presence of cognitive/physical impairments following a stroke
Unstable psychiatric illness (defined as hospitalization within the previous year for depression, bipolar disorder, or post-traumatic stress disorder)
Active/unstable illness (defined as hospitalization within the last 5 years for diabetes, heart disease, rheumatoid arthritis, seizure disorder, kidney or liver disease)
Chronic infectious disease lasting 6 months or longer requiring hospitalization in the last year
Current involvement in any clinical trial
Pregnancy
Use of antibiotics, probiotics, immunomodulatory medications, or fecal microbiome transplant in the 90 days prior to screening.

Enrollment

Veterans meeting eligibility criteria attended an in-person enrollment session. During the enrollment visit, participants completed questionnaires assessing their medical history and lifestyle including smoking history, medications, alcohol use, and comorbidities. Participants also completed the National Cancer Institute's Dietary History Questionnaire III (DHQ III) (15,16) and a questionnaire assessing their known exposures to chemicals, biologics, and pharmaceuticals during their Gulf War deployment as well as their branch of service.

Lastly, participants completed the Kansas GWI case definitions assessment. Participants self-reported individual symptoms within the six GWI symptom domains (fatigue, pain, gastrointestinal, cognition, skin, and respiratory) and rated the severity of those symptoms as no, mild, moderate, or severe impacts on daily life in the previous six months. Deployed GWVs with GWI must have endorsed one or more moderate to severe symptoms in the fatigue domain and at least two of the other five symptom domains. Participants not meeting these criteria were classified as controls.

Sample collection and study follow-up

Stool samples were collected weekly for eight weeks. Participants were provided with stool collection kits and were trained by the research team on how to properly collect, store, and ship specimens. Participants collected samples within a five-day window each week and were asked to ship specimens overnight within 24 hours of collection. Samples not received within 72 hours were rejected. The study team called participants weekly to remind them to collect and ship their samples. During the weekly check-in, participants were asked about antibiotic or

probiotic use in the last week, about gastrointestinal symptoms, and if they had questions or comments for the study team.

DNA extraction and sequencing

The full laboratory methods for this study have previously been published.(14,17) Briefly, total genomic DNA was extracted using a phenol:chloroform:isoamyl alcohol and bead-beating protocol with additional enzymatic lysis containing mutanolysin, lysostaphin, and lysozyme to assist in lysing gram-positive cell walls. Samples were purified using the NucleoSpin Gel and PCR cleanup kit according to the manufacturer's directions (Macherey-Nagel, Germany) and stored at -80°C. DNA was quantified using the Qubit dsDNA kit (Invitrogen, Waltham, MA, USA) on the Biotek Synergy HTX (Biotek Instruments, Winooski, VT, USA). Samples were then sequenced using 16S rRNA sequencing of the V4 region on the Illumina MiSeq at the University of Wisconsin Biotechnology Center. Purified DNA was normalized to 5 ng/μL, amplified using barcoded primers for the 16S V4 region, and sequenced using 2x250 paired end reads.

Microbial analysis and statistics

Raw sequences were processed into amplicon sequencing variants (ASVs) using QIIME2(18) following the "Moving Pictures" protocol.(19) DADA2(20) was used for the quality control steps. Taxonomy was assigned using the GreenGenes(21) database and assigned to the genus level whenever possible. Statistical analyses were conducted using R version 4.1.0.(22) Alpha diversity was assessed using the Shannon, Inverse Simpson's diversity indices; richness was assessed using the Chao1 index. The Bray-Curtis dissimilarity matrix, which was used to assess beta diversity, was visualized using non-metric multidimensional scaling (NMDS). Permutational Analysis of Variance (PERMANOVA) was used to estimate

associations between GWI status and beta diversity.(23) In addition to GWI vs controls, alpha and beta diversity between those with and without gastrointestinal symptoms was also assessed. The Quasi-Conditional Association Test and Generalized Estimating Equation (QCAT-GEE) (24) was used to test for differentially abundant taxa (subset to the 100 most prevalent ASVs) between those with GWI, after adjustment for age, added sugars, the presence of gastrointestinal symptoms, pesticide exposure, and the Healthy Eating Index – 2015 (HEI). The HEI is a measure of overall diet quality (independent of food quantity) and is scored by assigning points to 13 dietary components with a maximum of 100 points.(25). A QCAT-GEE analysis was also done by HEI. The Benjamini-Hochberg correction for the false discovery rate (FDR) (26) was applied ($\alpha=0.05$). QCAT-GEE provides three tests for differences by GWI status: the positive part (differences in abundance of each taxa), the zero part (differences in the presence/absence of taxa), and the two part (combining the positive and zero parts). A p-value of ≤ 0.05 was used for all statistical tests.

Results

A total of 69 GWVs, 36 with GWI and 33 healthy controls, were included in the final analysis (Figure 1). The 69 patients provided 498 stool samples over eight weeks. The average age for the full cohort was 56.3 years; those with GWI were younger (54.2 years) than those without GWI (58.8 years, $p=0.006$). Six (9%) participants were female, three in each group. A majority of participants were members of the Army ($N=46$, 67%), followed by the Marines ($N=12$, 17%), Navy and Air Force (each $N=5$, 7%), and the Coast Guard ($N=1$, 1%). Fifty-nine (86%) participants identified as White. Full demographics data can be found in Table 2. Those with GWI reported significantly more moderate/severe symptoms across all symptom domains (Table 3). Twelve members of the GWI group reported experiencing three symptoms, 13 reported four symptoms, eight reported five symptoms, and three reported six symptoms. Six of those without GWI reported no symptoms, 11 reported one symptom, 12 reported two

symptoms, two reported three symptoms, and two reported four symptoms. All exposures except vaccinations were more prevalent in the GWI group (Table 4).

Figure 1. Flowchart of study enrollment

Table 2. Participant demographics by age group

	Full Cohort (N=69)	GWI (N=36)	Control (N=33)	p-value
Age (years)	56	59	54	0.006*
Gender				
Male	63 (91%)	33 (92%)	30 (91%)	
Female	6 (9%)	3 (8%)	3 (9%)	0.91
Branch				
Army	46 (67%)	26 (72%)	20 (67%)	
Navy	5 (7%)	1 (3%)	4 (13%)	
Marines	12 (17%)	7 (19%)	5 (17%)	
Air Force	5 (7%)	2 (6%)	3 (10%)	
Coast Guard	1 (1%)	0 (0%)	1 (3%)	0.41
Race*				
White	59 (86%)	31 (86%)	28 (85%)	
Other	8 (12%)	4 (11%)	4 (12%)	
Preferred not to answer	2 (3%)	1 (3%)	1 (3%)	0.99

Table 3. Reported moderate/severe symptoms and GWI domain by group

	Full Cohort (N=69)	GWI (N=36)	Control (N=33)
--	--------------------	------------	----------------

Symptom Domain			
Fatigue	46 (67%)	36 (100%)	10 (30%)
Pain	36 (52%)	33 (92%)	3 (9%)
Neurologic	43 (62%)	31 (86%)	12 (36%)
Skin	12 (17%)	10 (28%)	2 (6%)
Respiratory	14 (20%)	11 (31%)	3 (9%)
Gastrointestinal	26 (38%)	23 (64%)	3 (9%)
Number of Symptoms			
Zero	6 (7%)	0 (0%)	6 (18%)
One	11 (16%)	0 (0%)	11 (33%)
Two	12 (17%)	0 (0%)	12 (36%)
Three	14 (20%)	12 (33%)	2 (6%)
Four	15 (22%)	13 (36%)	2 (6%)
Five	8 (12%)	8 (22%)	0 (0%)
Six	3 (4%)	3 (8%)	0 (0%)

189

190 **Table 4. Reported exposures by group**

Exposure	Full Cohort (N=69)	GWJ (N=36)	Control (N=33)	p-value
Vaccinations	65 (94%)	33 (92%)	32 (97%)	0.62
Oil well fires	55 (80%)	32 (89%)	25 (76%)	0.21
Chemical/biologic weapons	27 (46%)	17 (47%)	10 (30%)	0.22

Other chemicals	23 (33%)	14 (39%)	9 (27%)	0.44
Pyridostigmine bromide	49 (71%)	29 (81%)	20 (61%)	0.11
Pesticides	24 (35%)	17 (47%)	7 (21%)	0.042*
Toxic embedded fragments	5 (7%)	5 (14%)	0 (0%)	0.055
Infectious disease	5 (7%)	4 (11%)	1 (3%)	0.36
Heat/high temperatures	11 (16%)	8 (22%)	3 (9%)	0.19

Overall, participants in both the control and GWI groups reported eating diets similar to the U.S. national average (59 pts) with the average Healthy Eating Index (HEI) at 61 pts (GWI: 58.2, Control: 63.8) using the National Cancer Institute's Dietary History Questionnaire III (DHQ III). Those in the GWI group did have a poorer diet and had a borderline significantly lower score (p-value:0.0535). Those in the GWI group also reported eating diets containing significantly more added sugars (GWI: 17.2 tsp/day, Control: 11.0 tsp/day, p-value: 0.044) and well above the recommended amount of added sugar (Table 5). No other significant differences were observed between the groups. Additionally, the mean dietary intakes for both groups fell below the USDAs recommended daily intakes for all macronutrients.(27)

Table 5. Mean dietary data by group

	Full Cohort (N=69)	GWI (N=36)	Control (N=33)	p-value	Recommended Value*
Healthy Eating Index	61.0	58.2	63.8	0.054	100 is highest score

Fiber					
Total Dietary Fiber (grams)	18.3	18.4	18.2	0.95	31 g/day
Soluble Dietary Fiber (grams)	7.1	7.0	7.2	0.88	No set recommendation s
Insoluble Dietary Fiber (grams)	11.2	11.3	11.0	0.86	No set recommendation s
Total Fruit (cups)	1.0	1.1	1.0	0.81	2.0
Total Vegetables (cups)	1.5	1.7	1.4	0.27	3.0
Total Legumes (cups)	0.1	0.1	0.1	0.74	Approx. 0.3 cups/day or 2 cups/week
Whole grains (ounces)	0.8	0.8	0.8	0.91	>3.5
Refined grains (ounces)	3.4	3.9	3.0	0.13	<3.5
Total protein foods (ounces)	5.4	5.7	5.0	0.46	6.0
Total Dairy (cups)	2.1	2.4	1.8	0.2	3.0
Added sugars (teaspoons)	14.2	17.2	11.0	0.044	<10% of calories/day or

					<13 tsp
Energy from total fat (%kcal)	35.4%	35.6%	35.2%	0.86	20-35%
Energy from carbohydrates (%kcal)	44.8%	44.9%	44.6%	0.9	45-65%
Energy from protein (% kcal)	15.9%	15.4%	16.4%	0.32	10-35%
Energy from saturated fats (% kcal)	12.1%	12.0%	12.1%	0.92	<10%

The gut microbiota for all participants were dominated by members of the Firmicutes and Bacteroides phyla. Participants in the control group who reported experiencing moderate/severe gut-related symptoms had a very different gut microbial composition compared to all other groups. The ratio of Bacteroidetes to Firmicutes (B/F ratio) for the full cohort was 54.9% to 38.0% (Figure 2). Actinobacteria (2.3%), Verrucomicrobia (2.5%), and Proteobacteria (1.8%) were the only other phyla consisting of over 1% of the ASVs.

Figure 2. Relative abundance bar plot of the top 8 phyla by aggregated by week and GWI status.

The Shannon, Inverse Simpson, and Chao1 alpha diversity index measures all showed a very stable microbiome over time, regardless of GWI status. Linear mixed-effects modeling of the three alpha diversity measures confirmed these findings ($p=0.28$) (Figure 3). While no differences between groups were statistically significant, a visual inspection of the data shows

slightly lower alpha diversity amongst those reporting moderate to severe gut symptoms regardless of GWI status (Figure 4).

Figure 3. Linear mixed-effects modeling of alpha diversity by week. Red line is those with GWI and blue line is the controls. Lighter bands represent the 95% confidence interval of the regression line.

Figure 4. Alpha diversity by week and GWI status stratified by the presence of moderate/severe gastrointestinal symptoms (Yes/No).

To assess beta diversity, the Bray-Curtis dissimilarity was calculated and visualized using NMDS (Figure 5). Visual inspection showed a great deal of overlap of the ellipses indicating the centroids are in similar places; however, the PERMANOVA analysis indicated group status (GWI vs. controls, and GWI +GI, GWI, Control +GI, and control) significantly explained 1.2% and 3% of the variation in the data respectively (p-value:0.001 for both). However, PERMANOVA assumes homoscedasticity and the data showed significantly different dispersions (p=0.002 via betadisper in vegan). To confirm results, analysis of similarities (ANOSIM) was done which also found GWI status to significantly impact the gut microbiota as well (p <0.0001). The microbiota was very stable over the duration of the study with “week” not being associated with beta diversity using either PERMANOVA or ANOSIM (p=1.0 and 0.999 respectively).

Figure 5. NMDS plot of the Bray-Curtis dissimilarity matrix by week and GWI status

The QCAT-GEE test for differential lineages identified 9 lineages at the family level that were significantly different between those with GWI and the healthy controls. This model was adjusted for the subjects’ age, the presence of gastrointestinal symptoms, pesticide exposure during deployment, HEI score, and added sugar consumption (in teaspoons). Three lineages (Lachnospiraceae, Oscillospiraceae, and Ruminococcaceae) were more prevalent in those without GWI and were significant across the positive-, zero-, and two-part tests (Table 6). Additionally, we tested for whether there were differences in lineages associated with eating

habits (HEI score) those experiencing moderate to severe gastrointestinal symptoms (adjusted for GWI status) (Table 7). Lineages associated with the Eggerthellaceae, Erysipelatoclostridiaceae, Lachnospiraceae, and Oscillospiraceae families were all significant across the positive-, zero-, and two-part tests with Erysipelatoclostridiaceae and Lachnospiraceae associated with decreasing HEI scores.

Table 6. Bacterial taxa associated with GWI adjusted for covariates at the family level

Taxa	Positive Part	Zero Part	Two-Part	Direction*
Acidaminococcaceae	0.001	0.849	0.008	↑
Eggerthellaceae	0.446	0.028	0.059	↑
Erysipelotrichaceae	0.0735	0.002	0.012	↓
Lachnospiraceae	0.001	0.0001	0.001	↓
Marinifilaceae	0.0001	0.214	0.0001	↑
Oscillospiraceae	0.007	0.004	0.0001	↓
Peptostreptococcaceae	0.986	0.015	0.035	↓
Ruminococcaceae	0.002	0.041	0.001	↓
Veillonellaceae	0.019	0.317	0.061	↑

P-values reported have FDR correction applied.

*Up arrows indicate higher abundance in those with GWI, down arrows indicate higher abundance in controls

Table 7. Bacterial taxa associated with HEI scores among those with gastrointestinal symptoms adjusted for GWI status at the family level

Taxa	Positive Part	Zero Part	Two-Part	Direction*
Acidaminococcaceae	0.058	0.012	0.007	↑

Christensenellaceae	0.499	0.019	0.040	↑
Eggerthellaceae	0.038	0.039	0.001	↑
Erysipelatoclostridiaceae	0.05	0.04	0.015	↓
Lachnospiraceae	0.002	0.0001	0.0001	↓
Marinifilaceae	0.813	0.001	0.001	↓
Oscillospiraceae	0.001	0.0001	0.0001	↑
Peptostreptococcaceae	0.379	0.005	0.15	↑
Ruminococcaceae	0.701	0.0001	0.001	↑
Veillonellaceae	0.961	0.039	0.939	↑

P-values reported have FDR correction applied.

*Up arrows indicate increasing abundance with increasing HEI scores, down arrows indicate increase abundance with lower HEI scores

Discussion

This longitudinal cohort study of GWVs with and without GWI found participants with GWI had significantly different microbiomes from those without GWI. Furthermore, the gut microbiota was relatively stable over the eight weeks, regardless of GWI status, which was not surprising given no intervention was part of the study. The gut microbiota of both those with and without GWI consisted almost entirely of members of the Firmicutes and Bacteroidetes phyla across all time points. While there was no difference in alpha diversity between the groups, there was a difference in beta diversity by both group and the presence of gut symptoms. Similar to alpha diversity, beta diversity was not impacted by sampling week. We also identified several lineages that differed significantly by GWI status including members of the families Lachnospiraceae, Oscillospiraceae, and Ruminococcaceae which were all more prevalent in

those without GWI. Last, it appears those participants reporting gastrointestinal symptoms with healthier eating habits had differing gut microbiota from those with less healthy diets.

While limited data on the impact of GWI and the gut microbiome exists, several studies have observed greater abundances in Lachnospiraceae in those without GWI.(13) However, in a pilot study of Veterans with GWI, Ruminococcaceae was found to be associated with GWI, while in our study, it was associated with the healthy participant group. One potential reason for this difference is the smaller sample sizes observed in the pilot study of Veterans with GWI (n=16 with stool samples, 3-5 per group).(13) Other studies assessing the gut microbiome in GWI mouse models have also found Ruminococcaceae to be more abundant in the healthy controls.(12)(28) Previous research on Crohn's disease and inflammatory bowel disease (IBD) have found members of the Ruminococcaceae family to be decreased in those with the disease state,(29) consistent with our findings. Several members of the Ruminococcaceae family are important in microbe-mediated carbohydrate metabolism breaking down starches and cellulose in the lower GI tract.(30) They are critical producers of acetate(31), a necessary short-chain fatty acid (SCFA) for gut health as well as serving as a critical nutrient for butyrate producing bacteria in the gut.(32)(33) While not assessed in our study, acetate is a potential mediator of the gut-brain axis.(34) Animals models of multiple sclerosis (MS) have shown acetate supplementation can help reduce clinical symptoms associated with MS in mice.(35) In our cohort, 31 (86.1%) of those with GWI endorsed experiencing moderate to severe neurologic symptoms. Lachnospiraceae has also been shown to be in decreased abundance in those with GWI in both human and mouse models.(13)(28)(36) Additionally, it has been shown to be reduced in those with ulcerative colitis(37), ME/CFS(38), and Crohn's disease(39). Members of the Lachnospiraceae family are also key carbohydrate metabolizers producing butyrate.(37)

SCFAs, including butyrate and acetate, are crucial to gut health. They are the end products of carbohydrate metabolism primarily produced through anaerobic fermentation of dietary fiber in the gut. They also promote epithelial barrier function, cell proliferation, and act as

a nutrient source for many beneficial microbes in the gut.(32) Dietary interventions, either through food or supplements, to increase SCFAs have been proposed as potential treatments for a wide range of gut disorders including GWI. Prior studies have shown Western and high fat diets are associated with GWI.(12)(28)(40) Additionally, high sugar diets have been associated with decreased SCFAs, increased gut membrane permeability, and decreased microbial diversity in mouse models of ulcerative colitis.(41) In our study, the diets of all participants fell below the daily recommended amounts of dietary fiber, vegetables, whole grains, and fruits. Additionally, participants in both groups met or exceeded the recommended amount of energy from fats (including saturated fats). Those with GWI consumed significantly more added sugars than those without GWI and had a slightly lower HEI score. Poor overall diet was associated with ten differentially abundant lineages, most notable Lachnospiraceae and Ruminococcaceae. Ruminococcaceae increased with higher HEI scores, while Lachnospiraceae trended towards greater abundance in those with lower scores. However, 33 ASVs were associated with this lineage, with some being more abundant in those with higher HEI scores (*Dorea*, *Anaerostipes*, *Lachnoclostridium*, and *Ruminococcus*) and some more abundant in those with lower scores (*Agathobacter*, *Blautia*, and *Rosburia*). Prior research has shown Lachnospiraceae to have a complicated relationship to gut health with diet having a large influence on abundance.(37) This, in conjunction with the low levels of Lachnospiraceae and Ruminococcaceae found in those with GWI, indicates dietary interventions focused on improving diet and increasing SCFA production may benefit Veterans with GWI.

Our study has several strengths and limitations. This is the first longitudinal cohort study on GWI assessing the role of the gut microbiome and diet. We had a low dropout rate with 86% of participants completing the study and a majority completing the dietary questionnaire (DHQ III). One limitation to the study was the DHQ III. While food frequency questionnaires like the DHQ III are standard means of collecting dietary data(42), it is cumbersome to complete and relies heavily on self-reported data of foods consumed over the last month. Studies have shown

self-report dietary intakes underestimate the true energy consumption and individuals tend to report intakes closer to their perceived norms as opposed to actual consumption.(43)
Participants in our cohort likely ate worse diets than reported.

Conclusions

Future studies assessing the role of diet and the gut microbiome should strive to collect more detailed dietary data through the use of food diaries to reduce recall bias. Additionally, research building upon these findings would benefit from the use of shotgun metagenomics sequencing and the addition of metabolomics. Both metabolomics and metagenomic data would provide insight into the functional role these microbiota play in the gut of those with GWI. Furthermore, these methods would allow researchers to examine the role of SCFAs. Last, these findings along with the existing literature on GWI, point to the role of dietary interventions in alleviating the gastrointestinal symptoms frequently experienced by those with GWI. Studies assessing dietary and other interventions to manipulate the gut microbiome are needed to examine the safety and efficacy of treatments aimed at reducing GWI symptoms.

Acknowledgments

The authors would like to thank Catherine Shaughnessy for her help during the recruitment, enrollment, and data collection phases of this study.

References

1. Fukuda K, Nisenbaum R, Stewart G, Thompson WW, Robin L, Washko RM, et al. Chronic multisymptom illness affecting Air Force veterans of the Gulf War. *JAMA*. 1998 Sep 16;280(11):981–8.
2. Gronseth GS. Gulf war syndrome: a toxic exposure? A systematic review. *Neurol Clin*. 2005 May;23(2):523–40.
3. Minshall D. Gulf War Syndrome: a review of current knowledge and understanding. *J R Nav Med Serv*. 2014;100(3):252–8.
4. Steele L. Prevalence and patterns of Gulf War illness in Kansas veterans: association of symptoms with characteristics of person, place, and time of military service. *Am J Epidemiol*. 2000 Nov 15;152(10):992–1002.
5. Chester JE, Rowneki M, Van Doren W, Helmer DA. Progression of intervention-focused research for Gulf War illness. *Mil Med Res*. 2019 Oct 18;6(1):31.
6. Zundel CG, Krengel MH, Heeren T, Yee MK, Grasso CM, Janulewicz Lloyd PA, et al. Rates of Chronic Medical Conditions in 1991 Gulf War Veterans Compared to the General Population. *Int J Environ Res Public Health*. 2019;16(6).
7. Institute of Medicine. Chronic Multisymptom Illness in Gulf War Veterans: Case Definitions Reexamined. Washington (DC): National Academies Press (US); 2014.
8. Maule AL, Janulewicz PA, Sullivan KA, Krengel MH, Yee MK, McClean M, et al. Meta-analysis of self-reported health symptoms in 1990-1991 Gulf War and Gulf War-era veterans. *BMJ Open*. 2018 Feb 13;8(2):e016086.
9. Frémont M, Coomans D, Massart S, De Meirleir K. High-throughput 16S rRNA gene sequencing reveals alterations of intestinal microbiota in myalgic encephalomyelitis/chronic fatigue syndrome patients. *Anaerobe*. 2013 Aug;22:50–6.
10. Lakhan SE, Kirchgessner A. Gut inflammation in chronic fatigue syndrome. *Nutr Metab*. 2010 Oct 12;7:79–79.
11. Alhasson F, Das S, Seth R, Dattaroy D, Chandrashekar V, Ryan CN, et al. Altered gut microbiome in a mouse model of Gulf War Illness causes neuroinflammation and intestinal injury via leaky gut and TLR4 activation. *PloS One*. 2017;12(3):e0172914.
12. Angoa-Pérez M, Zagorac B, Francescutti DM, Winters AD, Greenberg JM, Ahmad MM, et al. Effects of a high fat diet on gut microbiome dysbiosis in a mouse model of Gulf War Illness. *Sci Rep*. 2020 Jun 12;10(1):9529.
13. Janulewicz PA, Seth RK, Carlson JM, Ajama J, Quinn E, Heeren T, et al. The Gut-Microbiome in Gulf War Veterans: A Preliminary Report. *Int J Environ Res Public Health*. 2019 Oct 4;16(19).
14. Keating JA, Shaughnessy C, Baubie K, Kates AE, Putman-Buehler N, Watson L, et al. Characterising the gut microbiome in veterans with Gulf War Illness: a protocol for a longitudinal, prospective cohort study. *BMJ Open*. 2019 Aug 19;9(8):e031114.

15. National Cancer Institute. Diet History Questionnaire, Veriosn III [Internet]. Available from: <https://epi.grants.cancer.gov/dhq3/>
16. National Cancer Institute, Applied Research Program. Diet*Calc Analysis Program. 2005.
17. Eggers S, Malecki KM, Peppard P, Mares J, Shirley D, Shukla SK, et al. Wisconsin microbiome study, a cross-sectional investigation of dietary fibre, microbiome composition and antibiotic-resistant organisms: rationale and methods. *BMJ Open*. 2018 Mar 27;8(3):e019450.
18. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol*. 2019 Aug 1;37(8):852–7.
19. “Moving Pictures” tutorial [Internet]. QIIME2docs. Available from: <https://docs.qiime2.org/2021.4/tutorials/moving-pictures/>
20. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods*. 2016 Jul 1;13(7):581–3.
21. McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, et al. An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J*. 2011/12/01 ed. 2012 Mar;6(3):610–8.
22. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2021. Available from: <https://www.R-project.org/>
23. Dixon P. VEGAN, a package of R functions for community ecology. *J Veg Sci*. 2003 Dec 1;14(6):927–30.
24. Tang Z-Z, Chen G, Alekseyenko AV, Li H. A general framework for association analysis of microbial communities on a taxonomic tree. *Bioinforma Oxf Engl*. 2017 May 1;33(9):1278–85.
25. U.S. Department of Agriculture Food and Nutrition Service. Healthy Eating Index [Internet]. Available from: <https://www.fns.usda.gov/healthy-eating-index-hei>
26. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B Methodol*. 1995 Jan 1;57(1):289–300.
27. U.S. Department of Agriculture and U.S. Department of Health and Human Services. Dietary Guidelines for Americans, 2020-2025. In: 9th ed. 2020. Available from: [dietaryguidelines.gov](https://www.dietaryguidelines.gov)
28. Bose D, Saha P, Mondal A, Fanelli B, Seth RK, Janulewicz P, et al. Obesity Worsens Gulf War Illness Symptom Persistence Pathology by Linking Altered Gut Microbiome Species to Long-Term Gastrointestinal, Hepatic, and Neuronal Inflammation in a Mouse Model. *Nutrients*. 2020 Sep 10;12(9):2764.

29. Morgan XC, Tickle TL, Sokol H, Gevers D, Devaney KL, Ward DV, et al. Dysfunction of the intestinal microbiome in inflammatory bowel disease and treatment. *Genome Biol.* 2012 Apr 16;13(9):R79–R79.
30. Flint HJ, Bayer EA, Rincon MT, Lamed R, White BA. Polysaccharide utilization by gut bacteria: potential for new insights from genomic analysis. *Nat Rev Microbiol.* 2008 Feb;6(2):121–31.
31. Chassard C, Bernalier-Donadille A. H₂ and acetate transfers during xylan fermentation between a butyrate-producing xylanolytic species and hydrogenotrophic microorganisms from the human gut. *FEMS Microbiol Lett.* 2006 Jan;254(1):116–22.
32. Parada Venegas D, De la Fuente MK, Landskron G, González MJ, Quera R, Dijkstra G, et al. Short Chain Fatty Acids (SCFAs)-Mediated Gut Epithelial and Immune Regulation and Its Relevance for Inflammatory Bowel Diseases. *Front Immunol.* 2019;10:277.
33. Duncan SH, Hold GL, Barcenilla A, Stewart CS, Flint HJ. *Roseburia intestinalis* sp. nov., a novel saccharolytic, butyrate-producing bacterium from human faeces. *Int J Syst Evol Microbiol.* 2002 Sep;52(Pt 5):1615–20.
34. Silva YP, Bernardi A, Frozza RL. The Role of Short-Chain Fatty Acids From Gut Microbiota in Gut-Brain Communication. *Front Endocrinol.* 2020;11:25.
35. Chevalier AC, Rosenberger TA. Increasing acetyl-CoA metabolism attenuates injury and alters spinal cord lipid content in mice subjected to experimental autoimmune encephalomyelitis. *J Neurochem.* 2017 Jun;141(5):721–37.
36. Seth RK, Kimono D, Alhasson F, Sarkar S, Albadrani M, Lasley SK, et al. Increased butyrate priming in the gut stalls microbiome associated-gastrointestinal inflammation and hepatic metabolic reprogramming in a mouse model of Gulf War Illness. *Toxicol Appl Pharmacol.* 2018/05/09 ed. 2018 Jul 1;350:64–77.
37. Vacca M, Celano G, Calabrese FM, Portincasa P, Gobbetti M, De Angelis M. The Controversial Role of Human Gut Lachnospiraceae. *Microorganisms.* 2020 Apr 15;8(4):573.
38. Lupo GFD, Rocchetti G, Lucini L, Lorusso L, Manara E, Bertelli M, et al. Potential role of microbiome in Chronic Fatigue Syndrome/Myalgic Encephalomyelitis (CFS/ME). *Sci Rep.* 2021 Mar 29;11(1):7043.
39. Maukonen J, Kolho K-L, Paasela M, Honkanen J, Klemetti P, Vaarala O, et al. Altered Fecal Microbiota in Paediatric Inflammatory Bowel Disease. *J Crohns Colitis.* 2015 Dec 1;9(12):1088–95.
40. Holton KF, Ramachandra SS, Murray SL, Baron M, Baraniuk JN. Effect of the low glutamate diet on inflammatory cytokines in veterans with Gulf War Illness (GWI): A pilot study. *Life Sci.* 2021 Sep 1;280:119637.
41. Laffin M, Fedorak R, Zalasky A, Park H, Gill A, Agrawal A, et al. A high-sugar diet rapidly enhances susceptibility to colitis via depletion of luminal short-chain fatty acids in mice. *Sci Rep.* 2019 Aug 23;9(1):12294–12294.

- 455 42. Naska A, Ligiou A, Ligiou P. Dietary assessment methods in epidemiological research:
456 current state of the art and future prospects. F1000Research. 2017 Jun 16;6:926–926.
- 457 43. Schoeller DA. How accurate is self-reported dietary energy intake? Nutr Rev. 1990
458 Oct;48(10):373–9.
- 459