1 Self-management of depression and anxiety amongst frail 2 older adults in the United Kingdom: a qualitative study

3

4 Pushpa Nair^{1*}, Kate R. Walters¹, Su Aw², Rebecca L. Gould³, Kalpa Kharicha⁴,
5 Marta Buszewicz¹, Rachael Frost¹

6

7

8 ¹ Department of Primary Care and Population Health, University College London Medical
9 School (Royal Free Campus), London, United Kingdom.

10 ²National University of Singapore, Saw Swee Hock of Public Health, Singapore.

11 ³Division of Psychiatry, University College London, London, United Kingdom.

12 ⁴ Campaign to End Loneliness, What Works Centre for Wellbeing, The Evidence Quarter,
13 Albany House, London, United Kingdom.

14

15

16 *Corresponding author

17 Email: pushpa.nair@nhs.net (PN)

18

19

20

21

22

23

24 Abstract

25 **Objectives:** Depression and anxiety are common in frail older people and are associated
 26 with high levels of morbidity and mortality, yet they typically face greater barriers to
 27 accessing mental health treatments than younger people and express preferences for self-
 28 managing their symptoms. This study aims to explore frail older adults' experiences of self-
 29 managing symptoms of depression and/or anxiety.

30 **Design:** Qualitative semi-structured interviews, exploring experiences of depression and/or
 31 anxiety, ways participants self-managed these and the contexts within which this took place.
 32 Interviews were audio-recorded and transcribed verbatim.

33 **Participants:** 28 frail older adults in the United Kingdom, purposively sampled for
 34 neighbourhood, frailty and symptoms of anxiety/depression.

35 **Analysis:** Thematic analysis to inductively derive themes from the data.

36 **Results:** Our findings suggest that frail older adults find maintaining routines, engaging in
 37 meaningful and creative activities and socialising important to self-manage depression and
 38 anxiety. These could all be adapted to the level of frailty experienced. Stoicism, acceptance,
 39 faith or addressing the perceived cause were helpful in some situations and for some
 40 personalities. Distraction and avoidance were helpful for more severe symptoms or where
 41 the causes of symptoms could not be resolved. Self-management strategies were less well-
 42 established for anxiety symptoms, especially when linked to newer health fears and worries
 43 about the future.

44 **Conclusions:** Developing services and sources of information that support and facilitate key
 45 therapeutic components of self-management, which align with older adults' preferred coping
 46 styles and take into account levels of frailty, may be a way of supporting frail older people
 47 waiting for mental health treatments or those who prefer not to access these. Greater
 48 awareness of anxiety and how it can be self-managed in frail older people is needed.

49

50 Introduction

51 Worldwide, approximately 11% of adults aged 65+ years are frail [1]. Although
 52 lacking a universal definition, frailty is generally conceptualized as a dynamic process
 53 involving deficiencies across multiple physiological systems, with diminishing reserves and
 54 increased vulnerability to poor health outcomes [2], encompassing symptoms such as weight
 55 loss, muscle weakness, low physical activity and low energy levels [3]. However,
 56 psychological aspects of frailty are often overlooked [4].

57 Depression has a prevalence of 9% in adults aged 75+ years [5], and anxiety 15% in
 58 those aged 60+ [6]. Frail older adults are four times more likely to experience anxiety and/or
 59 depression than non-frail older adults [7]. Both are associated with higher levels of functional
 60 impairment [8, 9], cognitive decline [9, 10, 11] and healthcare service utilization [12, 13],
 61 especially when combined with frailty [14], and both remain commonly under-diagnosed [15].
 62 Depression in older adults has additionally been linked to greater risk of falls [16] and self-
 63 neglect [17], with a reciprocal relationship with frailty [18] and higher relapse rates than in
 64 earlier life [19]. The mental health of older adults has become an even more urgent concern
 65 considering the negative impact of the Covid-19 pandemic on wellbeing in this group [20].

66 Uptake of mental health services amongst older adults is poor - up to 70% of adults
 67 aged 55+ in the United States (US) with mood disorders do not access mental health
 68 services [21]. Within the National Health Service (NHS) in the United Kingdom (UK), mental
 69 health services can be freely accessed through a GP referral or self-referral; however only
 70 6.2% of national referrals to psychological therapy services within UK primary care settings
 71 are for those aged 65+, and this drops to 2.1% for those aged >75 [22], with an even greater
 72 reduction in referrals seen during the recent Covid-19 pandemic [23]. There is evidence that
 73 many older adults view depression as a normal part of ageing and often prefer self-
 74 management approaches over seeking professional help [24, 25]. Practical difficulties such
 75 as limitations in mobility and transport costs have also been cited as contributing to lower
 76 levels of access, particularly amongst frailer older adults [26]. There is evidence that frail

77 older people prefer to access mental health treatments through their GP, but experience
78 problems accessing their GP or finding information from alternative sources [27]. In addition,
79 there is evidence that healthcare professionals experience a wide range of barriers in
80 referring older people to psychological therapies [27]. Those who are referred for
81 psychological therapies can find they are on waiting lists as long as 12 months in some
82 areas, with little support available to them whilst waiting for this service [28]. Waiting times
83 have been further exacerbated by the Covid-19 pandemic [23], with certain services only
84 offering remote access, potentially leading to the digital exclusion of some older adults [29].

85 The value of current treatments for mood disorders in frailer populations is also
86 uncertain. The efficacy of antidepressants is less clear in frailer populations [30], with
87 multiple medications a known contributor to frailty itself [31]. Despite this, there is evidence
88 that the oldest old are more likely to be prescribed medication than be referred for
89 psychological therapies compared to those who are younger [32]. Psychological therapies in
90 frail older people may also be less effective, may need to be adapted to account for mild
91 cognitive impairment [33, 34], or may be less acceptable; for example, a US survey showed
92 that those aged 85+ preferred supportive treatment for anxiety rather than Cognitive
93 Behavioral Therapy [35].

94 In this context, self-management strategies may play a significant role in improving
95 mood disorders in this population [36]. Self-management is defined as 'taking increased
96 responsibility for one's own health, behaviour and well-being' [37]. It can encompass lifestyle
97 changes, problem solving, goal-setting and symptom monitoring [38, 39]. It is important to
98 distinguish self-management, which focuses more on maintaining wellbeing and reducing
99 relapses [39], from self-help, which aims to treat episodes of depression or anxiety using
100 remote or instructional CBT-based approaches [40]. Whilst help-seeking is also a part of
101 self-management, barriers to this have been extensively documented in older people [25,
102 41]. Ways that older people can support themselves align with a preference for feeling
103 independent at a time of increasing dependence [27], fear of stigma and a potential
104 inclination to deal with emotional problems on their own [25, 42, 43]. Self-management thus

has the potential to improve older adults' quality of life by facilitating feelings of empowerment and self-efficacy [38, 42].

Self-management in older populations has most commonly been studied in connection with long-term medical conditions [43, 44, 45], with studies showing good self-management outcomes. Programmes adapting the collaborative chronic care model (CCM), which includes self-management as a key component, to mental health care [46] have also shown promising outcomes for depression across all age groups [47, 48, 49]. However, most studies specifically exploring self-management of mood disorders have focused on younger populations and found that a wide range of strategies are used [50, 51], such as physical activity, social engagement, being proactive and having knowledge about the condition [39, 52].

Limited qualitative literature exploring aspects of self-management of mood disorders in older adults have focused mainly on depression in 'younger old' (<75 years) non-frail groups, highlighting the importance of strategies such as socializing, going out, engaging in religion, exercising and keeping up hobbies (findings summarized in three qualitative meta-analyses [25, 53, 54]). More recent studies [41, 42, 55, 56] have highlighted that a sense of identity, meaning and experiential knowledge are particularly important for older adults' capacity to self-manage depression. There is much less literature exploring self-management techniques for *anxiety* in older adults. Furthermore, we know little about the capacity and motivation of *frail* older people to self-manage depressive symptoms, despite the higher prevalence of symptoms in this subgroup, and much less about how they view and self-manage anxiety symptoms. Knowing more about their self-management techniques may help develop supportive resources that recommend acceptable strategies for alleviating mood symptoms in this population.

In this study, we therefore aimed to explore self-management strategies for both anxiety and depression amongst the frailer 'older old' group (75+ years), for whom some strategies (e.g. exercise, socializing) may be compromised by reduced mobility and multiple health comorbidities.

133

134 **Research Design and Methods**

135 The voices of frail older adults are often absent in research [57] and we therefore
136 chose a qualitative methodology that sought to elicit what helped them self-manage
137 symptoms, how frailty influenced this and whether their strategies differed from those of
138 younger-old adults.

139 **Ethics statement**

140 The study was approved by NHS Camden and Kings Cross Research Ethics
141 Committee (ref 17/LO/1963) and research was carried out in accordance with the ethical
142 principles for medical research involving human subjects expressed in the World Medical
143 Association Declaration of Helsinki [58]. Informed written consent was obtained for all
144 participants involved in the study.

145 **Participants**

146 We recruited 28 participants in total from five UK general practices, across one semi-
147 rural (n=10), one suburban (n=4) and three urban areas (n=14), in order to sample
148 neighbourhoods with different ethnicities, socioeconomic backgrounds, and variable access
149 to mental health services. The UK has a free national healthcare system (NHS) and referrals
150 to mental health services are primarily accessed through general practice, although patients
151 are also able to self-refer. Data was collected in 2018.

152 **Inclusion Criteria**

153 Practice staff and general practitioners (GPs) searched their patient lists to identify
154 older adults (aged 75+), classed as moderately to severely frail according to their practice's
155 frailty register, who were currently experiencing symptoms of depression and/or anxiety. We
156 asked practices to include a mix of those with and without a previously recorded diagnosis of

depression and/or anxiety in their medical notes, as depression and anxiety are often under-diagnosed in this population and we wanted to also include those who were currently experiencing symptoms but did not have a formal diagnosis and had not accessed previous help.

Exclusion Criteria

Older adults with advanced dementia, and those who had <6 months life expectancy were excluded.

Procedure

Practices posted invitation letters (with pre-paid postal reply slips) to patients who met the above inclusion criteria, and included study information leaflets containing brief screening questions for symptoms of anxiety and depression to encourage participants to self-identify (adapted from the 2-item Patient Health Questionnaire and 2-item Generalized Anxiety Disorder scale, which are both validated in older populations [59, 60] , but shortened and with the 2-week timescale changed to 'recently'). Interested participants returned reply slips to the two primary investigators (PN and RF) and were then telephone-screened to ensure they met the inclusion criteria, by using the same questions as in the leaflet to confirm symptoms, and asking about functional difficulties to confirm frailty. Of the 263 postal invites sent out, we received 41 positive responses, of which 28 were deemed eligible for study inclusion through screening, and 42 negative responses.

Data Collection

We collected data through individual semi-structured interviews, conducted face-to-face at participants' homes (or another location if requested). Interviews were conducted by PN (n=9, an academic GP, not disclosed to participants unless directly asked) and RF (n=19, a health services researcher). Terms such as 'low mood' and 'worry' were used at the beginning of interviews to reduce any perceived stigma, but terminology was adapted to reflect participants' own language as interviews progressed. Each interviewer observed 1-2 of the other's interviews to ensure a consistent approach. Interviews lasted approximately

one hour (range 34-89 minutes) and were audio-recorded. The interview topic guide was developed with the input of the whole research team, including our two patient and public involvement (PPI) representatives (who were older adults from diverse backgrounds interested in mental health research in older people) and was modified as interviews progressed. We asked interviewees about their typical week, past and present experiences of low mood and anxiety, any coping and self-management strategies, social/carer networks, and their views on treatments, help-seeking and suggestions for improvements in mental health services (the full topic guide can be found in [61]). As there were rich data across a wide range of themes, this current paper focuses on the contexts and forms of self-management of mood symptoms. Data related to help-seeking, current health care treatments and service improvement have been published elsewhere [61]. At the end of each interview, we collected demographic data. Participants were provided with a list of relevant support services, an Independent Age depression guide [62] and a £20 voucher. Audio-recorded interview data were transcribed verbatim by an external transcription service, and anonymized and verified for accuracy by the interviewer.

Data analysis

We used thematic analysis to analyze our data from a constructivist perspective, which emphasizes the importance of multiple perspectives, contexts and values [63]. Our team included two health services researchers (RF, KK), three academic GPs (KW, MB, PN), two psychologists (RG, SA) and two PPI members (WD and SL). All transcripts were read by PN and RF and at least one other team member. PN and RF developed a thematic framework (which can be found in [61]), which was refined at a team meeting. PN, RF and SA coded transcripts line-by-line using NVivo 12 [64], according to the thematic framework. After coding three pilot transcripts, PN, RF and SA met to check for coding consistency and refine the thematic framework. Relevant headings from the thematic framework included: current experiences of depression/anxiety, influencing contextual factors, impact of physical health/frailty, life history and self-management strategies. PN and RF had regular meetings with the team to discuss, agree on and refine emerging analytical themes. Saturation was

Demographics	Number of Participants
Age mean; range	80.71 years; 75-88 years
Gender (male:female)	9:19

considered to have occurred after 26 interviews, with minor and wider collective themes accounted for, and two further interviews were conducted to corroborate this.

Results

We recruited 28 participants with a mean age of 80.71 (SD 4.07, range 75-88). Participants were approximately two-thirds female (n=19) and predominantly White British (n=22). Ten participants were receiving Pension Credit, an indicator of low income in the UK (see Table 1 for demographic details). Whilst only eight participants reported receiving a formal diagnosis from a clinician (in their medical notes) of depression, anxiety or both, telephone screening identified nearly half the participants as experiencing current symptoms of both anxiety and depression, with the rest experiencing one or the other.

Table 1. Demographic Characteristics of Study Participants

Neighbourhood	Urban	14
	Suburban	4
	Semi-rural	10
Ethnicity	White British	22
	Black Caribbean	4
	Indian	1
	White Irish	1
Country of birth	United Kingdom	22
	Ireland	1
	Trinidad	2
	Jamaica	2
	Mauritius	1
Ethnicity	White British	22
	Black Caribbean	4
	Indian	1
	White Irish	1
Marital status	Married	8
	Widowed	10
	Separated	2
	Divorced	7
	Single	1
Living situation	Alone	17
	With spouse	8
	With other family	3
Housing	Owner-occupied	16
	Council rented	7
	Housing association rented	2
	Sheltered housing	2
	Private rented	1
Age completed education	Before 15 years	8
	15-17 years	7
	17-20 years	6
	Over 21 years	7
Pension type	State	28
	Employer	17
	Private	3
	Pension credits	10
Diagnosis (self-reported)	Depression	4
	Anxiety	4
	Depression & Anxiety	3
	No diagnosis	16
	Unsure	1
Telephone screening	Depression symptoms	10
	Anxiety symptoms	5
	Both	13

225

226

Participants reported a range of experiences of depression, ranging from transient

227

periods of feeling down, which varied depending on physical health and levels of functioning,

to constant levels of low mood, which were often underpinned by family conflict, financial difficulties or past traumatic life experiences. In terms of anxiety, some participants reported frequently worrying about ‘silly little things’, whilst others had strong health-related fears or fears about the future, and a few reported constant levels of worry about everything. As many participants experienced both anxiety and depression, it was sometimes difficult to distinguish differences in the way that they self-managed each, but this is highlighted where possible in the results. This paper reports on the wide range of approaches used by participants to manage symptoms of depression and anxiety, incorporating four key themes: 1.) Impact of frailty 2.) Engaging in meaningful routines and activities; 3.) Socializing; 4.) Internal mental and emotional coping strategies.

Impact of frailty

Participants reported changes in perceived identity resulting from functional decline, illness and fears over losing independence, which negatively affected their mood and wellbeing:

‘Old age isn’t like ‘flu. You get worse, like. That’s what worries me, and I think I don’t want to be a nuisance.’ (White British Female, age 81-85, depression and anxiety symptoms)

Many mentioned not being able to use strategies that were previously helpful in managing their mood due to changes in functional levels. One participant reported no longer being able to read books due to a deterioration in her eyesight, whilst others reported not being able to go out as much due to limitations in mobility or chronic health conditions. This also impacted on their ability to socialise and exercise, as well as continuing with previously enjoyed hobbies, such as cooking and shopping. For participants who had become carers, their changes in circumstances often meant that they were no longer able to engage in activities that had previously been important to them for maintaining wellbeing.

257 *'I used to do country dancing. And, as I say, now it's – well I suppose your*
 258 *wings are clipped, ain't they, apart from when he goes to the Day Center'.*
 259 (White British Female, age 81-85, anxiety symptoms)
 260
 261 *'I think the only thing that I'm missing out on is to go out and have, you know, do*
 262 *some exercise or some clubbing or something. And then I can come home, and, you*
 263 *know, feel like I'm somebody.'* (Black Caribbean Female, age 76-80, anxiety and
 264 depression symptoms)

265
 266 However, within the limitations of their frailty, participants described a wide range of
 267 strategies that helped them cope with symptoms of low mood and/or anxiety. They had
 268 developed these over their life-time and were able to implement them according to their
 269 current levels of functioning, as outlined below.

270

271 **Engaging in meaningful routines and activities**

272 **Maintaining a routine**

273 The ability to maintain a routine emerged as integral to many participants' sense of
 274 identity. Even if participants could not do some of the activities they previously enjoyed,
 275 being able to socialize or keep the house tidy were all seen as representative of their ability
 276 to continue with 'normal' life.

277

278 *'The more you go out, the more you become a normal, still remain a normal,*
 279 *active person'.* (White British Female, age 81-85, anxiety symptoms)

280

281 Participants often used downward social comparisons, comparing themselves favorably to
 282 other older adults who had lost their independence, and who seemed to represent 'the other'
 283 they feared turning into. Despite these activities often being more difficult to uphold or having
 284 to be adapted as people became frailer, they provided some structure to the day and were
 285 used by some to combat low motivation and keep negative thoughts at bay.

286

287 *'I've got a routine that I keep each morning. I do that, not because I'm*
 288 *pedantic, but because if I don't, I'm letting everything slide, you know.'* (White
 289 British Male, age 86-90, depression symptoms)
 290
 291 Family, neighbours and carers were generally identified as helpful in assisting them to keep
 292 to a routine and therefore supporting mental wellbeing. However, this needed to be carefully
 293 balanced with not appearing too dependent, and some feared that sharing any hint of being
 294 unable to keep up these activities with their families would either worry them or result in loss
 295 of their independence.
 296
 297 *'I want to do things for myself, continue as long as I can, you know. I don't*
 298 *want them (family) running and worrying and whatever...'* (Black Caribbean
 299 Female, age 76-80, anxiety and depression symptoms)

300 **Meaningful and creative activities**

301 In addition to upholding a routine, activities that promoted a sense of meaning and
 302 creativity were also important for mood and wellbeing, particularly for participants whose
 303 frailty limited their ability to go out. Creative activities, such as poetry, painting, writing,
 304 patchwork, music and photography, helped participants express and make sense of their
 305 emotions, and often helped to contextualize earlier life experiences and memories.
 306
 307 *'That's why I think I like patchwork because you use old material, you know,*
 308 *and there's a lot of memories in the old material'.* (White British Female, age
 309 81-85, anxiety symptoms)
 310
 311 *'I also write dotty poems and things like that, I've got quite a collection of*
 312 *them...But when I feel, you know, in that mood, sometimes it comes out that*
 313 *way.'* (White British Male, age 86-90, depression symptoms)
 314
 315 Some male participants, in particular, had taken on new self-directed projects, which
 316 were often sedentary and sometimes facilitated by the use of technology; for
 317 example, researching local history, programming or constructing family trees.

318

319 *‘Everyday, most days, and I’m doing the family history. I’m just tracing right*

320 *back to my ancestors.’* (White British Male, age 86-90, depression symptoms)

321

322 Searching for meaning could manifest itself as a need to be productive, have a sense of

323 purpose or contribute towards something, for example by volunteering or finding a new

324 project in life.

325

326 *‘Maybe if some people, instead of sitting down and twiddle their thumb, could find*

327 *something to do. You know, it would help. An interest. Find an interest in life.’* (Black

328 Caribbean Female, age 86-90, depression symptoms)

329

330 *‘And I think they should do more things, not silly things, not like birthday cards*

331 *and like that, but things – even if you’re knitting squares for people all to sew*

332 *together, to send to Africa or something, you know.’* (White British Female,

333 age 81-85, anxiety symptoms)

334

335 The desire to be productive could offset feelings of dependency, associated with

336 participants’ frailty. However, these wishes were sometimes expressed as hypothetical

337 statements rather than as practical plans, either because participants were not aware of

338 available services or organizations that could help them realize their aspirations or because

339 their frailty prevented them from doing so.

340

341 *‘I mean if I could help somebody else you know, I feel, you know, and then*

342 *probably that would help me’.* (White British Female, age 76-80, depression

343 symptoms)

344

345 *‘I mean I do want to get out of this, you know, and sort of to feel...like a normal*

346 *person sort of thing, doing things or helping somebody. That’s the thing.*

347 *Instead of just somebody helping me you know.’* (White British Female, age

348 76-80, depression and anxiety symptoms)

349

Socializing

Camaraderie and connectivity

Despite participants' frailty, many found ways to stay connected to family and friends, for example via the telephone or through attending local groups, while a few also used social media, particularly if they difficulties going out. Many participants cited speaking to friends, peers and spouses as useful forms of informal support. The value of this seemed to be underpinned by a sense of shared life experience and camaraderie, both in reviewing the past but also in facing the existential uncertainty of the future:

'But I think most of us are very philosophical about things like that, because we're together and we're in the same boat. So we laugh about it, you know. Sometimes the humor is very black. But, it helps, you know.' (White British Male, age 86-90, depression symptoms)

Many valued the reciprocal exchange of advice, as it drew on personal experiences and enhanced feelings of altruism. In addition to supporting mental wellbeing, exchange of advice around services providing help with practical things, such as finances, could also help people to overcome problems negatively affecting their mental health.

'So a lot of connection happens out of just, you know, pulling together on shared things'. (White British Male, age 76-80, depression symptoms)

Some participants remained depressed or anxious despite having good social links, particularly if they were experiencing more severe symptoms, although many still highlighted the positive aspects of having social support in keeping them going, in contrast to those who were more socially isolated and did not have significant family or friend networks.

'I mean some of these poor people here [in sheltered housing block], they really are on their own, full stop, yes. And I don't know how they do it.' (White British Male, age 86-90, depression symptoms)

Control

Interestingly, attitudes to formalized peer support groups or Day Centers, where people were grouped on the basis of age alone, were much less positive. Many participants expressed resistance to being labelled as 'old', did not identify with others in the group, or were frequently irritated by them.

'The girls keep saying to me, 'Go to the old people's clubs'. And to be honest with you, I wouldn't want to go up there and sit with all them farts and you know eat fish and chips or a roll and coffee. It's not me'. (White British Male, age 86-90, depression symptoms)

Some participants displayed a preference for being around younger people, who they felt were less likely to 'make them miserable'. Being able to exert some control over whom they talked to, where and when, seemed to be an important aspect of 'informal' support that made it work for them, and many disliked the artificiality of organized groups. Some participants felt that setting up more *relaxed* environments or spaces, such as that provided within sheltered housing, might be useful.

'Yes, I think there should be more places set up where people can go and talk'. (White British Female, age 76-80, depression symptoms)

Shared interest groups were also mentioned as potentially useful and were more acceptable than 'old people's groups'. Participants tended to rely on recommendations from friends and family, rather than actively seeking out information about local groups themselves.

Internal mental and emotional coping strategies

Participants demonstrated a range of internal strategies to deal with their symptoms of depression and anxiety. These strategies were varied and depended on participants' personalities, values and life experiences.

Endurance, resilience and stoicism

Many participants reported a range of previous traumatic life experiences, including evacuation and bombing during the Blitz, family bereavements (with four participants having experienced the death of adult children), and physical, sexual and emotional abuse. Many highlighted their generation's way of dealing with trauma, characterized by resilience developed through stoicism, and how this had shaped their own coping mechanisms, particularly for depression. For some, this underpinned a positive way of self-managing depressive symptoms. Because they had survived more severe episodes in the past, they felt that they knew from experience that periods of low mood were likely to pass with time if they remained resilient and rode it out.

'I think we've all come from a generation that's fairly stoic, with the war years and that...Mum didn't have to do everything for you and all the rest of it. And so when these things come along, you've got that experience, it helps you cope.' (White British Male, age 86-90, depression symptoms)

However, others felt that this approach was not helpful and that not dealing with the trauma at the time had caused them to repress painful memories, which consequently had lasting negative effects on their mental health.

[Referring to her brother's suicide] 'We didn't deal with it as a family...at one point I was trying, I was saying that I thought it would be really good for us to have some family therapy. But they wouldn't all agree.' (White British Female, age 71-75, depression and anxiety symptoms)

Drawing on life experience appeared to be employed less for anxiety symptoms, which were often exacerbated by newer health concerns and existential fears about the future (e.g. preparedness around death and dying, financial cost of funerals, sorting out inheritance) - worries that were much less familiar, and for which they needed to develop new coping strategies.

439

440 *'I think, "Ooh, I'd better get the will sorted out.".....Silly things that really don't*

441 *matter because I won't be here anyway. I want it all to run smoothly and [for*

442 *my children] to get their fair share of what's due to them.'* (White British Male,

443 age 76-80, depression and anxiety symptoms)

444

445 *'I know we're getting old and I can see the future is not good ahead. And I just*

446 *can't see where we're going and what we're doing.'* (White British Female, age

447 86-90, depression & anxiety)

448 **Acceptance**

449 Many participants demonstrated an acceptance of their situation, recognizing that

450 some things were outside of their control.

451

452 *'But there are certain things that you have no power over. And I think when*

453 *you realize that, you can, you can be more content'.* (White British Male, age

454 76-80, depression and anxiety symptoms)

455

456 Some felt that accepting your emotions and being allowed to occasionally wallow or have a

457 'good cry' were helpful outlets for their feelings.

458

459 *'Sometimes I think oh what's the matter with me and then I sort of get that feeling –*

460 *it's a frightened feeling. And then I cry and then I feel better.'* (White British Female,

461 age 81-85, depression and anxiety symptoms)

462 **Addressing the cause**

463 Many of those living with depression in particular, felt that they were aware of the

464 causes of their depression, particularly if they had experienced previous episodes. They

465 often demonstrated pragmatic outlooks, locating perceived solutions within their internal

466 control rather than in external interventions (e.g. psychological therapies or antidepressants).

467

468 *'If it's, you know, the reason why you're depressed you try to look at the, the*
469 *cause and you know, focus on that really, and not rely on medication'.* (White
470 British Female, age 76-80, depression symptoms)
471
472 *'But I think I can rationalise it a bit more now than I used to. You know, I've had that*
473 *much more experience. And, of course, that earlier [depressive] episode in life has*
474 *taught me a lot.'* (White British Male, age 86-90, depression symptoms)
475
476 However, those who reported experiencing more severe depression and/or anxiety often
477 expressed a sense of hopelessness or resignation, particularly if they lacked social support,
478 as they were aware of the causes but were unable to resolve these (e.g. past trauma, family
479 conflict). Anxiety, in particular, seemed more difficult to rationalize, and participants had
480 difficulty recognizing, processing and gaining perspective on their symptoms. Often, they
481 could not identify why they felt anxious, which meant they could not address its cause, or it
482 was a realistic anxiety (e.g. fear of falling, death, deteriorating health, worries about getting
483 dementia), or they minimized it and put it down to 'silly little things'. Consequently they often
484 downplayed their symptoms and rarely sought help unless their symptoms were severe.
485
486 *'Now I find that the slightest thing, it doesn't matter what it is, the smallest*
487 *thing, which is ridiculous, is enough for me to then start to worry.'* (White
488 British Male, age 76-80, depression & anxiety symptoms)
489
490 *'But I am terrified at not dying and people thinking I'm dead. ...And being*
491 *buried alive. Or even being in a coffin alive or even being in a body bag alive.*
492 *And in fact generally I feel it's a claustrophobic thing. That can actually bring*
493 *on really strong feelings of panic and phobia, which is irritating you know,*
494 *because it's very irrational.'* (White British Male, age 81-85, depression and
495 anxiety symptoms)

Distraction and avoidance

Nearly all participants described using distraction and avoidance to not dwell on things that made them feel sad or anxious; for example, watching television, playing Mahjong or doing crossword puzzles. Whilst most accepted that this was not a long-term fix, distraction was employed to a greater degree by those who felt that their problems lacked solutions, for example, those with chronic pain, financial problems or multiple health issues.

‘As soon as I start playing Mahjong, I forget what’s going on. I forget my [chronic] pain. I forget I’m down...’ (White British Female, age 76-80, depression symptoms)

‘I just put the television [on] to keep my mind off [financial problems, multiple chronic health conditions]. I just watch television and I, I don’t even remember, I put the television so loud.’ (South Asian Female, age 76-80, depression and anxiety symptoms)

Avoidance of triggering situations (such as going out or using public transport) appeared to be viewed as a short-term solution by some, especially those experiencing anxiety, but could lead to increased isolation and was not viewed as helpful in the long-term.

‘But because that is my fear, you know. I don’t, wouldn’t like to go out and anything, anything happen, and I’m outside on the road and nobody don’t know who I am. And so that’s why I stay indoors.’ (Black Caribbean Female, age 76-80, anxiety symptoms)

Having faith

Eight female participants, nearly all of whom were from Christian backgrounds, mentioned their faith as important in helping them cope with their depression and /or anxiety symptoms. Some felt being *listened to* by God and being able to have an internal dialogue were useful, whilst others felt that being able to see their lives as part of God’s plan and

having trust in Him were important in gaining perspectives on their situations. For some, the act of prayer in itself was seen as therapeutic.

'By the time I've finished with that prayer, I'm well.' (Black Caribbean Female, age

76-80, anxiety symptoms)

'I think my Christian faith and getting grace of strength that helped me keep going

through it all.' (White British Female, age 76-80, depression symptoms)

Discussion

We interviewed 28 frail older adults (aged 75+), who described a number of approaches and strategies for self-managing depression and/or anxiety symptoms. Upholding personal routines and engaging in meaningful activities were important in nurturing a sense of identity and independence, which participants feared losing as part of their frailty. Meaningful activities for frailer older people were often those that were creative or productive, such as craftwork, photography, music, writing and researching, which could easily be undertaken at home in the context of their frailty. In our study, frail older adults found fostering their own social connections important in promoting a sense of camaraderie and positivity, which helped them manage their mood. This was facilitated through local groups and increased use of phones and social media where participants left the home less often. Internal mental and emotional coping strategies employed were varied and shaped according to earlier life experiences, values and contexts. Endurance and stoicism could foster a sense of resilience towards current and future adverse circumstances for some, but for others could lead to unresolved issues. Acceptance of situations beyond their control and attempting to solve underlying causes were used to manage depressive symptoms, but anxiety was harder to rationalize and cope with as it rarely had a clear cause. Participants also used distraction and avoidance, which was helpful for severe symptoms and situations

that could not be changed, but this could be maladaptive if it stopped people doing activities such as going out.

Our findings suggest that frail older people use similar strategies to self-manage low mood as robust younger-old people, with sociability, self-efficacy, resilience and life experience being important themes [25, 53, 54]. Frailty limited the use of some self-management strategies, such as going out or exercising, but participants were able to find alternative activities that were meaningful to them. This is in agreement with existing research that meaningful activities reduce depression in older adults [65] and that meaning is an important enabler for self-management [55]. However, previous studies have focused mainly on outside interventions or ways to increase engagement with existing services. In line with the recent literature [42, 55, 56], results from our study suggest that interventions should facilitate older adults to draw on the wealth of internal mental and emotional coping strategies utilized to manage their mood without accessing formal support; a finding that has also been highlighted as important in the management of loneliness in older adults [66] .

Whilst previous research has suggested that self-management may be helpful for anxiety [67], this relies on older adults having the tools to successfully manage anxiety. A major finding from our study suggests that anxiety, often underpinned by newer health fears and worries about the future, may be more difficult to self-manage than depression as most frail older adults interviewed had not developed effective coping strategies for this other than distraction and avoidance [68]. Only 2 participants in our sample mentioned relaxation techniques such as controlled breathing and meditation, suggesting simple advice or resources to raise awareness of these kind of techniques may be helpful. Findings from a sister paper [61] have also highlighted uncertainty about the validity of seeking professional help for anxiety amongst frail older adults. The conclusion from both papers, in keeping with other international qualitative studies on anxiety in older adults [69, 70], suggests more work needs to be done to raise awareness of anxiety as a diagnosis and find helpful ways of ameliorating anxiety symptoms in this population.

Strengths and weaknesses

We captured a wide range of views from older adults from a variety of socioeconomic backgrounds, degrees of frailty and symptom severity. We placed an equal focus on both symptoms of anxiety and depression, which are often interconnected and demonstrate high levels of comorbidity [71]. Our topic guide and thematic framework were developed with a range of team member and PPI perspectives. We found no noteworthy differences in self-management according to socioeconomic background or gender. However, we had a low response rate from more severely frail older adults (e.g. from nursing homes), older men, ethnic minority groups and non-English speakers. In our sample, relatively few reported a formal diagnosis of anxiety and/or depression, which may reflect low rates of help-seeking and diagnosis in this population, or that the experiences of those with more severe symptoms may not have been fully explored. We were unable to access participants' medical notes to verify self-reported diagnoses, and we did not collect specific data on current or previous treatments, although about two thirds of our sample reported having tried antidepressants. Furthermore, as participants were all currently experiencing some degree of low mood and/or anxiety symptoms, the longer-term impact of these self-management strategies was unclear, and we did not objectively assess symptom severity. It is therefore possible that individuals may perceive their strategies to be effective when they are not in the longer-term, and future prospective studies are needed to explore the effectiveness of different self-management strategies. Finally, the results of this study reflect the views of a high-income Western country with free national healthcare and may not be transferrable to lower-income countries, which may have higher levels of frailty and more limited resources [72].

Implications

Information and advice for frail older adults to address depression and anxiety should communicate techniques that are acceptable and helpful (e.g. maintaining routines, meaningful activities, finding ways to socialize despite limited mobility), as well as

highlighting the benefits and limitations of self-management techniques that may vary according to personality, situation and symptoms (e.g. acceptance, stoicism, distraction). Frail older adults should be encouraged to reflect on what has helped them in the past and what is meaningful for them. Further awareness needs to be raised regarding anxiety and ways in which frail older people can self-manage such symptoms. However, as frail older adults rarely seek mental health information from sources other than their GP [27, 61] (whom they may delay seeing until symptoms are severe), multiple avenues are likely to be needed to communicate these, such as through voluntary community and social enterprise (VCSE) organizations, healthcare settings (e.g. waiting room posters/leaflets, pharmacists) and interactions with healthcare professionals.

 GPs might help frail older adults maintain their independence for longer by early referral to multidisciplinary services and relevant social prescribing services. As in other studies, we found that meaningful social activities were important [73], and that older adults preferred informal social groups of their own choosing rather than formalized groups which collectively labelled them as 'old' [74, 75]. Previous studies have highlighted a high degree of internalized ageism in older adults' attitudes [25], which is often associated with negative health outcomes [76]. Our study highlighted that control over who they talked to and in what environment was important for our participants. Creating community spaces for frail older adults to get together or offering inter-generational groups that center around a common interest may be useful for improving wellbeing in this group. For those with mobility limitations, these spaces may need to be incorporated closer to home, e.g. within sheltered housing, with encouragement to use remote forms of communication such as phone and video calling. VCSE workers would be well-placed to help frail older adults identify available appropriate services and could receive training to provide basic advice on self-management techniques. A recent study looked at the benefits of a guided self-management approach for anxiety in older veterans [77]. This may help with recognition and understanding of these symptoms which were clearly troubling for many of our participants.

Conclusion

There is an urgent need for mental health support for older adults, given the context of an increasingly ageing global population, and the need is even greater in wake of the Covid-19 pandemic [20], which has also put additional pressures on mental health services and increased waiting times [23]. For frail older adults with mood disorders, self-management may have an important and acceptable role in maintaining wellbeing, preventing relapse and providing an adjunct to other treatments, as well as offering an alternative to formal treatment for those with less severe symptoms, those who are on waiting lists for therapy or who are not able or willing to engage with services. Our research suggests that frailty limits some strategies used to self-manage mood, but that people adapt their self-management techniques to account for this, for example, focusing more on sedentary, creative pursuits or maintaining routines. Helping frail older adults maintain independence and engage in meaningful and social activities were all identified as potentially important in the self-management of depression and anxiety in this population. Supporting frail older adults' internal coping strategies may be useful for dealing with depression; however, those with anxiety may need further advice and techniques to manage their symptoms. Self-management support is likely to be more helpful when tailored to the individual, taking into account personal preferences, life experiences, levels of functioning and available resources.

Acknowledgements

We would like to thank our PPI representatives, Wes Dowridge and Shamime Lakda, for their valuable input throughout this project.

References

1. Collard RM, Boter H, Schoevers RA, Oude Voshaar RC. Prevalence of frailty in community-dwelling older persons: a systematic review. *J Am Geriatr Soc*. 2012;60(8):1487-92. doi: 10.1111/j.1532-5415.2012.04054.x.
2. Clegg A, Hassan-Smith Z. Frailty and the endocrine system. *Lancet Diabetes Endocrinol*. 2018;6(9):743–52.
3. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56(3):M146-56.
4. Grenier A. Constructions of frailty in the English language, care practice and the lived experience. *Ageing Soc*. 2007;27(3):425–45.
5. Rodda J, Walker Z, Carter J. Depression in older adults. *BMJ*. 2011;343(sep28 1):d5219.
6. Bryant C, Jackson H, Ames D. The prevalence of anxiety in older adults: methodological issues and a review of the literature. *J Affect Disord*. 2008;109(3):233–50.
7. Ní Mhaoláin AM, Fan CW, Romero-Ortuno R, Cogan L, Cunningham C, Kenny R-A, et al. Frailty, depression, and anxiety in later life. *Int Psychogeriatr*. 2012;24(8):1265–74.
8. Blackburn P, Wilkins-Ho M, Wiese BS. Depression in older adults: Diagnosis and management. *British Columbia Medical Journal*. 2017;59(3):171–7.
9. Potter GG, McQuoid DR, Whitson HE, Steffens DC. Physical frailty in late-life depression is associated with deficits in speed-dependent executive functions: Frailty and executive dysfunction in LLD. *Int J Geriatr Psychiatry*. 2016;31(5):466–74.

10. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*. 2020;396(10248):413–46.
11. Pocklington C. Depression in older adults. *British Journal of Medical Practitioners*. 2017;10(1).
12. Meeks TW, Vahia IV, Lavretsky H, Kulkarni G, Jeste DV. A tune in “a minor” can “b major”: a review of epidemiology, illness course, and public health implications of subthreshold depression in older adults. *J Affect Disord*. 2011;129(1–3):126–42.
13. Wolitzky-Taylor KB, Castriotta N, Lenze EJ, Stanley MA, Craske MG. Anxiety disorders in older adults: a comprehensive review. *Depress Anxiety*. 2010;27(2):190–211.
14. Dent E, Hoogendijk EO. Psychosocial factors modify the association of frailty with adverse outcomes: a prospective study of hospitalised older people. *BMC Geriatr*. 2014;14:108.
15. Mitchell AJ, Rao S, Vaze A. Do primary care physicians have particular difficulty identifying late-life depression? A meta-analysis stratified by age. *Psychother Psychosom*. 2010;79(5):285–94.
16. Lee D-CA, Lalor AF, Russell G, Stolwyk R, Brown T, Mcdermott F, et al. Understanding temporal relationships between depression, falls, and physical activity in a cohort of post-hospitalized older adults: A breakthrough or a conundrum? *International Psychogeriatrics*. 2017;29(10):1681–92.
17. Hansen MC, Flores DV, Coverdale J, Burnett J. Correlates of depression in self-neglecting older adults: A cross-sectional study examining the role of alcohol abuse and pain in increasing vulnerability. *J Elder Abuse Negl*. 2016;28(1):41–56.
18. Soysal P, Veronese N, Thompson T, Kahl KG, Fernandes BS, Prina AM, et al. Relationship between depression and frailty in older adults: A systematic review and meta-analysis. *Ageing Res Rev*. 2017;36:78–87.

19. Herron JW, Mitchell A. Depression and antidepressant prescribing in the elderly. *Prescriber*. 2018;29(3):12–7.
20. De Pue S, Gillebert C, Dierckx E, Vanderhasselt M-A, De Raedt R, Van den Bussche E. The impact of the COVID-19 pandemic on wellbeing and cognitive functioning of older adults. *Sci Rep*. 2021;11(1):4636.
21. Byers AL, Arean PA, Yaffe K. Low use of mental health services among older Americans with mood and anxiety disorders. *Psychiatr Serv*. 2012;63(1):66–72.
22. NHS Digital. Psychological Therapies: reports on the use of IAPT services. 2020. Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/psychological-therapies-report-on-the-use-of-iapt-services>.
23. Purrington J, Beail N. The impact of Covid-19 on access to psychological services. *Adv Ment Health Intellect Disabil*. 2021;15(4):119–31.
24. Burroughs H, Lovell K, Morley M, Baldwin R, Burns A, Chew-Graham C. Justifiable depression': How primary care professionals and patients view late-life depression? A qualitative study. *Family Practice*. 2006;23:369–77.
25. Nair P, Bhanu C, Frost R, Buszewicz M, Walters K. A systematic review and thematic synthesis of older adults' perceptions, beliefs and attitudes about depression and its treatment. *The Gerontologist*. 2019;1–12.
26. Van Der Weele GM. Response to an unsolicited intervention offer to persons aged >75 years after screening positive for depressive symptoms: a qualitative study. *International Psychogeriatrics*. 2012;24(2):270–7.
27. Frost R, Beattie A, Bhanu C, Walters K, Ben-Shlomo Y. Management of depression and referral of older people to psychological therapies: a systematic review of qualitative studies. *Br J Gen Pract*. 2019;69(680):e171–81.
28. MIND. We Still need to talk. A Report on Access to Talking Therapies. 2013. Available from: https://www.mind.org.uk/media-a/4248/we-still-need-to-talk_report.pdf

29. MIND. Digital services for people with mental health problems and digital exclusion during the coronavirus pandemic. 2020. Available from: <https://www.mind.org.uk/media/6693/briefing-digital-services-during-coronavirus-2020-final.pdf>
30. Mallery L, MacLeod T, Allen M, McLean-Veysey P, Rodney-Cail N, Bezanson E, et al. Systematic review and meta-analysis of second-generation antidepressants for the treatment of older adults with depression: questionable benefit and considerations for frailty. *BMC Geriatr*. 2019;19(1):306.
31. Gutiérrez-Valencia M, Izquierdo M, Cesari M, Casas-Herrero Á, Inzitari M, Martínez-Velilla N. The relationship between frailty and polypharmacy in older people: A systematic review: Frailty and polypharmacy: A systematic review. *Br J Clin Pharmacol*. 2018;84(7):1432–44.
32. Walters K, Falcaro M, Freemantle N, King M, Ben-Shlomo Y. Sociodemographic inequalities in the management of depression in adults aged 55 and over: an analysis of English primary care data. *Psychol Med*. 2018;48(9):1504–13.
33. Gould RL, Coulson MC, Howard RJ. Efficacy of cognitive behavioral therapy for anxiety disorders in older people: a meta-analysis and meta-regression of randomized controlled trials. *J Am Geriatr Soc*. 2012;60(2):218–29.
34. Frost R, Bauernfreund Y, Walters K. Non-pharmacological interventions for depression/anxiety in older adults with physical comorbidities affecting functioning: systematic review and meta-analysis. *Int Psychogeriatr*. 2019;31(8):1121–36.
35. Mohlman J. A community based survey of older adults' preferences for treatment of anxiety. *Psychol Aging*. 2012;27(4):1182–90.
36. Polacsek M, Boardman GH, McCann TV. A theory on the components of depression self-management in older adults. *Qual Health Res*. 2021;31(1):160–71.
37. Department of Health. Supporting people with long term conditions to self-care. Department of Health. 2006;29. Available from:

765 <https://www.yearofcare.co.uk/sites/default/files/images/Supporting%20people%20with%20long%20term%20conditions.pdf>

766

767 38. Barlow J, Wright C, Sheasby J, Turner A, Hainsworth J. Self-management

768 approaches for people with chronic conditions: a review. *Patient Educ Couns*. 2002;48(2):177–87.

769

770 39. Houle J, Gascon-Depatie M, Bélanger-Dumontier G, Cardinal C. Depression self-

771 management support: a systematic review. *Patient Educ Couns*. 2013;91(3):271–9.

772

773 40. Duggal HS. Self-management of depression: Beyond the medical model. *Perm J*. 2019;23.

774

775 41. Polacsek M, Boardman GH, McCann TV. Help-seeking experiences of older adults

776 with a diagnosis of moderate depression. *Int J Ment Health Nurs*. 2019;28(1):278–87.

777

778 42. Polacsek M, Boardman GH, McCann TV. Factors influencing self-management of

779 depression in older adults: a qualitative study. *Aging Ment Health*. 2020;24(6):939–46.

780

781 43. Reynolds K, Medved M, Mackenzie CS, Funk LM, Koven L. Older adults' narratives

782 of seeking mental health treatment: Making sense of mental health challenges and “muddling through” to care. *Qual Health Res*. 2020;30(10):1517–28.

783

784 44. Jonker AAGC, Comijs HC, Knipscheer KCPM, Deeg DJH. Promotion of self-

785 management in vulnerable older people: a narrative literature review of outcomes of the Chronic Disease Self-Management Program (CDSMP). *Eur J Ageing*. 2009;6(4):303–14.

786

787 45. Kastner M, Cardoso R, Lai Y, Treister V, Hamid JS, Hayden L, et al. Effectiveness of

788 interventions for managing multiple high-burden chronic diseases in older adults: a systematic review and meta-analysis. *CMAJ*. 2018;190(34):E1004–12.

789

790 46. Wagner EH, Austin BT, Davis M, Hindmarsh M, Schaefer J, Bonomi A. Improving

791 chronic illness care: Translating evidence into action. *Health Aff Millwood*. 2001;20(1):64–78.

792

47. Gilbody S, Bower P, Fletcher J, Richards D, Sutton AJ. Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes: A cumulative meta-analysis and review of longer-term outcomes. *Arch Intern Med*. 2006;166(21):2314–21.
48. Miller CJ, Grogan-Kaylor A, Perron BE, Kilbourne AM, Woltmann E, Bauer MS. Collaborative chronic care models for mental health conditions: cumulative meta-analysis and metaregression to guide future research and implementation. *Med Care*. 2013;51(10):922–30.
49. Woltmann E, Grogan-Kaylor A, Perron B, Georges H, Kilbourne AM, Bauer MS. Comparative effectiveness of collaborative chronic care models for mental health conditions across primary, specialty, and behavioral health care settings: systematic review and meta-analysis. *Am J Psychiatry*. 2012;169(8):790–804.
50. Van Grieken RA, van Tricht MJ, Koeter MWJ, van den Brink W, Schene AH. The use and helpfulness of self-management strategies for depression: The experiences of patients. *PLoS One*. 2018;13(10):e0206262.
51. Villaggi B, Provencher H, Coulombe S, Meunier S, Radziszewski S, Hudon C, et al. Self-management strategies in recovery from mood and anxiety disorders. *Glob Qual Nurs Res*. 2015;2:2333393615606092.
52. Turner A, Realpe AX, Wallace LM, Kosmala-Anderson J. A co-produced self-management programme improves psychosocial outcomes for people living with depression. *Ment Health Rev (Brighton)*. 2015;20(4):242–55.
53. Corcoran J, Brown E, Davis M, Pineda M, Kadolph J, Bell H. Depression in older adults: a meta-synthesis. *J Gerontol Soc Work*. 2013;56(6):509–34.
54. Holm AL, Severinsson E. Surviving depressive ill-health: a qualitative systematic review of older persons' narratives: Surviving depressive ill-health. *Nurs Health Sci*. 2014;16(1):131–40.

55. Polacsek M, Boardman GH, Mccann TV. A (2021a): Self Identity and Meaning in Life as Enablers for Older Adults to Self-Manage Depression. Issues in Mental Health Nursing.
56. Smit D, Peelen J, Vrijzen JN, Spijker J. An exploration of the conditions for deploying self-management strategies: a qualitative study of experiential knowledge in depression. BMC Psychiatry. 2020;20(1):210.
57. Benraad CEM, Kamerman-Celie F, van Munster BC, Oude Voshaar RC, Spijker J, Olde Rikkert MGM. Geriatric characteristics in randomised controlled trials on antidepressant drugs for older adults: a systematic review: Geriatric characteristics in antidepressant drug RCTs. Int J Geriatr Psychiatry. 2016;31(9):990–1003.
58. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. JAMA. 2013 Nov 27;310(20):2191-4. doi: 10.1001/jama.
59. Li C, Friedman B, Conwell Y, Fiscella K. Validity of the Patient Health Questionnaire 2 (PHQ-2) in identifying major depression in older people: Validity of phq-2 in detecting depression. J Am Geriatr Soc. 2007;55(4):596–602.
60. Wild B, Eckl A, Herzog W, Niehoff D, Lechner S, Maatouk I, et al. Assessing generalized anxiety disorder in elderly people using the GAD-7 and GAD-2 scales: results of a validation study. Am J Geriatr Psychiatry. 2014;22(10):1029–38
61. Frost R, Nair P, Aw S, Gould RL, Kharicha K, Buszewicz M, et al. Supporting frail older people with depression and anxiety: a qualitative study. Aging Ment Health. 2020;24(12):1977–84.
62. Independent Age. Dealing with depression: How to recognise the signs and the action you can take. 2017. Available from: <https://www.independentage.org/information/advice-guides-factsheets-leaflets/dealing-depression>

63. Lincoln Y, Guba E. Paradigmatic controversies, contradictions and emerging confluences. In: Denzin N, Lincoln, Y, editors. The Landscape of Qualitative Research: Theories and Issues. London: SAGE Publications Inc; 2003. p. 253–91.
64. QSR International Pty Ltd. NVivo qualitative data analysis software, Version 12. 2018. Available from: https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/home?_ga=2.154458939.2014665091.1643195778-2060129496.1643195778
65. Fiske A, Wetherell JL, Gatz M. Depression in older adults. *Annu Rev Clin Psychol.* 2009;5(1):363–89.
66. Kharicha K, Manthorpe J, Iliffe S, Chew-Graham CA, Cattan M, Goodman C, et al. Managing loneliness: a qualitative study of older people’s views. *Aging Ment Health.* 2021;25(7):1206–13.
67. Shepardson RL, Tapio J, Funderburk JS. Self-management strategies for stress and anxiety used by nontreatment seeking Veteran primary care patients. *Mil Med.* 2017;182(7):e1747–54.
68. Taylor SE, Stanton AL. Coping resources, coping processes, and mental health. *Annu Rev Clin Psychol.* 2007;3(1):377–401.
69. Kingstone T, Burroughs H, Bartlam B, Ray M, Proctor J, Shepherd T, et al. Developing a community-based psycho-social intervention with older people and third sector workers for anxiety and depression: a qualitative study. *BMC Fam Pract.* 2017;18(1).
70. Knight BG, Winterbotham S. Rural and urban older adults’ perceptions of 8 mental health services accessibility. *Aging & Mental Health.* 2019;0:1–7.
71. Braam AW, Copeland JRM, Delespaul PAEG, Beekman ATF, Como A, Dewey M, et al. Depression, subthreshold depression and comorbid anxiety symptoms in older Europeans: results from the EURODEP concerted action. *J Affect Disord.* 2014;155:266–72.

72. Siriwardhana DD, Hardoon S, Rait G, Weerasinghe MC, Walters KR. Prevalence of frailty and prefrailty among community-dwelling older adults in low-income and middle-income countries: a systematic review and meta-analysis. *BMJ Open*. 2018;8(3):e018195.
73. Forsman AK, Schierenbeck I, Wahlbeck K. Psychosocial interventions for the prevention of depression in older adults: systematic review and meta-analysis. *J Aging Health*. 2011;23(3):387–416.
74. Kharicha K, Iliffe S, Manthorpe J, Chew-Graham CA, Cattan M, Goodman C, et al. What do older people experiencing loneliness think about primary care or community based interventions to reduce loneliness? A qualitative study in England. *Health Soc Care Community*. 2017;25(6):1733–42.
75. Lee SH, Kim YB. Which type of social activities decrease depression in the elderly? An analysis of a population-based study in South Korea. *Iran J Public Health*. 2014;43(7):903–12.
76. Robertson G. Ageing and ageism: the impact of stereotypical attitudes on personal health and well-being outcomes and possible personal compensation strategies. *Self Soc*. 2017;45(2):149–59.
77. Gould CE, Carlson C, Wetherell JL, O'Hara R, Goldstein MK, Loup JR, et al. Guided self-management targeting anxiety and activity participation in older Veterans. *Aging Ment Health*. 2021;25(10):1913–22.

899

900

901

902

903

904

905

906

907