COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data

Authors: JOÃO F. PEREIRA

– Department of Epidemiology, Instituto Nacional de Saúde Dr. Ricardo Jorge, and
Department of Mathematics, University of Trás-os-Montes e Alto Douro
Portugal
(pereira.jpf96@gmail.com)

CONSTANTINO CAETANO

– Department of Epidemiology, Instituto Nacional de Saúde Dr. Ricardo Jorge and
Center for Computational and Stochastic Mathematics, Instituto Superior Técnico, University of Lisbon
Portugal
(constantino.caetano@insa-min-saude.pt)

LILIANA ANTUNES

– Department of Epidemiology, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, and
CEAUL – Centro de Estatística e Aplicações, Faculdade de Ciências, Universidade de Lisboa
Portugal
(liliana.antunes@insa-min-saude.pt)

PAULA PATRÍCIO

– Center for Mathematics and Applications (CMA), FCT NOVA and
Department of Mathematics, FCT NOVA,
Quinta da Torre, 2829-516 Caparica,
Portugal
(pcpr@fct.unl.pt)

MARIA LUÍSA MORGADO

– Department of Mathematics, University of Trás-os-Montes e Alto Douro, and
Center for Computational and Stochastic Mathematics, Instituto Superior Técnico, University of Lisbon
Portugal
(luisam@utad.pt)

BALTAZAR NUNES

– Department of Epidemiology, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, and
Centro de Investigação em Saúde Pública, Escola Nacional de Saúde Pública, Universidade NOVA de Lisboa, Lisbon, Portugal
(baltazar.nunes@insa.min-saude.pt)

*Corresponding author

1 Received: Month 0000 Revised: Month 0000 Accepted: Month 0000

2 Abstract:

3 • Using Portuguese diagnosis-related groups (DRGs) data we analysed the dynamics
4 of the COVID-19 hospitalisation in Portugal, with the ultimate goal of estimating
5 parameters to be used in COVID-19 mathematical modelling.

6 After processing the data, corresponding to the period between March 2020 and the
7 end of March 2021, an exploratory analysis was conducted, in which the length of
8 stay (LoS) in hospital care, the time until death (TuD), the number of admissions
9 and mortality count were estimated and described with respect to the age and gender
10 of the patients, but also by region of admission and evolution over time.

11 The median and mean of LoS was estimated to be 8 and 12.5 days (IQR:4-15) for non-
12 ICU patients and for ICU patients as 18 and 24.3 days (IQR:11-30). The percentage
13 of patients that, during their hospitalisation, required ICU care was determined to
14 be 10.9%. In-hospital mortality of non-ICU patients (22.6%) and of ICU patients
15 (32.8%) were also calculated. In a visual exploratory analysis we observed that they
16 changed with time, age group, gender and region.

17 Various univariable probability distributions were fitted to the LoS and TuD data,
18 using maximum likelihood estimation (MLE) but also maximum goodness-of-fit esti-
19 mation (MGE) to obtain the distribution parameters.

20 Key-Words:

21 • *COVID-19 hospitalisation; data analysis; model parametrisation; statistical distribu-*
22 *tions.*

23 AMS Subject Classification:

24 • 62-07,62E07,62F10,62P10

1. INTRODUCTION

In Portugal, the first case of COVID-19 was reported on March 2nd, 2020. On March 12, 2020 the Portuguese government declared the State of Emergency, and lockdown measures were implemented to stop the rapid increase of infections. From then onwards, several periods of confinement were implemented, according to the rise and fall in the number of cases and other epidemiological factors [2, 6].

Mathematical models are a suitable tool to develop scenarios to evaluate the impact of different control measures. For COVID-19, several mathematical models have been developed (see [11] and the references therein). The success of these models in capturing the transmission dynamics strongly depend on a good parametrisation. While it is expected that disease and biological parameters do not differ among different regions, other, as demographic and social related parameters, strongly depend on the population under study. Furthermore, there was a necessity for models to consider the hospitalisation dynamics of COVID-19, in order to assess health care burden. To this end, we analysed Portuguese DRG (diagnosis-related group) data with two main goals: to characterise hospital admissions and to estimate suitable parameters of interest.

In Portugal, the clinical data of hospitalized patients is logged according to the International Classification of Diseases, Clinical Modification and International Classification of Diseases, Tenth Revision Procedure Coding System (ICD-10-CM/PCS), in line with the 2020/2021 normative [3]. This unstructured data is then grouped into DRGs, with the objective of being used in healthcare management, pricing and decision-making by the hospital management and health professionals, but also by authorized national agencies that use it to perform various types of public health studies and statistical analysis on hospital treatment. The DRG system currently in use in Portugal is the *all-patient refined diagnosis-related groups* (hereafter APR-DRG) (*Administração Central do Sistema de Saúde* 2014). This system assigns codes to the patient which refers to a specific diagnosis, so that patients with similar resource usage and costs are clustered. According to Julio Souza *et.al* [19], the overall credibility of the Portuguese data is very high and the data can be considered credible and usable, despite some particular cases of diagnosis that have less cohesive data.

This paper is organized as follows: In Section 2 the original DRG data and the transformed data used in this study are explained as well as the description of the methodology used in the exploratory analysis and in the fitting of the distributions. In Section 3 the exploratory analysis is performed, in which the LoS and TuD, number of entries, fatalities and other parameters, for either ICU and non-ICU patients, is characterized and compared between genders, age groups and regions, as well as their evolution over time. In Section 4 the results from this study are discussed. An appendix section A is also provided, with an extended explanation of the data and all transformations carried out and also more extensive graphs and tables and other relevant information.

2. METHODS

2.1. Data

The data used in this study was obtained from the BI-MH platform of the SPMS/ACSS (Ministry of Health Shared Services / Central Administration of the Health System, IP) services of Portugal and is composed by the DRG data, with hospital discharges, having a COVID-19 diagnosis.

The DRG data was restructured in order to obtain the age of the patients and their respective age group, the regional health administration (ARS) to which the hospital belongs, LoS and TuD, as explained in the appendix A.1. This new data, containing 54685 entries, is described in appendix A.2, where the used variables are identified and explained. Because the DRG data is recorded only when a patient is discharged, a right truncation problem arises. In order to overcome this, we disregard admissions from the last two months, as explained in the appendix A.3. As such the considered data ranges are from 1 of March 2020 to 31 of March 2021. Additionally, from this restructured data, the admissions were aggregated by day. The type of hospitalisation is distinguished between:

- **Non-ICU patients:** patients who have not received treatment in intensive care units during their stay in the hospital;
- **ICU patients:** patients who received treatment intensive care units during their hospitalisation.

The regions used follow the areas of the regional health administrations (ARSs) established by the Portuguese national health service (SNS). The population numbers for each region were extracted from the Portuguese Central Administration of the Health Service (ACSS) website (data from June 2016 link1). The five considered ARSs that cover Continental Portugal are: ARS Norte, ARS Centro, ARS Lisboa e Vale do Tejo (LVT), ARS Alentejo and ARS Algarve. The Portuguese autonomous regions (Açores and Madeira) are not included in the study due to lack of data.

The population estimates (total and by gender) correspond to the resident population in Continental Portugal and were extracted from the census 2021 data, elaborated by *Statistics Portugal* (INE), the Portuguese official statistical institution, and is available at INE website at link2. The population numbers by age group correspond to the provisional resident population data obtained from the census data, available at the link3.

A table containing the number of hospital entries, the population number for Portugal and percentages calculated from these values was used in several

1 analysis, being presented in the appendix A.1 in table 3. The daily data is
2 presented in appendix A.2 tables 4,5,6,7 and 8.

3 Summary statistics complementary to the various Violin/box plots and
4 histograms are available in appendix A.4. Extra graphs of interest may be found
5 in appendix A.5. All data processing and analysis was done using *R* [14].

2.2. Distribution fitting

6 The probability distributions of the LoS and TuD, for non-ICU and ICU
7 patients, were also studied using the *R* *fitdistrplus* package [5]. The objective
8 of this analysis was to find the probability distribution that best describes the
9 observed data. To enable the use of the Weibull distributions the episodes with
10 0 days of stay were removed.

11 To choose the candidate distributions to be fitted to the data we used dif-
12 ferent tools of *fitdistrplus*. In addition to empirical plots, like density histograms
13 and the plots of the empirical cumulative distribution function, the descriptive
14 statistics provided helps in this choice, specially the Skewness-kurtosis plot (fig-
15 ure 11), also known as Cullen and Frey graph, with a nonparametric bootstrap
16 procedure to take into account the uncertainty in the estimation of kurtosis and
17 skewness from data.

18 The next step was to fit the chosen distributions to the data. The de-
19 fault method of estimation uses the maximum likelihood function, which was
20 used in all chosen distributions to be fitted. With the objective to give more
21 weight to the left tail of the distribution, as the data has many large outliers, the
22 maximum goodness-of-fit estimation using the left-tail Anderson-Darling (ADL)
23 distance was used in the two best fitted distributions, the gamma and log-normal
24 distributions.

25 To evaluate the fit of the candidate distribution, various goodness-of-fit
26 statistics (Kolmogorov-Smirnov, Cramer-von Mises and Anderson-Darling statis-
27 tic) and goodness-of-fit criteria (Aikake's and Bayesian information criterion) are
28 provided in the appendix A.4.

3. RESULTS

3.1. Exploratory data analysis

One of the most important parameters related to hospitalisation is the length of stay (LoS), as the rate of exits from hospital and the number of occupied beds are dependent on it. In figure 1 a violin/box plot, together with histograms, related to LoS for either type of hospitalisation are presented, obtaining a mean of 12.5 days and a median of 8 days (IQR:4-15) for non-ICU patients and a mean of 24.3 days and median of 18 days (IQR:11-30) for ICU patients (extended summary statistics available at appendix A.4 table 10). There is a clear difference in LoS between non-ICU and ICU patients.

Figure 1: Distribution of the LoS in hospital care. **(a)** Violin/box plot of the length of stay by type of hospitalisation, in which the white diamond represents the mean. **(b)** Distribution of the LoS for non-ICU and ICU patients (white diamonds represent the mean).

There are two possible hospitalisation outcomes, either the patient is discharged from hospital or he dies. Considering these outcomes a violin/box plot comparing time until death (TuD) and time until Discharge (TuDisc) is presented in figure 2, with summary statistics available in appendix A.4 in table 11. The TuD mean and median are 11.01 and 7 days for non-ICU, and 20.04 and 16 for ICU, respectively, and for TuDisc the mean and median are 12.98 and 9 for non-ICU and 26.39 and 19 days for ICU patient.

Three parameters that describe the hospitalisation dynamics were esti-

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 7

Figure 2: Violin/Box plot of the length of stay (LoS) for each outcome (Death or Discharge) and type of hospitalisation (non-ICU and ICU) (white diamonds represent the mean).

ated as follows:

$$\text{Percentage of ICU patients} = \frac{\text{ICU entries}}{\text{Total entries}} \times 100\% = 10.92\%$$

$$\text{Non-ICU COVID-19 fatality rate} = \frac{\text{non-ICU fatalities}}{\text{non-ICU entries}} \times 100\% = 22.60\%$$

$$\text{ICU COVID-19 fatality rate} = \frac{\text{ICU fatalities}}{\text{ICU entries}} \times 100\% = 32.84\%$$

- 1 In which the percentage of ICU patients represents the percentage of patients
- 2 that, at any point during hospital care, received ICU treatment, the non-ICU
- 3 COVID-19 fatality rate represents the percentage of patients that did not receive
- 4 any non-ICU care and that died in hospital care, and the ICU COVID-19 fatality
- 5 rate represents the percentage of patients that at any point during hospital care
- 6 received ICU treatment and died in hospital care.

3.1.1. Characterization over time

7 One of the characteristics of the COVID-19 pandemic is the peak in cases
8 that occurs in different periods of time. The number of hospitalizations follows
9 the number of cases and therefore is also prone to rapid peaks in hospitalisation.
10 This is specially problematic for health services since this could lead to a lack of
11 hospital capacity and affect the care to be provided.

12 We found an association between number of hospitalizations and the fatality
13 rate, the higher the number of hospital entries the higher the in-hospital fatality
14 rate and the decrease in the LoS of the patients that died, as seen in figure 3.
15 There is visible the increase in fatality rate with the increase in cases related to
16 the second surge in hospitalisation from September to February 2020.

17 Violin/box plots for either LoS and TuD over time are available in appendix
18 A.5, in figure 14 (a) and (b), linegraphs with hospital occupancy, discharges and

- 1 fatalities over time are also available in figure 15. Summary statistics for LoS
- 2 and TuD are displayed in appendix A.4 table 12 and 13.

Figure 3: (a)Entries, (b)percentage of ICU patients and (c)fatality rate, over time and by type of hospitalisation

3.1.2. Characterization by age

It is well known that COVID-19 affects more the elderly, increasing the risk of hospitalisation, severe illness and death. This was visible in the data as fatality rate increases with age, as shown in figure 5(b), as well as a higher percentage of hospitalized population (appendix A.5, figure 17(a) and (b)). The probability of being admitted to ICU was also higher for patients with 60 to 69 years of age, decreasing afterwards. This makes us conclude that the probability of entering ICU for older patients is lower.

The group from 10 to 15 years old is the one with the lower number of entries, but where the percentage of ICU patients and the fatality rate is higher than the 0 to 10 groups and the 15 to 20 group. This anomaly is not exclusive to Portugal and was reported by the Center of Disease Control and Prevention (CDC)[10].

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 9

Figure 4: Hospital entries by age: (a) entries and (b) percentage of patients requiring ICU treatment.

Figure 5: Mortality by age: (a) number of fatalities and (b) in-hospital fatality rate.

Figure 6: Hospital entries by age group and type of hospitalization.

3.1.3. Characterization by gender

1 Looking at gender differences in COVID-19 hospitalisation, there is a clear
2 difference in the total number hospitalisation, where males were more hospitalized
3 than females, with percentage of the population hospitalized in non-ICU of 0.51%

for males and 0.45% for females, and for ICU 0.08% for males and 0.03% for females. Male patients also have a higher fatality rate than females. For non-ICU, males patients have a 23.23% mortality rate and female patients have 21.95%, for ICU patients males have 33.81% and females 30.70 %.

Violin/box plots for the LoS considering gender and age groups are presented in appendix A.5 figure 20, where it is observable that the LoS are similar between group's gender, being the only exceptions the groups of the 25-30 and 30-35 for non-ICU patients were a large difference between genders arises. In the case of TuD there is also no observable difference, as seen in appendix A.5 figure 19.

In the Appendix A.5 figure 22 multiple graphs considering the differences between gender and type of hospitalization are presented. It is visible a lower number of entries and fatalities in females in general, with the females only being the group with more entries and fatalities in the 80 and over group for non-ICU patient type. Males have specially high numbers of entries and fatalities in ICU when compared to females.

Figure 7: Hospital entries by gender (F for female and M for male, white diamonds represent the mean). (a) Violin/box plot of the LoS by gender and type of hospitalisation, (b) total number of entries by gender and type of hospitalisation and (c) total number of fatalities by gender and type of hospitalisation, with fatality rates.

3.1.4. Characterization by region

The SARS-CoV2 geographical spread through Portugal was not uniform, with regions having increases and decreases in incidence in different periods, although the increase in cases in one region seems to spread to all other regions, as can be seen in appendix A.5 figure 25.

This leads to the differences in the percentage of the population admitted to hospital by region, observed in figure 8, with the *LVT* region being the one with the biggest percentage of hospitalization rate, followed by the *Norte* and *Centro* regions, other very populous regions. This could be explained by the higher population density of these regions that makes social distancing less efficient and by the transfer of patients from smaller hospitals located in less populated regions to larger ones, in big cities like Lisboa (*LVT* region), Porto (*Norte* region) and Coimbra (*Centro*).

No major difference was found between LoS and TuD by region, as seen in appendix A.5 figure 25. Looking at the percentage of patients requiring ICU treatment the *Centro* region has the lower and the *Algarve* region the higher percentage.

Figure 8: Number of (a)entries and (b)fatalities, by region and type of hospitalisation.

Figure 9

Figure 10: Percentage of ICU patients, by region.

3.2. Distribution fitting

An analysis of the distribution of the LoS was conducted as explained in section 2. Using the Cullen and Frey graphs (figure 11) we chose the distributions that best fitted each sample of the LoS type of hospitalisation, non-ICU and ICU. From the interpretations of those graphs the gamma, lognormal and Weibull distributions were the ones chosen to be fitted to the data. A table with the summary statistics used to plot these graphs is available in the appendix A.6 table 14.

After fitting the previously chosen distributions, for each type of hospitalisation (non-ICU and ICU), using the maximum likelihood estimation method, we compared them by using goodness-of-fit plots and the goodness of fit statistics and criteria supplied in the *fitdistrplus* package. We also estimated the gamma and log-normal distribution by fitting using the Left-tail AD variant of the Anderson-Darling distance. Tables with all the results are present in appendix A.6 (see 16 and 17).

The graphs in figure 12 represent all fitted theoretical distributions to the data for non-ICU and ICU patients. All estimated parameters are available in the appendix A.6 in table 15. For non-ICU patients the lognormal distribution (mean-log 2.1195 and sd-log 0.9496) was the distribution with the lower goodness-of-fit(GoF) statistics, except for the Anderson-Darling statistic that was lower in the lognormal ADL distribution (mean-log 2.1465 and sd-log 0.9886). The lognormal distribution (mean-log 2.8583 and sd-log 0.8655) also had lower GoF statistics for ICU patients, excepting for Kolmogorov-Smirnov and Anderson-Darling statistics that were lower for lognormal ADL distribution (mean-log

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 13

1 2.8986 and sd-log 0.8565).

Figure 11: Cullen and Frey graphs for (a)Non-ICU and (b)ICU patients.

Figure 12: Fitted distributions to the LoS for (a)non-ICU (Hosp) and (b)ICU patients.

2 The same analysis was carried out to the TuD, with the Cullen and Frey
3 graphs and statistics being presented in figure 26 and table 18, and a table with
4 the parameters for all fitted distribution in tables 19, in appendix A.6. Compar-

ing goodness of fit statistics, for the non-ICU patients the lognormal distribution (mean-log 1.9684 and sd-log 0.9873) had lower values for Anderson-Darling statistic and Alkaike's and Bayesian Information Criterion, with the gamma distribution (shape 1.2314 and rate 0.1088) having lower values in the Kolmogorov-Smirnov and Cramer-von Mises statistics. For ICU patients the gamma distribution (shape 1.5799 and rate 0.0779) had lower values for all but Kolmogorov-Smirnov and Cramer-von Mises statistic that were lower for the gamma ADL distribution (rate 1.7360 and rate 0.0888).

Figure 13: Fitted distributions to the time until death for (a)non-ICU and (b)ICU patients.

4. DISCUSSION

From the analysis of Portuguese DRG data the mean length of stay (LoS) was estimated as 12.5 (IQR: 4-15) for patients without ICU hospitalization and 24.3 days (IQR: 11-30) for patients who received ICU treatments, with it being higher in the case of recovery when comparing with in the case of hospital fatality. No meaningful differences were found in LoS between genders and regions, but differences were found between age groups and over time, with older groups having a higher LoS. The mean time until death (TuD) and time until discharged (TuDisc) was also estimated as 11.01 and 12.94 for non-ICU patients and 20.06 and 26.39 days for ICU patients.

The percentage of patients that required entry in ICU was estimated as 10.92% and the fatality rate as 22.60% for non-ICU patients and 32.84% for ICU, with the percentage of ICU patients changing depending in patient age and the region of hospitalization. The fatality rate was different between age group, gender and region, for either non-ICU and ICU patients.

Various distributions were fitted to the density plots of the LoS and TuD, for either non-ICU and ICU patients, with the estimated distributions parameters being presented in the Appendix A.6 in table 15 for LoS and table 15 for TuD. Log-normal and log-normal ADL have better goodness-of-fit statistics for LoS of non-ICU and ICU and for TuD of non-ICU, gamma and gamma ADL have better fits for TuD of ICU patients.

In [9], the authors consider various distributions that were fitted to the hospitalized patients clinical database of Belgium, with the length of stay for patients that recover being estimated as between 5 days (in the young population) to 15.7 (in the elderly) and the median length of stay for patients that die between 5.7 days (in the elderly) and 12.2 days (in the working age population). Our study distinguishes between non-ICU and ICU patients but the LoS, TuD and TuDisc follow the trends presented by [9], increasing with the age of the patient. This study also refers that the LoS is higher for males, what was not observed in our study. A lognormal distribution was used to describe the LoS in hospital (log-mean = 2.480 and log-sd = 0.913; IBM = 76,865) and in ICU (log-mean = 2.183 and log-sd = 1.052; IBM = 7312), the LoS for recovery and death was also analysed.

From a retrospective cohort study of Portuguese surveillance data, Peixoto[16] identifies a consistent increase in risk of ICU admission with age until the 70-79 age group, with decreased risk for subsequent age groups. In our study the risk increases until age 60-75 age group, decreasing afterward, as can be seen in Figure 4(b). In regards to mortality, Peixoto[16] claims a constant increase of risk of death with age, which is also visible in our data, specially in in Figure 5(b).

In her paper Vekaria [13] estimates the various LoS, for the UK, using three different statistical methods. With LoS for non-ICU patients being between 8.4 and 9.3 days and for ICU being between 16.2 and 35.2 days. The study also presents the time from hospital entry to ICU as being between 2 to 4 days and the time from ICU exit to full hospital release being between 1.8 and 8.4 days.

In Rees [15] paper a systematic review of early evidence on LoS of patients was conducted. This paper estimates the parameters from all published papers up to 12 April 2020. According to this the mean length of stay, for countries outside of China, is 11.0 days in hospital (non it ICU) and 12.2 in ICU. The LoS in hospitalisation, and the decrease in length of stay with time in the beginning of the pandemic reported in this study is also visible in our data.

One of the limitations of the data used is the lack of distinction between length of stay in ICU care and the overall time in hospital care for a patient that received ICU treatment, as the LoS refers to the total time in hospital. Unfortunately, there are not very many studies presenting LoS, as Rees [15] also report, LoS is normally not the primary measure of interest in studies which report it, and most of them consider LoS in ICU as time in ICU hospitalisation, what makes comparing them with our data difficult. Another limitation comes from the fact that the database BI-MH does not include hospitalizations in private hospitals or other non-public healthcare providers. Other factor to take in consideration, as it overestimates entries in hospital care, is that hospital transfers count as new entries and therefore a patient can have multiple consecutive entries.

Our objective with this study was to elucidate the COVID-19 hospitalization dynamics in Portugal. We took advantage of the DRG data to explore time in hospital care and other important parameters providing the necessary

1 tools and knowledge for the construction of better COVID-19 models that con-
2 sider hospitalization. The parameters here presented were already used in the
3 COVID-19 in-CTRL model [2].

ACKNOWLEDGMENTS

4 The authors acknowledge financial support from the Fundação para a Ciência
5 e Tecnologia—FCT through project 692 2^a edição Research 4 covid, project name
6 Projeção do Impacte das medidas Não-farmacológicas de Controlo e mitigação da
7 epidemia de COVID-19 em Tempo Real (COVID-19 in-CTRL). The first author
8 also acknowledges FCT within the grant with reference "COVID_inCTRL/DEP/01/2020".
9 The second author also acknowledges FCT within the PhD grants "DOCTOR-
10 ATES 4 COVID-19", number 2020.10172.BD. The third author also acknowl-
11 edges FCT within the PhD grant "DOCTORATES 4 COVID-19" with refer-
12 ence 2020.10252.BD. The fourth also acknowledges FCT within the Strategic
13 Project UIDB/00297/2020 (Centro de Matemática e Aplicações, FCT NOVA).
14 The fifth author also acknowledges FCT within projects UIDB/04621/2020 and
15 UIDP/04621/2020.

REFERENCES

- 16 [1] ANGELICO R.; TRAPANI S.; MANZIA, T. M.; LOMBARDINI, L.; TISONE G.
17 AND CARDILLO, M. (2020) The COVID-19 outbreak in Italy: Initial implications
18 for organ transplantation programs, *R Foundation for Statistical Computing* doi:
19 10.1111/ajt.15904.
- 20 [2] CAETANO C.; MORGADO M. L.; PATRÍCIO P.; PEREIRA J. F. AND NUNES B.
21 (2021) Mathematical Modelling of the Impact of Non-Pharmacological Strategies
22 to Control the COVID-19 Epidemic in Portugal, *Mathematics* **Vol. 9**, No. 10, p.
23 1084, doi: 10.3390/math9101084.
- 24 [3] CENTRO DE TERMINOLOGIAS CLÍNICAS (2020/2021) Catálogo Português
25 ICD-10-CM/PCS V5.0, 16-12-2020 link: [https://www.ctc.min-saude.pt/wp-](https://www.ctc.min-saude.pt/wp-content/uploads/2020/03/DocEsp_ICD10CMPCSV5.0.pdf)
26 [content/uploads/2020/03/DocEsp_ICD10CMPCSV5.0.pdf](https://www.ctc.min-saude.pt/wp-content/uploads/2020/03/DocEsp_ICD10CMPCSV5.0.pdf)
- 27 [4] CUCINOTTA D. AND VANELLI M. (2020) WHO Declares COVID-19 a
28 Pandemic *Acta Bio Medica Atenei Parmensis* **Vol. 91**, 157–160, doi:
29 10.23750/abm.v9i1.9397
- 30 [5] DELIGNETTE-MULLER M. L. AND DUTANG C. (2015) fitdistrplus: An R Package
31 for Fitting Distributions, *Journal of Statistical Software, Articles* **Vol. 64**, No. 4,
32 1–34, doi: 10.18637/jss.v064.i04.
- 33 [6] DIÁRIO DA REPÚBLICA ELETRÓNICO (2020) Legislação Compilada-COVID-19,
34 url: <https://dre.pt/legis/lacao-covid-19-upo>.

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 17

- 1 [7] DIRECTORATE-GENERAL OF HEALTH (PORTUGUESE: DIREÇÃO-GERAL DA
2 SAÚDE; DGS) (2021) Situation Report Nº 394 — 31/03/2021
- 3 [8] EUROSTAT - EUROPEAN COMMISSION (2020) Statistical regions in the European
4 Union and partner countries: NUTS and statistical regions 2021: 2020 edition,
5 *Publications Office* doi: 10.2785/72829
- 6 [9] FAES C.; ABRAMS S.; VAN BECKHOVEN D.; MEYFROIDT G.; Vlieghe E.;
7 HENS N. AND BELGIAN COLLABORATIVE GROUP ON COVID-19 HOSPITAL
8 SURVEILLANCE (2020) Time between Symptom Onset, Hospitalisation and Re-
9 recovery or Death: Statistical Analysis of Belgian COVID-19 Patients *International*
10 *Journal of Environmental Research and Public Health* **Vol. 17** No. 20, 7560, doi:
11 10.3390/ijerph17207560
- 12 [10] HAVERS F. *et al.* (2009) Hospitalization of Adolescents Aged 12–17 Years with
13 Laboratory-Confirmed COVID-19 — COVID-NET *MMWR. Morbidity and Mor-*
14 *tality Weekly Report*, **2021** No. 23, 851–857, doi: 10.15585/mmwr.mm7023e1
- 15 [11] HARJULE P.; TIWARI V. AND KUMAR A. (2021) Mathematical models to predict
16 COVID-19 outbreak : An interim review, *Journal of Interdisciplinary Mathematics*
17 **Vol. 24**, No. 2, 259-284, doi: 10.1080/09720502.2020.1848316.
- 18 [12] KRETZSCHMAR M.; WALLINGA J. (2009) Mathematical Models in Infectious Dis-
19 ease Epidemiology. In: Krämer A., Kretzschmar M., Krickeberg K. (eds) Modern
20 Infectious Disease Epidemiology *Statistics for Biology and Health*, 209–221, doi:
21 10.1007/978-0-387-93835-6_12
- 22 [13] VEKARIA, B., OVERTON, C., WIŚNIEWSKI, A. *et al.* (2021) COVID-19 length
23 of hospital stay: a systematic review and data synthesis *BMC Infect Dis* **Vol. 21**
24 No. 700, doi: 10.1186/s12879-021-06371-6
- 25 [14] R CORE TEAM (2020) R: A Language and Environment for Statistical Com-
26 puting, *American Journal of Transplantation* **Vol. 20**, No. 7, 1780–1784, url:
27 <https://www.R-project.org/>
- 28 [15] REES, E.M., NIGHTINGALE, E.S., JAFARI, Y. *et al.* (2020) Hospital Length of
29 Stay For COVID-19 Patients: Data-Driven Methods for Forward Planning *BMC*
30 *Med* **Vol. 18** No. 270, doi: 10.1186/s12916-020-01726-3
- 31 [16] RICOCA PEIXOTO V.; VIEIRA A.; AGUIAR P.; SOUSA P.; CARVALHO C.;
32 THOMAS D.; ABRANTES A. AND NUNES C. (2020) COVID-19 :Determinants of
33 Hospitalization, ICU and Death among 20,293 reported cases in Portugal *medRxiv*
34 *2020.05.29.20115824* doi: 10.1101/2020.05.29.20115824
- 35 [17] SARMADI M.; KAKHKI S.; FOROUGHI M.; ABADI T.S.H.; NAYYERI S.;
36 MOGHADAM V.K. AND RAMEZANI M. (2020) Hospitalization period of COVID-19
37 for future plans in hospital, *British Journal of Surgery* **Vol. 107**, No. 10 e427–e428,
38 doi: 10.1002/bjs.11871.
- 39 [18] SILVA C.J.; CRUZ C.,TORRES D.F.M.; MUÑUZURI A.P.; CARBALLOSA A.;
40 AREA I., NIETO J.J.; FONSECA-PINTO R.; PASSADOURO R.; DOS SANTOS E.S.;
41 ABREU W. AND MIRA J. (2021) Optimal control of the COVID-19 pandemic:
42 controlled sanitary deconfinement in Portugal *Scientific Reports* **Vol.11**, No. 1,
43 doi: 10.1038/s41598-021-83075-6
- 44 [19] SOUZA J.; PIMENTA D.; CABALLERO I. AND FREITAS A. (2020) Measuring
45 data credibility and medical coding: a case study using a nationwide Portuguese
46 inpatient database *Software Quality Journal* **Vol. 28** No. 3, 1043–1061, doi:
47 10.1007/s11219-020-09504-3

- 1 [20] Mangia C., Russo A., Civitelli S. and Gianicolo E. A. L. (2020). differenze
2 sesso/genere nella letalità COVid-19: cosa dicono e non dicono i dati *Epidemi-*
3 *ologia & Prevenzione*, No. 44(5-6 Suppl 2) 400–406 doi:10.19191/EP20.5-6.S2.145

A. APPENDIX

A.1. Data processing

1 The original Portuguese DRG data used in this was extracted from the BI-
 2 MH platform of the SPMS/ACSS services of Portugal and provided to us with
 3 the following structure:

Variable	Description
Health institution	Institution where the entries took place
Gender	Gender of the patient
Date of birth	Birth date of the patient
Hospital process	Anonymised data related to the hospital process number
Hospital episode	Anonymised data related to the hospital episode number
Destination after discharge	Destination of the patient after medical discharge, including death.
Discharge date	Date the patient was discharged from the hospital
Entry date	Date the patient entered hospital care
ICU	Whether or not the patient was in ICU care
Type of episode	Time of stay in hospital care groups (short term, normal, long term and prolonged evolution)
Diagnostic code	Diagnosis-related groups (DRG) code, in this case all entries have the U071 code, that identifies COVID-19 diagnosis.
Diagnostic	Diagnostic represented by the Diagnostic code, in this case all entries have the COVID-19 diagnostic
Diagnostic order	Order of diagnostic of the patient in that day

Table 1: Variable identification of the provided DRG data

4 The original DRG data, explained in the previous table 1, was transformed
 5 before use as to have the variable identified in table 2. The next list presents all
 6 modifications that were performed:

- 7 • From the "Health institution" column, all institutions were distributed

- 1 by the ARS (Regional Health Administration) to which they belong and
- 2 grouped in the "Region" column.
- 3 • From the "Date of birth" column the age of the patient at the time of entry
- 4 was determined and introduced in the "Age" column;
- 5 • The "ICU" column had the name changed to "Local" and the designations
- 6 changed, with "non-ICU" representing non-ICU patients and "ICU" for
- 7 patients that received any ICU care;
- 8 • A column with the length of stay was added, being calculated by subtracting
- 9 the entry date to the discharge date;
- 10 • Using the "Age" column, the patients were divided by age groups in a
- 11 column named "Group", with ages from 0 to 80 housed in 5 year gaps
- 12 groups and the 61 and up being grouped in a single group;
- 13 • Unused columns were removed in order to facilitate the analysis.

A.2. Variables identification

14 In table 2 the variables used in the study are explained. They were ob-
15 tained from the restructuring of the original DRG data, as is explained in section
16 A.1.

Variable	Description
Region	ARS of the health institution (Norte, Centro, LVT, Alentejo e Algarve)
Age	Age of the patient, at the time of entry
Gender	Gender of the patient
Entry date	Patient entry date to the health institution
Discharge date	Patient discharge date from the health institution
Destination after discharge	Destination after medical discharge, includes as a outcome death.
Length of stay	Length of stay in health care, corresponds to number of days between the discharge date and the entry date, in days
Group	Age group corresponding to the age of the patient

Table 2: Variable identification of the used data, restructured from the DRG data

17 The data obtained was grouped by gender, age group, time and region, for
18 either non-ICU and ICU patients, having obtained the table 3, with the entries

1 variable corresponding to the admission date and fatalities using the discharge
2 date.

3 The regions represent the ARS to which the hospital where the entry oc-
4 curred belongs to. With the population numbers used being relative to the
5 population that is covered by each region (link1). Data from the Portuguese
6 autonomous regions were not considered.

7 The total number of population by gender refers to the number of residents,
8 for Continental Portugal, according to the Portuguese 2021 census data. With
9 the total population being 9,857,593 people (link3). The number of individuals
10 for each age group corresponds to the annual mean resident population by age
11 group, according to the provisional resident population indicator, obtained from
12 the 2021 census. The sum of the population by group is 10 298 252, being slightly
13 lower than the, more up to date, total population relative to the 2021 preliminary
14 census data.

15 The Fatalities corresponds to the date of discharge for patients that died.
16 With the fatality rate being calculated as:

$$\text{Fatality rate} = \frac{\text{Fatalities}}{\text{Entries}} * 100\%,$$

17 and the percentage of ICU patients as:

$$\text{Percentage of ICU patients} = \frac{\text{ICU Entries}}{\text{Total Entries}} * 100\%.$$

	Entries			Percentage of Entries (%)			Population			Percentage of Population (%)			Fatalities			Fatality rate (%)			Percentage of ICU patients (%)		
	non-ICU	ICU	Total	non-ICU	ICU	Total	non-ICU	ICU	Total	non-ICU	ICU	Total	non-ICU	ICU	Total	non-ICU	ICU	Total	non-ICU	ICU	Total
Total	2380	1876	5256	48.576	31.413	40.925	510908	4097	5256	0.404	0.006	0.5548	523	576	579	22.058	32.836	23.723	10.0207		
Gender	2403	406	2809	51.1424	68.507	51.0475	465785	51	2809	0.5314	0.0874	0.6188	578	138	713	23.229	33.8135	24.7268	14.1108		
	Female	346	27	0.7103	0.4521	0.6821	415539	1	27	0.0833	0.0005	0.0838	1	1	2	0.289	3.7037	0.3302	7.2386		
Age Groups	[0,5]	373	17	0.3501	0.0351	0.3501	45341	0	17	0.0000	0.0000	0.0000	0	0	0	0.0000	0.0000	0.0000	0.0000		
	(5,10]	114	15	0.233	0.2512	0.2359	476290	0	15	0.024	0.0032	0.027	0	1	1	0	6.6667	0.752	11.6279		
	(10,15]	231	12	0.4742	0.2009	0.4444	506433	0	12	0.0456	0.0024	0.048	1	0	1	0.4329	0	0.4115	4.9883		
	(15,20]	471	25	0.9609	0.4186	0.907	531810	0	25	0.0886	0.0047	0.0933	1	0	1	0.2129	0	0.2016	5.0403		
	(20,25]	734	43	1.5068	0.72	1.4209	515282	0	43	0.1424	0.0083	0.1508	5	1	6	0.6812	2.3226	0.7722	5.3411		
	(25,30]	861	76	1.7675	1.2726	1.7134	532250	0	76	0.1615	0.0143	0.1757	8	7	15	0.9292	9.2105	1.0009	8.111		
	(30,35]	985	122	1.9707	2.0429	2.0243	615083	0	122	0.1901	0.0198	0.18	17	12	29	1.7299	9.8301	2.6197	11.0208		
	(35,40]	1301	217	2.6707	3.6336	2.7759	724463	0	217	0.1796	0.03	0.2005	28	24	52	2.1322	11.0399	3.4296	14.2901		
	(40,45]	1745	326	3.5811	5.1411	4.3661	956702	0	326	0.2407	0.0437	0.2844	46	34	80	2.8343	15.6707	5.1411	17.4157		
	(45,50]	2125	440	4.2921	7.4682	4.7015	790701	0	440	0.2408	0.0529	0.2938	49	70	119	3.2924	15.6007	6.3013	18.5501		
Month	(50,55]	2869	640	5.7604	10.7167	6.3015	790701	0	640	0.3078	0.0906	0.4884	162	137	309	5.7672	22.9088	8.0591	18.5501		
	(55,60]	3497	854	7.1788	14.301	7.5955	655391	0	854	0.5336	0.1303	0.6639	287	222	509	8.207	25.9953	11.6985	19.0277		
	(60,65]	4142	1002	8.5029	16.7783	9.4066	602210	0	1002	0.6878	0.1664	0.8542	505	369	874	12.1922	36.8263	16.9607	19.479		
	(65,70]	4711	902	9.6709	15.1038	10.2642	536388	0	902	0.8783	0.1682	1.0464	824	389	1213	17.401	43.1264	21.6105	16.0698		
	(70,75]	5595	696	11.4856	11.6544	11.5041	428706	0	696	1.3051	0.1623	1.4674	1472	361	1833	26.3002	51.8678	29.1369	11.0634		
	(75,80]	18469	564	35.9403	9.4441	35.7191	661224	0	564	2.8088	0.0853	2.9541	7540	920	7860	39.7491	56.7376	40.2996	2.8874		
	(80,85]	1631	373	3.3482	0.2458	3.6046	3857593	0	373	0.0165	0.0038	0.0203	45	66	111	21.1327	28.4182	25.305	18.0128		
	(85,90]	1434	258	2.9587	0.4961	3.4548	346616	0	258	0.0146	0.0031	0.0177	32	72	104	14.6667	22.6667	18.6667	10.3219		
	(90,95]	1083	125	2.188	0.2822	2.4931	21724	0	125	0.0108	0.0013	0.0121	149	26	175	14.0169	20.8	14.7929	12.3415		
	(95,100]	1037	146	2.1288	2.4447	2.1623	9857593	0	146	0.0105	0.0015	0.012	141	34	175	13.5969	23.2877	14.7929	12.3415		
Regions	Alentejo	859	109	1.7634	1.8252	1.7701	9857593	0	109	0.0087	0.0011	0.0098	111	23	134	12.922	21.009	13.843	11.2603		
	Azores	1363	233	2.798	3.9015	2.9185	9857593	0	233	0.0138	0.0024	0.0162	233	64	297	17.0946	27.4678	18.699	14.7411		
	Braga	642	111	1.3179	1.8587	1.377	9857593	0	111	0.0087	0.0011	0.0098	99	29	128	15.4206	26.1201	16.9887	14.7411		
	Castella	3832	586	7.8405	9.8125	8.079	9857593	0	586	0.0389	0.0099	0.0488	776	164	940	20.2305	27.9803	21.2766	13.2639		
	Coimbra	1740	975	3.4897	13.7602	5.5067	9857593	0	975	0.0783	0.0099	0.0884	172	347	519	22.1189	35.5807	25.0259	11.1876		
	Lisboa	7740	1574	15.8607	19.3742	15.3674	9857593	0	1574	0.1574	0.0144	0.1718	1574	352	1926	15.7406	25.4541	20.5944	15.7406		
	LVT	13294	1415	27.2905	23.6839	26.8077	9857593	0	1415	0.1349	0.0144	0.1492	3032	549	3581	27.2906	38.7986	28.4248	9.072		
	Norte	6191	658	12.7091	11.0181	12.5245	9857593	0	658	0.0628	0.0067	0.0695	1331	205	1536	21.499	31.155	22.4266	10.672		
	Portugal	1133	138	2.3239	2.3108	2.3242	9857593	0	138	0.0115	0.0014	0.0129	123	52	175	10.8561	37.6812	13.7687	10.8576		
	Setúbal	1706	256	3.4021	4.2867	3.5587	106726	0	256	0.0232	0.1535	1.1768	556	86	642	32.5909	33.5938	32.7217	13.0479		
Norte	Açores	1189	243	2.4408	4.069	2.6186	451006	0	243	0.0266	0.0539	0.3175	273	102	375	22.9005	41.9753	26.1872	16.0693		
	Algarve	7681	599	15.7679	10.0301	15.1413	1480664	0	599	0.5188	0.0405	0.5592	2600	232	2992	26.8194	38.7312	27.6812	7.2843		
	LVT	19624	2406	40.2849	40.288	40.2853	3658671	0	2406	0.5802	0.0657	0.6019	4433	845	5278	22.8897	35.1205	25.4652	10.9215		
	Norte	18513	2408	35.0032	41.3262	38.3967	3128844	0	2408	0.5253	0.079	0.6712	3689	686	4385	19.265	28.201	20.8699	11.763		

Table 3: Table with various data in relation to gender, age group, time and region.

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 23

Table 4: Table with daily admissions (Part I)

Date	Entries	Discharges	non-ICU entries	non-ICU discharges	Non-ICU in hospital	ICU entries	en-ICU exits	ICU in hospital	Non-ICU deaths	ICU deaths	sum non-ICU deaths	sum ICU deaths
2020-03-01	5	0	5	0	5	0	0	0	0	0	0	0
2020-03-02	7	0	7	0	12	0	0	0	0	0	0	0
2020-03-03	12	1	11	1	22	1	0	1	0	0	0	0
2020-03-04	13	0	12	0	34	1	0	2	0	0	0	0
2020-03-05	14	0	8	0	42	6	0	8	0	0	0	0
2020-03-06	13	1	13	1	54	0	0	8	0	0	0	0
2020-03-07	16	0	15	0	69	1	0	9	0	0	0	0
2020-03-08	16	1	15	1	83	1	0	10	0	0	0	0
2020-03-09	18	1	16	1	98	2	0	12	0	0	0	0
2020-03-10	20	0	16	0	114	4	0	16	0	0	0	0
2020-03-11	25	2	20	2	132	5	0	21	0	0	0	0
2020-03-12	23	9	19	9	142	4	0	25	0	0	0	0
2020-03-13	29	11	26	11	157	3	0	28	0	0	0	0
2020-03-14	31	4	25	4	178	6	0	34	0	0	0	0
2020-03-15	26	3	15	3	190	11	0	45	0	0	0	0
2020-03-16	37	7	30	6	214	7	1	51	0	1	0	1
2020-03-17	44	10	33	9	238	11	1	61	1	0	1	1
2020-03-18	53	9	40	8	270	13	1	73	0	1	1	2
2020-03-19	65	14	51	13	308	14	1	86	3	0	4	2
2020-03-20	68	20	49	19	338	19	1	104	4	1	8	3
2020-03-21	95	19	76	19	395	19	0	123	4	0	12	3
2020-03-22	97	12	78	12	461	19	0	142	6	0	18	3
2020-03-23	105	19	85	18	528	20	1	161	5	0	23	3
2020-03-24	119	31	92	26	594	27	5	183	5	5	28	8
2020-03-25	142	31	116	28	682	26	3	206	11	3	39	11
2020-03-26	136	30	113	27	768	23	3	226	6	2	45	13
2020-03-27	145	53	116	45	839	29	8	247	15	5	60	18
2020-03-28	172	41	145	39	945	27	2	272	14	1	74	19
2020-03-29	131	38	109	35	1019	22	3	291	15	2	89	21
2020-03-30	157	72	136	65	1090	21	7	305	23	4	112	25
2020-03-31	170	85	139	80	1149	31	5	331	21	3	133	28
2020-04-01	152	76	137	70	1216	15	6	340	18	3	151	31
2020-04-02	163	92	138	72	1282	25	20	345	21	8	172	39
2020-04-03	129	97	109	84	1307	20	13	352	13	3	185	42
2020-04-04	134	66	115	60	1362	19	6	365	24	4	209	46
2020-04-05	127	60	118	51	1429	9	9	365	16	5	225	51
2020-04-06	125	110	113	98	1444	12	12	365	24	3	249	54
2020-04-07	114	137	105	124	1425	9	13	361	28	5	277	59
2020-04-08	122	120	111	112	1424	11	8	364	19	3	296	62
2020-04-09	88	97	77	86	1415	11	11	364	22	3	318	65
2020-04-10	107	102	94	93	1416	13	9	368	32	1	350	66
2020-04-11	102	70	92	60	1448	10	10	368	23	1	373	67
2020-04-12	90	58	79	49	1478	11	9	370	24	6	397	73
2020-04-13	90	109	80	96	1462	10	13	367	25	5	422	78
2020-04-14	116	128	107	113	1456	9	15	361	28	4	450	82
2020-04-15	103	125	96	110	1442	7	15	353	26	2	476	84
2020-04-16	94	111	87	97	1432	7	14	346	23	5	499	89
2020-04-17	106	129	94	113	1413	12	16	342	24	7	523	96
2020-04-18	75	72	67	60	1420	8	12	338	23	4	546	100
2020-04-19	55	66	51	59	1412	4	7	335	19	2	565	102
2020-04-20	80	113	73	96	1389	7	17	325	25	2	590	104
2020-04-21	61	99	55	92	1352	6	7	324	20	3	610	107
2020-04-22	58	105	52	89	1315	6	16	314	21	3	631	110
2020-04-23	62	98	55	86	1284	7	12	309	9	3	640	113
2020-04-24	67	110	57	93	1248	10	17	302	21	4	661	117
2020-04-25	61	55	53	49	1252	8	6	304	17	1	678	118
2020-04-26	47	52	45	46	1251	2	6	300	18	2	696	120
2020-04-27	42	93	38	83	1206	4	10	294	12	4	708	124
2020-04-28	38	99	36	83	1159	2	16	280	14	5	722	129
2020-04-29	45	85	42	76	1125	3	9	274	9	1	731	130
2020-04-30	52	85	45	79	1091	7	6	275	17	1	748	131
2020-05-01	40	60	36	51	1076	4	9	270	11	0	759	131
2020-05-02	40	52	36	45	1067	4	7	267	10	5	769	136
2020-05-03	34	40	33	40	1060	1	0	268	17	0	786	136
2020-05-04	49	104	41	91	1010	8	13	263	9	2	795	138
2020-05-05	33	106	27	91	946	6	15	254	10	1	805	139
2020-05-06	41	86	39	69	916	2	17	239	10	0	815	139
2020-05-07	52	66	51	61	906	1	5	235	10	0	825	139
2020-05-08	29	88	25	68	863	4	20	219	7	3	832	142
2020-05-09	39	40	33	34	862	6	6	219	5	2	837	144
2020-05-10	34	26	30	22	870	4	4	219	9	3	846	147
2020-05-11	34	83	25	75	820	9	8	220	9	1	855	148
2020-05-12	38	67	35	56	799	3	11	212	7	2	862	150
2020-05-13	50	81	41	70	770	9	11	210	2	2	864	152
2020-05-14	49	73	46	60	756	3	13	200	4	2	868	154
2020-05-15	48	56	44	50	750	4	6	198	3	0	871	154
2020-05-16	37	38	31	34	747	6	4	200	9	2	880	156
2020-05-17	33	22	30	21	756	3	1	202	8	1	888	157
2020-05-18	25	74	24	66	714	1	8	195	3	3	891	160
2020-05-19	31	65	27	55	686	4	10	189	9	1	900	161
2020-05-20	44	72	39	62	663	5	10	184	7	0	907	161
2020-05-21	21	42	20	35	648	1	7	178	3	1	910	162
2020-05-22	36	83	34	70	612	2	13	167	1	3	911	165
2020-05-23	37	35	31	30	613	6	5	168	7	2	918	167
2020-05-24	34	15	28	12	629	6	3	171	3	2	921	169
2020-05-25	33	57	30	46	613	3	11	163	2	1	923	170
2020-05-26	39	46	37	37	613	2	9	156	5	2	928	172
2020-05-27	66	66	61	62	612	5	4	157	7	0	935	172
2020-05-28	55	60	49	55	606	6	5	158	6	2	941	174
2020-05-29	38	73	36	65	577	2	8	152	9	1	950	175
2020-05-30	33	25	30	25	582	3	0	155	5	0	955	175
2020-05-31	16	18	14	18	578	2	0	157	5	0	960	175

Table 5: Table with daily admissions (Part II)

Date	Entries	Discharges	non-ICU entries	non-ICU discharges	Non-ICU in hospital	ICU entries	ICU exits	ICU in hospital	Non-ICU deaths	ICU deaths	sum non-ICU deaths	sum ICU deaths
2020-06-01	49	54	41	47	572	8	7	158	9	2	969	177
2020-06-02	40	45	36	40	568	4	5	157	1	0	970	177
2020-06-03	42	51	38	44	562	4	7	154	4	2	974	179
2020-06-04	40	56	38	49	551	2	7	149	5	0	979	179
2020-06-05	35	65	31	56	526	4	9	144	11	1	990	180
2020-06-06	27	32	26	30	522	1	2	143	3	1	993	181
2020-06-07	28	18	21	17	526	7	1	149	4	0	997	181
2020-06-08	35	43	29	39	516	6	4	151	4	0	1001	181
2020-06-09	37	45	33	43	506	4	2	153	4	0	1005	181
2020-06-10	26	31	25	26	505	1	5	149	7	1	1012	182
2020-06-11	35	7	29	7	527	6	0	155	0	0	1012	182
2020-06-12	40	64	36	58	505	4	6	153	3	5	1015	187
2020-06-13	39	22	35	20	520	4	2	155	2	2	1017	189
2020-06-14	23	16	21	15	526	2	1	156	0	0	1017	189
2020-06-15	51	63	43	50	519	8	13	151	6	1	1023	190
2020-06-16	29	41	26	38	507	3	3	151	2	1	1025	191
2020-06-17	36	57	31	45	493	5	12	144	5	2	1030	193
2020-06-18	46	46	42	40	495	4	6	142	7	1	1037	194
2020-06-19	46	62	42	51	486	4	11	135	1	0	1038	194
2020-06-20	47	23	41	19	508	6	4	137	5	1	1043	195
2020-06-21	39	9	34	8	534	5	1	141	3	1	1046	196
2020-06-22	53	64	48	53	529	5	11	135	3	3	1049	199
2020-06-23	53	53	47	46	530	6	7	134	3	1	1052	200
2020-06-24	53	42	45	39	536	8	3	139	5	2	1057	202
2020-06-25	46	51	39	47	528	7	4	142	5	1	1062	203
2020-06-26	45	52	38	45	521	7	7	142	3	3	1065	206
2020-06-27	38	25	35	23	533	3	2	143	6	0	1071	206
2020-06-28	37	13	31	12	552	6	1	148	4	0	1075	206
2020-06-29	37	62	29	54	527	8	8	148	7	2	1082	208
2020-06-30	31	36	27	32	522	4	4	148	5	0	1087	208
2020-07-01	39	59	34	50	506	5	9	144	5	3	1092	211
2020-07-02	33	46	31	42	495	2	4	142	7	4	1099	215
2020-07-03	51	65	45	61	479	6	4	144	8	1	1107	216
2020-07-04	37	23	33	18	494	4	5	143	7	1	1114	217
2020-07-05	37	22	31	19	506	6	3	146	5	2	1119	219
2020-07-06	44	59	39	54	491	5	5	146	7	1	1126	220
2020-07-07	39	37	35	31	495	4	6	144	5	3	1131	223
2020-07-08	40	58	37	48	484	3	10	137	11	2	1142	225
2020-07-09	35	52	32	46	470	3	6	134	3	0	1145	225
2020-07-10	40	45	34	38	466	6	7	133	7	2	1152	227
2020-07-11	23	25	23	18	471	0	7	126	5	5	1157	232
2020-07-12	25	8	21	8	484	4	0	130	1	0	1158	232
2020-07-13	36	38	32	29	487	4	9	125	4	2	1162	234
2020-07-14	30	42	29	39	477	1	3	123	6	0	1168	234
2020-07-15	36	39	30	34	473	6	5	124	4	0	1172	234
2020-07-16	29	37	25	31	467	4	6	122	3	0	1175	234
2020-07-17	30	51	24	42	449	6	9	119	3	0	1178	234
2020-07-18	26	20	24	17	456	2	3	118	5	1	1183	235
2020-07-19	28	13	24	12	468	4	1	121	4	1	1187	236
2020-07-20	32	44	27	39	456	5	5	121	4	3	1191	239
2020-07-21	29	39	27	31	452	2	8	115	5	2	1196	241
2020-07-22	27	33	25	30	447	2	3	114	3	1	1199	242
2020-07-23	33	52	29	43	433	4	9	109	5	2	1204	244
2020-07-24	32	47	28	39	422	4	8	105	3	0	1207	244
2020-07-25	24	18	20	16	426	4	2	107	3	0	1210	244
2020-07-26	28	12	28	10	444	0	2	105	2	1	1212	245
2020-07-27	19	34	18	32	430	1	2	104	2	0	1214	245
2020-07-28	23	30	19	26	423	4	4	104	2	1	1216	246
2020-07-29	22	35	19	30	412	3	5	102	4	0	1220	246
2020-07-30	21	42	18	36	394	3	6	99	5	3	1225	249
2020-07-31	20	58	18	45	367	2	13	88	1	1	1226	250
2020-08-01	22	9	18	9	376	4	0	92	1	0	1227	250
2020-08-02	22	6	21	4	393	1	2	91	1	2	1228	252
2020-08-03	15	28	12	24	381	3	4	90	1	0	1229	252
2020-08-04	16	29	13	26	368	3	3	90	3	0	1232	252
2020-08-05	20	27	18	23	363	2	4	88	4	0	1236	252
2020-08-06	21	32	19	27	355	2	5	85	6	1	1242	253
2020-08-07	43	49	42	41	356	1	8	78	1	3	1243	256
2020-08-08	23	16	21	15	362	2	1	79	4	1	1247	257
2020-08-09	24	12	21	10	373	3	2	80	4	1	1251	258
2020-08-10	29	37	24	36	361	5	1	84	2	0	1253	258
2020-08-11	29	19	25	17	369	4	2	86	3	0	1256	258
2020-08-12	16	25	15	20	364	1	5	82	4	0	1260	258
2020-08-13	13	27	9	22	351	4	5	81	5	0	1265	258
2020-08-14	18	27	15	24	342	3	3	81	3	0	1268	258
2020-08-15	25	13	18	12	348	7	1	87	4	1	1272	259
2020-08-16	20	6	18	6	360	2	0	89	2	0	1274	259
2020-08-17	27	33	22	25	357	5	8	86	4	2	1278	261
2020-08-18	20	28	20	26	351	0	2	84	1	1	1279	262
2020-08-19	29	29	21	25	347	8	4	88	4	0	1283	262
2020-08-20	25	32	22	30	339	3	2	89	2	1	1285	263
2020-08-21	22	14	19	12	346	3	2	90	2	0	1287	263
2020-08-22	20	12	18	11	353	2	1	91	4	0	1291	263
2020-08-23	25	17	18	14	357	7	3	95	4	2	1295	265
2020-08-24	23	24	20	19	358	3	5	93	3	2	1298	267
2020-08-25	25	34	21	30	349	4	4	93	8	0	1306	267
2020-08-26	31	37	27	35	341	4	2	95	2	0	1308	267
2020-08-27	36	25	28	19	350	8	6	97	4	1	1312	268
2020-08-28	32	42	27	35	342	5	7	95	1	0	1313	268
2020-08-29	30	12	26	12	356	4	0	99	3	0	1316	268
2020-08-30	23	12	20	12	364	3	0	102	4	0	1320	268
2020-08-31	29	30	24	25	363	5	5	102	2	0	1322	268

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 25

Table 6: Table with daily admissions (Pare III)

Date	Entries	Discharges	non-ICU entries	non-ICU discharges	Non-ICU in hospital	ICU entries	ICU exits	ICU in hospital	Non-ICU deaths	ICU deaths	sum non-ICU deaths	sum ICU deaths
2020-09-01	34	23	32	16	379	2	7	97	2	2	1324	270
2020-09-02	44	29	35	25	389	9	4	102	1	1	1325	271
2020-09-03	47	38	39	36	392	8	2	108	4	1	1329	272
2020-09-04	37	29	30	25	397	7	4	111	4	1	1333	273
2020-09-05	43	16	39	15	421	4	1	114	3	0	1336	273
2020-09-06	38	10	28	9	440	10	1	123	3	1	1339	274
2020-09-07	47	37	46	31	455	1	6	118	4	0	1343	274
2020-09-08	46	38	35	30	460	11	8	121	2	0	1345	274
2020-09-09	50	24	44	24	480	6	0	127	3	0	1348	274
2020-09-10	49	35	42	31	491	7	4	130	6	2	1354	276
2020-09-11	53	33	48	29	510	5	4	131	4	0	1358	276
2020-09-12	46	24	39	23	526	7	1	137	5	1	1363	277
2020-09-13	49	11	41	10	557	8	1	144	4	1	1367	278
2020-09-14	49	45	39	38	558	10	7	147	2	2	1369	280
2020-09-15	51	42	42	37	563	9	5	151	4	1	1373	281
2020-09-16	51	54	49	46	566	2	8	145	9	2	1382	283
2020-09-17	49	44	42	41	567	7	3	149	5	1	1387	284
2020-09-18	62	48	55	43	579	7	5	151	3	2	1390	286
2020-09-19	55	31	49	27	601	6	4	153	10	2	1400	288
2020-09-20	49	19	41	15	627	8	4	157	3	3	1403	291
2020-09-21	67	47	62	41	648	5	6	156	6	1	1409	292
2020-09-22	68	53	58	47	659	10	6	160	4	0	1413	292
2020-09-23	76	44	64	41	682	12	3	169	5	1	1418	293
2020-09-24	71	36	59	34	707	12	2	179	6	0	1424	293
2020-09-25	60	70	51	63	695	9	7	181	7	1	1431	294
2020-09-26	52	26	44	25	714	8	1	188	6	0	1437	294
2020-09-27	65	21	53	17	750	12	4	196	4	1	1441	295
2020-09-28	54	42	46	37	759	8	5	199	7	1	1448	296
2020-09-29	65	65	51	57	753	14	8	205	10	1	1458	297
2020-09-30	69	50	60	46	767	9	4	210	4	0	1462	297
2020-10-01	68	59	58	50	775	10	9	211	6	2	1468	299
2020-10-02	64	65	53	58	770	11	7	215	13	2	1481	301
2020-10-03	65	43	61	39	792	4	4	215	13	2	1494	303
2020-10-04	76	32	69	30	831	7	2	220	7	1	1501	304
2020-10-05	91	37	78	28	881	13	9	224	10	5	1511	309
2020-10-06	96	52	81	46	916	15	6	233	7	3	1518	312
2020-10-07	113	78	97	71	942	16	7	242	7	2	1525	314
2020-10-08	84	77	73	69	946	11	8	245	11	1	1536	315
2020-10-09	122	91	110	79	977	12	12	245	3	5	1539	320
2020-10-10	87	31	79	25	1031	8	6	247	11	2	1550	322
2020-10-11	74	31	57	25	1063	17	6	258	12	2	1562	324
2020-10-12	98	86	81	73	1071	17	13	262	16	3	1578	327
2020-10-13	105	80	95	74	1092	10	6	266	7	0	1585	327
2020-10-14	129	109	112	93	1111	17	16	267	12	1	1597	328
2020-10-15	127	125	109	110	1110	18	15	270	23	3	1620	331
2020-10-16	132	101	115	88	1137	17	13	274	12	6	1632	337
2020-10-17	131	46	110	41	1206	21	5	290	17	2	1649	339
2020-10-18	139	37	118	33	1291	21	4	307	15	2	1664	341
2020-10-19	156	123	128	111	1308	28	12	323	14	4	1678	345
2020-10-20	183	114	160	103	1365	23	11	335	14	1	1692	346
2020-10-21	174	114	148	102	1411	26	12	349	13	4	1705	350
2020-10-22	189	138	166	119	1458	23	19	353	31	4	1736	354
2020-10-23	203	144	185	122	1521	18	22	349	18	6	1754	360
2020-10-24	184	72	155	65	1611	29	7	371	22	2	1776	362
2020-10-25	151	57	135	52	1694	16	5	382	26	4	1802	366
2020-10-26	240	165	206	152	1748	34	13	403	19	4	1821	370
2020-10-27	198	175	176	154	1770	22	21	404	22	5	1843	375
2020-10-28	208	172	176	156	1790	32	16	420	28	4	1871	379
2020-10-29	222	182	192	164	1818	30	18	432	39	5	1910	384
2020-10-30	260	211	228	196	1850	32	15	449	32	4	1942	388
2020-10-31	249	94	221	83	1988	28	11	466	29	4	1971	392
2020-11-01	240	82	199	69	2118	41	13	494	37	9	2008	401
2020-11-02	226	200	198	184	2132	28	16	506	37	3	2045	404
2020-11-03	234	250	212	216	2128	22	34	494	46	7	2091	411
2020-11-04	310	251	278	235	2171	32	16	510	39	3	2130	414
2020-11-05	330	258	289	230	2230	41	28	523	36	10	2166	424
2020-11-06	262	250	232	228	2234	30	22	531	43	5	2209	429
2020-11-07	281	102	244	96	2382	37	6	562	38	5	2247	434
2020-11-08	251	112	216	101	2497	35	11	586	44	7	2291	441
2020-11-09	272	256	252	224	2525	20	32	574	37	16	2328	457
2020-11-10	262	277	234	240	2510	28	28	574	54	8	2382	465
2020-11-11	344	283	313	254	2569	31	29	576	56	9	2438	474
2020-11-12	299	287	265	258	2576	34	29	581	47	12	2485	486
2020-11-13	293	284	263	260	2579	30	24	587	43	1	2528	487
2020-11-14	318	127	286	104	2761	32	23	596	48	14	2576	501
2020-11-15	262	155	223	139	2845	39	16	619	66	11	2642	512
2020-11-16	292	310	260	289	2816	32	21	630	54	6	2696	518
2020-11-17	289	331	262	298	2780	27	33	624	60	10	2756	528
2020-11-18	319	303	270	280	2770	49	23	650	59	6	2815	534
2020-11-19	324	276	291	248	2813	33	28	655	36	8	2851	542
2020-11-20	337	349	303	319	2797	34	30	659	52	5	2903	547
2020-11-21	289	125	271	112	2956	18	13	664	49	6	2952	553
2020-11-22	306	144	254	126	3084	52	18	698	59	9	3011	562
2020-11-23	289	323	261	297	3048	28	26	700	64	5	3075	567
2020-11-24	308	314	279	287	3040	29	27	702	49	11	3124	578
2020-11-25	320	313	281	283	3038	39	30	711	53	14	3177	592
2020-11-26	318	288	285	265	3058	33	23	721	44	8	3221	600
2020-11-27	302	387	266	341	2983	36	46	711	53	14	3274	614
2020-11-28	310	133	273	119	3137	37	14	734	41	9	3315	623
2020-11-29	250	138	228	119	3246	22	19	737	55	12	3370	635
2020-11-30	278	363	252	326	3172	26	37	726	51	11	3421	646

Table 7: Table with dmissions aggregataed by day (Part IV)

Date	Entries	Discharges	non-ICU entries	non-ICU discharges	Non-ICU in hospital	ICU entries	en-ICU exits	ICU in hospital	Non-ICU deaths	ICU deaths	sum non-ICU deaths	sum ICU deaths
2020-12-01	293	156	256	146	3282	37	10	753	61	5	3482	651
2020-12-02	290	349	262	312	3232	28	37	744	52	10	3534	661
2020-12-03	273	366	246	321	3157	27	45	726	48	5	3582	666
2020-12-04	334	372	301	339	3119	33	33	726	55	11	3637	677
2020-12-05	259	168	231	152	3198	28	16	738	60	8	3697	685
2020-12-06	255	148	235	132	3301	20	16	742	61	12	3758	697
2020-12-07	268	397	249	347	3203	19	50	711	55	8	3813	705
2020-12-08	266	168	234	151	3286	32	17	726	59	10	3872	715
2020-12-09	281	394	254	356	3184	27	38	715	67	7	3939	722
2020-12-10	236	356	209	323	3070	27	33	709	78	7	4017	729
2020-12-11	257	403	234	353	2951	23	50	682	55	14	4072	743
2020-12-12	249	166	231	150	3032	18	16	684	77	11	4149	754
2020-12-13	251	134	227	120	3139	24	14	694	64	10	4213	764
2020-12-14	249	321	224	286	3077	25	35	684	60	12	4273	776
2020-12-15	273	320	244	288	3033	29	32	681	57	11	4330	787
2020-12-16	272	317	248	282	2999	24	35	670	59	10	4389	797
2020-12-17	226	287	200	255	2944	26	32	664	51	11	4440	808
2020-12-18	268	339	241	308	2877	27	31	660	58	6	4498	814
2020-12-19	214	154	197	138	2936	17	16	661	58	10	4556	824
2020-12-20	261	101	232	87	3081	29	14	676	43	5	4599	829
2020-12-21	226	356	196	317	2960	30	39	667	48	8	4647	837
2020-12-22	294	372	264	336	2888	30	36	661	58	12	4705	849
2020-12-23	297	373	279	334	2833	18	39	640	48	6	4753	855
2020-12-24	232	294	199	267	2765	33	27	646	50	6	4803	861
2020-12-25	221	111	203	101	2867	18	10	654	49	8	4852	869
2020-12-26	264	117	231	101	2997	33	16	671	49	10	4901	879
2020-12-27	250	109	228	95	3130	22	14	679	46	12	4947	891
2020-12-28	277	295	250	262	3118	27	33	673	44	17	4991	908
2020-12-29	337	323	306	304	3120	31	19	685	51	5	5042	913
2020-12-30	291	332	271	293	3098	20	39	666	49	17	5091	930
2020-12-31	362	366	325	324	3099	37	42	661	42	12	5133	942
2021-01-01	302	145	280	133	3246	22	12	671	50	11	5183	953
2021-01-02	299	138	272	122	3396	27	16	682	53	13	5236	966
2021-01-03	343	118	307	102	3601	36	16	702	44	13	5280	979
2021-01-04	335	312	294	280	3615	41	32	711	58	12	5338	991
2021-01-05	369	335	324	295	3644	45	40	716	61	11	5399	1002
2021-01-06	389	331	354	293	3705	35	38	713	67	10	5466	1012
2021-01-07	409	351	354	318	3741	55	33	735	73	12	5539	1024
2021-01-08	473	390	418	349	3810	55	41	749	72	10	5611	1034
2021-01-09	431	166	378	149	4039	53	17	785	74	12	5685	1046
2021-01-10	389	167	350	149	4240	39	18	806	79	13	5764	1059
2021-01-11	404	399	356	352	4244	48	47	807	92	18	5856	1077
2021-01-12	496	414	450	374	4320	46	40	813	96	15	5952	1092
2021-01-13	472	398	424	352	4392	48	46	815	89	24	6041	1116
2021-01-14	502	417	461	367	4486	41	50	806	94	16	6135	1132
2021-01-15	483	494	445	452	4479	38	42	802	110	11	6245	1143
2021-01-16	476	265	416	242	4653	60	23	839	83	14	6328	1157
2021-01-17	457	224	415	206	4862	42	18	863	93	9	6421	1166
2021-01-18	470	467	424	426	4860	46	41	868	112	18	6533	1184
2021-01-19	557	559	507	513	4854	50	46	872	122	19	6655	1203
2021-01-20	554	478	506	438	4922	48	40	880	116	15	6771	1218
2021-01-21	545	461	506	428	5000	39	33	886	124	12	6895	1230
2021-01-22	560	579	503	534	4969	57	45	898	164	19	7059	1249
2021-01-23	549	360	507	319	5157	42	41	899	146	19	7205	1268
2021-01-24	488	279	444	254	5347	44	25	918	136	19	7341	1287
2021-01-25	596	565	543	518	5372	53	47	924	142	17	7483	1304
2021-01-26	626	664	574	606	5340	52	58	918	159	26	7642	1330
2021-01-27	565	653	507	600	5247	58	53	923	160	16	7802	1346
2021-01-28	590	596	530	549	5228	60	47	936	141	12	7943	1358
2021-01-29	539	647	492	588	5132	47	59	924	148	17	8091	1375
2021-01-30	542	397	499	372	5259	43	25	942	162	17	8253	1392
2021-01-31	499	319	454	290	5423	45	29	958	168	16	8421	1408
2021-02-01	489	635	441	591	5273	48	44	962	149	17	8570	1425
2021-02-02	531	615	496	577	5192	35	38	959	125	17	8695	1442
2021-02-03	515	678	468	630	5030	47	48	958	126	20	8821	1462
2021-02-04	478	599	440	561	4909	38	38	958	154	13	8975	1475
2021-02-05	425	697	393	636	4666	32	61	929	127	20	9102	1495
2021-02-06	387	294	346	276	4736	41	18	952	122	10	9224	1505
2021-02-07	310	257	279	231	4784	31	26	957	115	13	9339	1518
2021-02-08	320	598	293	555	4522	27	43	941	116	15	9455	1533
2021-02-09	354	579	327	532	4317	27	47	921	99	15	9554	1548
2021-02-10	352	568	301	508	4110	51	60	912	120	22	9674	1570
2021-02-11	272	523	246	481	3875	26	42	896	104	14	9778	1584
2021-02-12	272	611	244	543	3576	28	68	856	93	24	9871	1608
2021-02-13	227	221	210	193	3593	17	28	845	78	19	9949	1627
2021-02-14	175	165	155	144	3604	20	21	844	62	12	10011	1639
2021-02-15	181	474	154	423	3335	27	51	820	73	16	10084	1655
2021-02-16	181	458	165	405	3095	16	53	783	92	11	10176	1666
2021-02-17	167	416	152	370	2877	15	46	752	66	19	10242	1685
2021-02-18	151	352	138	314	2701	13	38	727	51	7	10293	1692
2021-02-19	170	451	150	406	2445	20	45	702	51	5	10344	1697
2021-02-20	144	120	128	104	2469	16	16	702	44	10	10388	1707
2021-02-21	97	110	86	94	2461	11	16	697	47	8	10435	1715
2021-02-22	121	348	109	314	2256	12	34	675	40	4	10475	1719
2021-02-23	105	297	91	257	2090	14	40	649	36	9	10511	1728
2021-02-24	104	262	96	226	1960	8	36	621	28	8	10539	1736
2021-02-25	101	265	94	224	1830	7	41	587	33	14	10572	1750
2021-02-26	80	280	66	237	1659	14	43	558	24	11	10596	1761
2021-02-27	83	96	72	82	1649	11	14	555	21	9	10617	1770
2021-02-28	57	60	51	50	1650	6	10	551	25	5	10642	1775

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 27

Table 8: Table with daily admissions (Part 5)

Date	Entries	Discharges	non-ICU entries	non-ICU discharges	Non-ICU in hospital	ICU entries	ICU exits	ICU in hospital	Non-ICU deaths	ICU deaths	sum non-ICU deaths	sum ICU deaths
2021-03-01	86	244	80	203	1527	6	41	516	23	11	10665	1786
2021-03-02	70	200	58	172	1413	12	28	500	30	5	10695	1791
2021-03-03	71	181	63	156	1320	8	25	483	22	4	10717	1795
2021-03-04	67	176	57	153	1224	10	23	470	24	5	10741	1800
2021-03-05	68	209	64	180	1108	4	29	445	18	5	10759	1805
2021-03-06	60	53	55	45	1118	5	8	442	19	6	10778	1811
2021-03-07	38	41	35	32	1121	3	9	436	17	5	10795	1816
2021-03-08	45	184	40	155	1006	5	29	412	17	11	10812	1827
2021-03-09	52	149	43	123	926	9	26	395	13	6	10825	1833
2021-03-10	50	127	47	109	864	3	18	380	12	3	10837	1836
2021-03-11	41	112	35	91	808	6	21	365	10	4	10847	1840
2021-03-12	33	132	28	111	725	5	21	349	16	3	10863	1843
2021-03-13	33	39	30	32	723	3	7	345	7	7	10870	1850
2021-03-14	27	27	25	22	726	2	5	342	11	4	10881	1854
2021-03-15	29	95	28	71	683	1	24	319	6	6	10887	1860
2021-03-16	38	111	33	89	627	5	22	302	9	5	10896	1865
2021-03-17	41	82	35	65	597	6	17	291	6	7	10902	1872
2021-03-18	39	79	37	59	575	2	20	273	6	3	10908	1875
2021-03-19	35	100	31	73	533	4	27	250	8	5	10916	1880
2021-03-20	24	18	22	11	544	2	7	245	3	3	10919	1883
2021-03-21	28	24	24	18	550	4	6	243	6	5	10925	1888
2021-03-22	40	76	36	65	521	4	11	236	5	1	10930	1889
2021-03-23	37	79	32	57	496	5	22	219	6	7	10936	1896
2021-03-24	34	63	31	56	471	3	7	215	5	2	10941	1898
2021-03-25	34	60	29	51	449	5	9	211	5	0	10946	1898
2021-03-26	32	80	28	62	415	4	18	197	3	3	10949	1901
2021-03-27	25	25	22	20	417	3	5	195	4	4	10953	1905
2021-03-28	13	21	10	18	409	3	3	195	5	2	10958	1907
2021-03-29	22	63	20	56	373	2	7	190	2	1	10960	1908
2021-03-30	25	50	23	35	361	2	15	177	4	1	10964	1909
2021-03-31	34	61	32	48	345	2	13	166	8	3	10972	1912

A.3. Representation problem

As each entry is only introduced in the database when the patient receives hospital discharged, there is a need to cut some days from the end of the data timeline, to ensure a correct representation of the data. In order to do that we calculated the mean, median and the quantiles for the length of stay, before trimming the data, for either hospitalisation and ICU (table 9).

	mean	median	25%	75%	90%	95%	99%	max	IQR
non-ICU	12.57	8	4	15	27	38	69	348	11
ICU	24.34	18	11.0	30.0	50.7	67.0	114.9	372	19

Table 9: Summary statistics of the LoS before data trimming.

Therefore if we remove the last 63 days from the data, we ensure that at least 95% of patients in hospital at any day are present in the data. To facilitate the analysis, instead of removing the last 63 days, we only consider entries with date of entry until 31 of March 2021. Due to the low numbers of entries, dates before 1 of March 2020 were removed. The first case of COVID-19 in Portugal was confirmed by the Portuguese authorities at March 2nd 2020, but a couple of entries with date of entry prior to that date appear in the data, this could be explained by in hospital infections. As a consequence from the original 55827, 1142 entries were removed, being considered a total of 54685 entries.

A.4. Summary statistics

The first summary statistics (table 10) related to the length of stay, for either type of hospitalisation was used in figure 1 to plot the violin/box plot and to expands the information visible in the histograms as they are cut at 100 days of length of stay.

LoS	mean	median	25%	75%	90%	95%	99%	max	IQR
Non-ICU	12.54	8	4	15	27	38	67	348	11
ICU	24.31	18	11.0	30.3	51.0	66.0	112.0	216	19.3

Table 10: Summary statistics of the LoS by type of hospitalisation (non-ICU and ICU patients).

Table 11 shows the summary statistics of the length of stay for patients that died, corresponding to the time until death (TuD), and for patients that were discharged from hospital, corresponding to the Time until Discharge (TuDisc). These statistics are complementary to the figure 2 violin/box plot, where a difference in LoS between groups with different outcome from hospitalisation seems to arise.

		mean	median	25%	75%	90%	95%	99%	max	IQR
TuD	Non-ICU	11.01	7	3	14	24	33	61	228	11
	ICU	20.04	16	9.0	26.0	39.0	53.0	95.4	199.0	17
TuDisc	Non-ICU	12.98	9	5	16	28	39	69	348	11
	ICU	26.39	19	12	34	56	71	115	216	22

Table 11: Summary statistics of the length of stay for patients that died (TuD) or where discharged (TuDisc), for either type of hospitalisation (non-ICU and ICU patients).

Tables with the summary statistics for LoS (table 12) and TuD (table 13), in respect to the month (month of entry for LoS and of exit for TuD), age group, gender and region, with violin/box plots build from LoS and TuD in figure 14.

Table 12: Summary statistics of the LoS, by type of hospitalisation (non-ICU and ICU), entry month, age group, gender and region.

Variable	Group	Local	median	mean	sd	25%	75%	90%	95%	Max	IQR
Month	2020-03	non-ICU	12.00	19.70	23.28	6.00	25.00	46.00	63.00	348.00	19.00
		ICU	24.00	32.43	29.59	13.00	40.00	65.80	92.40	202.00	27.00
	2020-04	non-ICU	11.00	15.79	19.62	5.00	21.00	34.00	44.00	288.00	16.00
		ICU	22.00	27.11	20.55	13.00	37.00	54.70	72.55	109.00	24.00
	2020-05	non-ICU	10.00	15.49	20.68	4.00	20.00	33.80	44.00	277.00	16.00
		ICU	22.00	26.26	21.71	12.00	34.00	45.00	61.40	166.00	22.00
	2020-06	non-ICU	9.00	14.09	17.86	4.00	17.00	31.00	44.00	165.00	13.00
		ICU	21.00	28.66	24.91	12.00	38.75	55.50	78.75	141.00	26.75
	2020-07	non-ICU	10.00	15.61	22.11	4.50	17.00	33.00	51.10	236.00	12.50
		ICU	17.00	27.18	35.27	9.00	30.00	61.40	72.80	216.00	21.00
	2020-08	non-ICU	10.00	17.76	23.86	5.00	20.00	42.90	64.00	192.00	15.00
		ICU	18.00	29.60	32.76	10.00	37.50	68.00	86.50	193.00	27.50
	2020-09	non-ICU	11.00	17.12	20.79	5.00	21.00	39.00	55.90	230.00	16.00
		ICU	22.00	30.08	27.08	13.00	34.00	63.60	85.00	168.00	21.00
	2020-10	non-ICU	9.00	13.28	15.15	5.00	16.00	28.00	41.00	214.00	11.00
		ICU	17.00	24.80	23.73	11.00	30.75	53.50	69.00	163.00	19.75
	2020-11	non-ICU	8.00	12.24	13.30	4.00	15.00	26.00	37.00	200.00	11.00

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 29

		ICU	18.00	24.52	22.32	10.00	31.00	52.60	68.30	200.00	21.00
		non-ICU	9.00	13.14	13.25	5.00	17.00	29.00	40.00	145.00	12.00
2020-12		ICU	18.00	22.49	17.14	11.00	30.00	45.00	58.00	116.00	19.00
		non-ICU	8.00	10.99	10.83	4.00	14.00	24.00	31.00	158.00	10.00
2021-01		ICU	16.00	21.64	19.11	10.00	26.00	44.60	61.00	158.00	16.00
		non-ICU	7.00	9.59	8.85	4.00	12.00	20.00	26.00	97.00	8.00
2021-02		ICU	17.00	22.02	17.97	10.00	28.00	44.00	60.15	119.00	18.00
		non-ICU	7.00	10.07	9.54	4.00	13.00	21.00	28.40	78.00	9.00
2021-03		ICU	17.00	19.31	13.76	10.00	25.00	38.30	48.30	75.00	15.00
Age Groups	[0,5)	non-ICU	3.00	6.28	11.10	1.00	6.00	13.00	20.00	101.00	5.00
	[5,10)	ICU	15.00	33.15	45.65	6.50	36.50	87.60	111.60	200.00	30.00
		non-ICU	4.00	5.91	6.27	1.00	8.00	12.40	17.20	34.00	7.00
	[10,15)	ICU	16.00	16.40	15.08	6.00	19.00	31.60	35.80	40.00	13.00
		non-ICU	3.00	7.13	9.79	1.25	9.00	18.40	30.00	47.00	7.75
	[15,20)	ICU	10.00	10.87	6.79	6.50	14.50	20.20	21.90	24.00	8.00
		non-ICU	3.00	7.96	19.75	2.00	7.00	17.00	29.50	277.00	5.00
	[20,25)	ICU	11.50	20.42	21.08	9.75	17.00	47.30	61.70	76.00	7.25
		non-ICU	4.00	7.08	13.05	2.00	7.00	15.00	26.00	225.00	5.00
	[25,30)	ICU	13.00	18.64	18.09	9.00	26.00	31.00	39.80	89.00	17.00
		non-ICU	3.00	7.49	13.90	2.00	7.00	18.00	28.00	288.00	5.00
	[30,35)	ICU	13.00	20.95	21.04	8.50	27.00	41.80	56.60	101.00	18.50
		non-ICU	4.00	7.12	9.38	2.00	8.00	15.00	26.00	66.00	6.00
	[35,40)	ICU	10.50	14.14	11.50	7.00	15.00	30.50	41.25	58.00	8.00
		non-ICU	5.00	8.88	12.84	3.00	10.00	19.00	29.00	165.00	7.00
	[40,45)	ICU	13.00	17.76	17.42	9.00	21.75	36.00	39.95	129.00	12.75
		non-ICU	6.00	9.17	11.02	4.00	10.00	19.00	28.00	143.00	6.00
	[45,50)	ICU	13.00	20.10	22.70	8.00	23.00	40.00	50.00	216.00	15.00
		non-ICU	7.00	9.92	11.29	4.00	11.00	21.00	30.00	213.00	7.00
	[50,55)	ICU	14.00	21.54	21.60	9.00	26.75	46.00	61.00	193.00	17.75
		non-ICU	8.00	10.86	11.88	5.00	12.00	23.00	32.00	132.00	7.00
	[55,60)	ICU	18.00	23.31	22.43	11.00	29.00	43.00	54.50	202.00	18.00
		non-ICU	8.00	11.11	12.50	5.00	13.00	23.00	32.00	167.00	8.00
	[60,65)	ICU	19.00	24.70	20.31	11.00	30.25	53.00	64.05	151.00	19.25
		non-ICU	8.00	12.26	13.80	5.00	14.00	26.00	37.00	239.00	9.00
	[65,70)	ICU	19.00	25.09	21.32	11.00	33.00	52.00	66.00	180.00	22.00
		non-ICU	9.00	13.05	15.30	5.00	15.00	27.00	38.00	228.00	10.00
	[70,75)	ICU	19.00	25.50	22.75	12.00	31.00	53.90	67.00	199.00	19.00
		non-ICU	10.00	14.06	15.39	5.00	17.00	30.00	42.00	348.00	12.00
	[75,80)	ICU	20.00	26.46	21.94	12.00	34.00	56.00	72.00	168.00	22.00
		non-ICU	10.00	14.32	15.19	5.00	18.00	31.00	41.00	230.00	13.00
	[80,150)	ICU	19.00	25.79	23.40	11.00	31.00	58.50	80.25	168.00	20.00
		non-ICU	9.00	13.44	14.54	5.00	17.00	29.00	39.00	267.00	12.00
		ICU	17.00	22.67	20.96	9.00	28.00	46.70	61.00	210.00	19.00
Gender	Female	non-ICU	8.00	12.32	14.51	4.00	15.00	27.00	37.00	348.00	11.00
		ICU	17.00	23.82	22.36	10.00	29.00	50.00	67.00	210.00	19.00
	Male	non-ICU	9.00	12.74	14.09	5.00	16.00	27.00	38.00	277.00	11.00
Region	Alentejo	ICU	18.00	24.53	21.82	11.00	31.00	51.00	66.00	216.00	20.00
		non-ICU	7.00	10.35	11.55	4.00	13.00	21.00	29.00	178.00	9.00
	Algarve	ICU	19.00	23.87	19.52	12.00	29.25	48.00	58.25	126.00	17.25
		non-ICU	8.00	13.12	16.93	4.00	14.00	28.20	44.00	163.00	10.00
	Centro	ICU	17.00	22.83	20.86	11.50	27.00	42.80	57.80	216.00	15.50
		non-ICU	9.00	12.88	12.15	5.00	16.00	26.00	36.00	135.00	11.00
	LVT	ICU	19.00	23.50	16.28	12.00	31.00	44.00	55.10	113.00	19.00
		non-ICU	9.00	13.28	15.03	4.00	17.00	29.00	40.00	348.00	13.00
	Norte	ICU	18.00	24.56	21.92	11.00	31.00	51.50	66.00	210.00	20.00
		non-ICU	8.00	11.77	14.31	4.00	14.00	26.00	36.00	277.00	10.00
		ICU	17.00	24.44	23.56	10.00	31.00	54.00	70.65	202.00	21.00

1

Table 13: Summary statistics of the TuD, by type of hospitalisation (non-ICU and ICU), month of discharge, age group, gender and region.

Variable	Group	Local	median	mean	sd	25%	75%	90%	95%	Max	IQR
Month	2020-03	non-ICU	3.00	4.59	4.22	2.00	6.00	9.00	11.40	24.00	4.00
		ICU	6.50	7.89	6.61	2.00	11.25	16.60	19.30	26.00	9.25
	2020-04	non-ICU	7.00	9.43	8.32	3.00	13.00	21.00	25.30	49.00	10.00
		ICU	12.00	14.24	9.83	7.00	22.00	29.80	32.90	38.00	15.00
	2020-05	non-ICU	12.00	18.06	16.78	6.00	25.25	46.00	52.00	75.00	19.25
		ICU	24.50	28.77	18.05	18.75	35.25	49.70	62.40	80.00	16.50
	2020-06	non-ICU	8.00	15.22	18.08	5.00	18.00	38.20	50.10	98.00	13.00
		ICU	26.00	32.21	29.02	10.00	43.00	75.60	91.00	101.00	33.00
	2020-07	non-ICU	8.00	15.46	19.48	4.50	18.50	33.20	44.10	135.00	14.00
		ICU	12.00	24.36	32.26	6.00	24.75	61.40	110.20	139.00	18.75
	2020-08	non-ICU	11.50	17.62	17.88	5.00	22.50	39.50	61.00	90.00	17.50
		ICU	15.50	28.06	35.07	9.25	29.00	56.50	73.65	151.00	19.75
	2020-09	non-ICU	8.00	11.34	12.33	4.00	14.00	21.30	37.10	85.00	10.00
		ICU	12.00	13.79	9.15	9.00	19.00	22.20	29.40	43.00	10.00
	2020-10	non-ICU	7.00	10.45	11.91	3.00	13.00	22.00	31.60	90.00	10.00
		ICU	13.00	17.87	18.22	6.50	24.00	37.60	46.00	141.00	17.50
	2020-11	non-ICU	6.00	9.36	11.60	3.00	12.00	19.00	26.55	163.00	9.00
		ICU	13.00	15.89	13.61	7.00	21.00	31.00	39.35	92.00	14.00
	2020-12	non-ICU	8.00	11.03	12.37	4.00	14.00	22.90	30.45	206.00	10.00
		ICU	17.00	19.01	12.10	10.00	26.25	35.00	40.25	82.00	16.25

	2021-01	non-ICU	6.50	10.20	11.91	3.00	13.00	23.00	32.00	149.00	10.00
		ICU	13.00	17.46	15.79	8.00	21.75	37.00	51.75	98.00	13.75
	2021-02	non-ICU	8.00	10.87	10.43	4.00	14.00	24.00	31.00	106.00	10.00
		ICU	17.00	20.44	17.46	12.00	25.00	32.40	45.70	180.00	13.00
	2021-03	non-ICU	14.00	19.73	20.59	7.00	25.00	41.00	58.20	178.00	18.00
		ICU	26.00	27.50	17.49	17.00	36.00	49.00	56.00	134.00	19.00
	2021-04	non-ICU	31.00	37.34	30.14	17.00	43.50	80.40	84.80	142.00	26.50
		ICU	35.00	39.84	24.17	20.00	64.00	75.60	84.20	89.00	44.00
	2021-05	non-ICU	96.50	114.75	86.61	52.25	159.00	200.40	214.20	228.00	106.75
		ICU	73.00	84.70	49.85	46.25	105.75	128.80	163.90	199.00	59.50
Age Groups	[0,5)	non-ICU	2.00	2.00	-	2.00	2.00	2.00	2.00	2.00	0.00
		ICU	18.00	18.00	-	18.00	18.00	18.00	18.00	18.00	0.00
	[5,10)	non-ICU	-	-	-	-	-	-	-	-	-
		ICU	-	-	-	-	-	-	-	-	-
	[10,15)	non-ICU	-	-	-	-	-	-	-	-	-
		ICU	1.00	1.00	-	1.00	1.00	1.00	1.00	1.00	0.00
	[15,20)	non-ICU	48.00	48.00	-	48.00	48.00	48.00	48.00	48.00	0.00
		ICU	-	-	-	-	-	-	-	-	-
	[20,25)	non-ICU	23.00	23.00	-	23.00	23.00	23.00	23.00	23.00	0.00
		ICU	-	-	-	-	-	-	-	-	-
	[25,30)	non-ICU	6.00	14.20	23.48	2.00	6.00	36.00	46.00	56.00	4.00
		ICU	30.00	30.00	-	30.00	30.00	30.00	30.00	30.00	0.00
	[30,35)	non-ICU	6.50	10.75	11.66	3.50	13.50	21.60	29.30	37.00	10.00
		ICU	7.00	8.29	4.27	6.00	11.50	13.40	13.70	14.00	5.50
	[35,40)	non-ICU	6.00	15.59	23.32	5.00	17.00	29.80	49.20	98.00	12.00
		ICU	17.00	24.83	32.34	7.50	23.50	40.30	77.55	121.00	16.00
	[40,45)	non-ICU	8.00	15.50	14.80	4.75	26.75	39.00	40.95	51.00	22.00
		ICU	11.50	20.75	27.95	7.25	23.50	37.40	46.10	139.00	16.25
	[45,50)	non-ICU	9.50	14.62	15.81	3.75	21.00	29.30	45.45	83.00	17.25
		ICU	16.00	26.24	29.27	8.00	30.00	61.80	78.60	142.00	22.00
	[50,55)	non-ICU	10.00	16.17	16.96	5.00	22.50	37.70	51.45	83.00	17.50
		ICU	20.00	24.91	22.40	12.00	30.75	46.00	60.70	141.00	18.75
Gender	Female	non-ICU	7.00	10.22	12.30	3.00	13.00	22.00	31.00	206.00	10.00
		ICU	15.00	18.99	18.87	8.00	24.00	37.00	53.00	180.00	16.00
	Male	non-ICU	8.00	11.74	13.19	4.00	15.00	25.00	34.00	228.00	11.00
		ICU	17.00	20.47	18.27	9.00	26.00	39.00	53.00	199.00	17.00
Region	Alentejo	non-ICU	5.00	7.83	10.97	3.00	10.00	16.00	22.25	178.00	7.00
		ICU	17.00	20.84	18.29	7.00	23.50	51.50	58.75	89.00	16.50
	Algarve	non-ICU	8.00	13.94	20.79	4.00	15.00	28.80	41.80	163.00	11.00
		ICU	16.00	18.92	11.30	11.25	26.00	30.00	38.75	64.00	14.75
	Centro	non-ICU	7.00	10.51	10.97	4.00	13.00	23.00	29.00	135.00	9.00
		ICU	16.00	18.50	13.15	10.00	24.25	34.00	40.45	96.00	14.25
	LVT	non-ICU	8.00	12.08	13.94	4.00	16.00	26.00	36.00	228.00	12.00
		ICU	15.00	19.56	17.76	8.00	26.00	38.60	54.60	180.00	18.00
	Norte	non-ICU	7.00	10.29	11.59	3.00	13.00	22.00	32.00	163.00	10.00
		ICU	16.00	21.19	21.40	8.00	27.00	43.00	59.50	199.00	19.00

A.5. Additional plots

Figure 14: Violin/box plots of the (a)LoS and (b)TuD, by month and type of hospitalisation (white diamonds represent the mean).

(a)

(b)

(c)

Figure 15: Evolution of patients (a)in-hospital, (b)discharged and (c)death with time, by type of hospitalization.

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 33

(a)

(b)

Figure 16: Comparing entries and exits by time, for (a)non-ICU and (b)ICU patients.

(a)

(b)

(c)

Figure 17: Characterization of the age groups. (a) Violin/box plot of the LoS by age groups (white diamonds represent the mean). Histograms of the number of entries, by age group, for (b) non-ICU and (c) ICU patients

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 35

(a)

(b)

(c)

Figure 18: (a) Violin/box plot of the TuD by age groups (white diamonds represent the mean). Histogram of the fatalities, by age group, of (b) non-ICU and (c) ICU patients

Figure 19: Violin/box plot of the TuD by gender and type of hospitalisation (white diamonds represent the mean).

(a)

(b)

Figure 20: Violin/box plot of the LoS by gender and type of hospitalisation (white diamonds represent the mean).

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 37

Figure 21: Violin/box plot of the TuD by gender and type of hospitalisation (white diamonds represent the mean).

Figure 22: (a,b,c,d)Histograms with entries and fatalities for ICU and non-ICU patients. Linegraphs of fatalities rates by group for (e)non-ICU and (f)ICU patients. Percentage of males in (g)non-ICU and (h)ICU patients, for age group. Fatality rate over time for (i)non-ICU and (j)ICU.

COVID-19 Hospitalisation in Portugal, the first year: Results from hospital discharge data 39

Figure 23: Violin/box plots of (a)LoS and (b)TuD by region (white diamonds represent the mean).

Figure 24: Linegraphs of fatalities rates by age group and region for (a) non-ICU(a) and (b)ICU patients. Fatality rate by region and over time for (c)non-ICU and (d)ICU (white diamonds represent the mean).

Figure 25: Linegraphs of entries by region for (a)non-ICU and (b)ICU patients entries, and fatalities for (c)non-ICU and (d)ICU patients fatalities

A.6. Distribution fitting

Local	Hosp	ICU
min	1	1
max	348	216
median	9	18
mean	12.81596	24.43704
estimated sd	14.33082	21.97512
estimated skewness	4.475019	2.608978
estimated kurtosis	45.76527	13.74304

Table 14: LoS summary statistics for the Cullen and Frey graph (Figure 11).

LoS		non-ICU	ICU
gamma	shape	1.3009479	1.236824
	rate	0.1015242	0.05061267
log-normal	mean log	2.119463	2.8582927
	sd log	0.9495999	0.8655452
Weibull	shape	1.263677	1.253865
	scale	13.09343	26.429118
gamma ADL	shape	1.5623940	1.236824
	rate	0.1403541	0.05061267
log-normal ADL	mean log	2.1465055	2.8986155
	sd log	0.9885536	0.8565485

Table 15: Estimated LoS distribution parameters.

	gamma	lognormal	Weibull
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.07854276	0.05344594	0.07335166
Cramer-von Mises statistic	63.00402211	15.96330257	66.41153408
Anderson-Darling statistic	365.04226391	127.87010687	446.79678352
Goodness-of-fit criteria			
Akaike's Information Criterion	336511.5	332306.2	337756.5
Bayesian Information Criterion	336529.0	332323.8	337774.0
	gamma ADL	lognormal ADL	
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.04724813	0.04920051	
Cramer-von Mises statistic	24.02657920	17.63190355	
Anderson-Darling statistic	Inf	145.71971535	
Goodness-of-fit criteria			
Akaike's Information Criterion	339100.5	332491.9	
Bayesian Information Criterion	339118.1	332509.4	

Table 16: LoS goodness-of-fit statistics for non-ICU patients.

	gamma	lognormal	Weibull
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.06457332	0.0566513	0.06895957
Cramer-von Mises statistic	2.8023735	1.76070045	8.34009006
Anderson-Darling statistic	19.9448961	12.46493868	50.56938674
Goodness-of-fit criteria			
Akaike's Information Criterion	49117.16	49102.09	49350.59
Bayesian Information Criterion	49130.54	49115.47	49363.97
	gamma ADL	lognormal ADL	
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.04424102	0.03867471	
Cramer-von Mises statistic	1.97564933	1.69481544	
Anderson-Darling statistic	32.65844445	16.87600111	
Goodness-of-fit criteria			
Akaike's Information Criterion	49360.57	49116.56	
Bayesian Information Criterion	49373.95	49129.94	

Table 17: LoS goodness-of-fit statistics for ICU patients.

Figure 26: TuD Cullen and Frey graphs for (a) Non-ICU and (b) ICU patients.

Local	Hosp	ICU
min	1	1
max	228	199
median	8	16
mean	11.33318	20.27451
estimated sd	11.33318	18.43436
estimated skewness	11.33318	3.072088
estimated kurtosis	37.3627	18.92779

Table 18: TuD Summary statistics for the Cullen and Frey graph (Figure 26).

TuD		non-ICU	ICU
gamma	shape	1.2314081	1.57990102
	rate	0.1087638	0.07792103
log-normal	meanlog	1.9684210	2.660706
	sdlog	0.9873343	0.898466
Weibull	shape	1.244128	1.246129
	scale	21.859276	21.898314
gamma ADL	shape	1.73385831	1.73596577
	rate	0.08887636	0.08882768
log-normal ADL	meanlog	2.734321	2.736472
	sdlog	0.939225	0.938759

Table 19: Estimated TuD distribution parameters.

	gamma	lognormal	Weibull
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.06756849	0.05607134	0.07106338
Cramer-von Mises statistic	10.79585720	5.31072150	10.18207602
Anderson-Darling statistic	70.37506509	46.28643673	78.49461693
Goodness-of-fit criteria			
Akaike's Information Criterion	73132.70	72288.43	73326.16
Bayesian Information Criterion	73147.26	72302.99	73340.71
	gamma ADL	lognormal ADL	
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.04833295	0.0518889	
Cramer-von Mises statistic	4.80060941	5.4628413	
Anderson-Darling statistic	66.41089597	52.02366989	
Goodness-of-fit criteria			
Akaike's Information Criterion	73448.81	72412.75	
Bayesian Information Criterion	73463.37	72427.31	

Table 20: TuD goodness-of-fit statistics for non-ICU patients.

	gamma	lognormal	Weibull
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.04668797	0.08914086	0.05729949
Cramer-von Mises statistic	0.92583551	2.88870410	1.77441464
Anderson-Darling statistic	5.93558166	17.15111131	11.48495042
Goodness-of-fit criteria			
Akaike's Information Criterion	15329.37	15401.71	15391.03
Bayesian Information Criterion	15340.51	15412.85	15402.17
	gamma ADL	lognormal ADL	
Goodness-of-fit statistics			
Kolmogorov-Smirnov statistic	0.03759531	0.07105014	
Cramer-von Mises statistic	0.39364935	2.46654705	
Anderson-Darling statistic	3.98784875	18.87662199	
Goodness-of-fit criteria			
Akaike's Information Criterion	15344.69	15421.03	
Bayesian Information Criterion	15355.83	15432.17	

Table 21: TuD goodness-of-fit statistics for ICU patients.

A.7. Links

- 1 In the main text of the paper links for websites are provided, them being:
- 2 • **link1:** [https://www.acss.min-saude.pt/wp-content/uploads/2016/10/](https://www.acss.min-saude.pt/wp-content/uploads/2016/10/SNS-RH-ARS-n1-dados-a-junho-de-2016.pdf)
 - 3 [SNS-RH-ARS-n1-dados-a-junho-de-2016.pdf](https://www.acss.min-saude.pt/wp-content/uploads/2016/10/SNS-RH-ARS-n1-dados-a-junho-de-2016.pdf);
 - 4 • **link2:** [https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&](https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&ind0corrCod=0011166&xlang=pt)
 - 5 [ind0corrCod=0011166&xlang=pt](https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&ind0corrCod=0011166&xlang=pt);
 - 6 • **link3:** [https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&](https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&contecto=pi&ind0corrCod=0011166&selTab=tab0)
 - 7 [contecto=pi&ind0corrCod=0011166&selTab=tab0](https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&contecto=pi&ind0corrCod=0011166&selTab=tab0).