


**Table S1: Demographic and clinical description of the cohort, Related to STAR Methods**

Study number	Age	Gender	Days sampled post infection	Vaccination status	Days since vaccination	Sequence	Severity
COV132	36-40	Male	9	Unvaccinated	-	Not sequenced	Severe
COV140	26-30	Female	1	Unvaccinated	-	Not sequenced	Mild
COV145	16-20	Female	3	Unvaccinated	-	Omicron	Mild
COV149	60-65	Female	3	Unvaccinated	-	Not sequenced	Mild
COV157	66-70	Male	4	Unvaccinated	-	Omicron	Moderate
COV158	66-70	Female	6	Unvaccinated	-	Not sequenced	Moderate
COV159	20-25	Male	6	Unvaccinated	-	Not sequenced	Mild
COV162	56-60	Male	5	Unvaccinated	-	Not sequenced	Moderate
COV165	60-65	Female	4	Unvaccinated	-	Not sequenced	Moderate
COV166	56-60	Male	4	Unvaccinated	-	Not sequenced	Severe
COV167	60-65	Female	8	Unvaccinated	-	Not sequenced	Severe
COV176	70-75	Male	2	Unvaccinated	-	Not sequenced	Moderate
COV178	60-65	Female	2	Unvaccinated	-	Omicron	Moderate
COV183	26-30	Male	2	Unvaccinated	-	Not sequenced	Mild
COV184	56-60	Male	1	Unvaccinated	-	Not sequenced	Moderate
COV185	46-50	Female	1	Unvaccinated	-	Not sequenced	Moderate
COV186	56-60	Male	10	Unvaccinated	-	Omicron	Moderate
COV187	30-35	Female	3	Unvaccinated	-	Not sequenced	Mild
COV189	46-50	Male	5	Unvaccinated	-	Not sequenced	Moderate
COV190	30-35	Female	4	Unvaccinated	-	Omicron	Mild
COV151	46-50	Male	1	One dose Ad26.CoV2.S	56	Omicron	Severe
COV182	36-40	Male	3	One dose Ad26.CoV2.S	Unknown	Not sequenced	Mild
COV131	56-60	Female	4	Two doses BNT162b2	109	Not sequenced	Severe
COV135	80-85	Male	5	Two doses BNT162b2	72	Omicron	Mild
COV138	66-70	Male	5	Two doses BNT162b2	Unknown	Not sequenced	Moderate
COV161	60-65	Female	5	Two doses BNT162b2	112	Not sequenced	Mild
COV181	70-75	Male	4	Two doses BNT162b2	163	Not sequenced	Severe



**Figure S1: Omicron BA.1 neutralization titers are substantially boosted in response to Omicron infection compared to infection with other VOCs or vaccination, Related to Figure 2**

(A) Neutralization  $ID_{50}$  titers from vaccinees two months post two-doses of BNT162b2 or one-dose of Ad26.COVS, individuals acutely infected with D614G, Beta, Delta or Omicron or individuals who were vaccinated with either BNT162b2 or Ad26.COVS and then infected with Omicron (BTI = breakthrough infection) against D614G (white) and Omicron BA.1 (orange) SARS-CoV-2 variants. Lines indicate geometric mean titer (GMT). (B) Fold difference of neutralization titer against Omicron BA.1 relative to D614G for each individual is shown with median fold difference indicated by the bold line. The dotted line indicates no fold difference.