

## Supplementary material

**Supplementary table:** Variables included in regression model

Variable	Definition	Responses	Component of model
Patient ID	Unique identifier	ID, URN, MRN	Model random factor
Age	Age at time of surgery	years	Model covariate
Gender	As identified by the patient	Male; Female; Not specified	Model covariate
BMI	Body mass index as per standard formula	kg/(m <sup>2</sup> )	Model covariate
Secondary diagnosis	Injuries to other knee structures	meniscus (lateral/medial); cartilage (femoral/tibial/patella); ligament (collateral/PCL); other	Model covariate
DateofInjury to DateofSurgery	Time delay between when the injury occurred and when surgery is performed	Weeks	Model covariate
Timepoint	Post-surgery delay of measurement of maximum knee flexion	Initial Maximum	Model covariate
SGTS	Was a SGTS system used?	Yes; No	Model primary predictor
Maximum extension	Minimum flexion angle under passive loading forcing the knee into extension. Relative to anatomical zero	Degrees	Model primary outcome

## Supplementary data: Regression model code (STATA)

```
1 import delimited "Filepath MasterSheet.csv", stringcols(1)
2
3 *Swap Filepath for local filepath
4
5 // Mixed effects logistic regression as per DAG figure (adjusted variables as covariates) (updated
6 // Sept 2021)
7
8 * encode categorical variables
9 label define sexcode 0 "Male" 1 "Female", replace
10 encode sex, gen(sexcode)
11 encode secondary, gen(secondcode)
12 label define sgtscode 0 "No" 1 "Yes", replace
13 encode sgts, gen(sgtscode)
14 label define timecode 0 "Initial" 1 "Maximum", replace
15 encode timepoint, gen(timecode)
16 label define loecode 0 "No" 1 "Yes", replace
17 encode loestatus, gen(loecode)
18 label define notchcode 0 "No" 1 "Yes", replace
19 encode notchplasty, gen(notchcode)
20 encode secmed, gen(secmedcode)
21 encode seclat, gen (seclatcode)
22 encode secother, gen(othercode)
23 encode presromrest, gen(presromcode)
24
25 *Check missing pattern of key variables
26 misstable summarize, all
27 misstable patterns ageatsurgery sexcode secondcode dos_physio bmi sgtscode timecode
28
29 *Sort data to summarize
30 sort timecode
31
32 *Create table summary by timepoint
33 *by timecode: summarize, separator(4)
34 table sgtscode notchcode, statistic (count notchcode)
35
36 table sgtscode, statistic(count ageatsurgery bmi dos_physio doi_dos initial_max)
37 table sgtscode, statistic(median ageatsurgery bmi dos_physio doi_dos initial_max)
38 table sgtscode, statistic(q1 ageatsurgery bmi dos_physio doi_dos initial_max)
39 table sgtscode, statistic(q3 ageatsurgery bmi dos_physio doi_dos initial_max)
40
41 *Set and perform melogit
42
43 *label define loecode 0 "No" 1 "Yes", replace
44 melogit loecode i.sgtscode#i.timecode c.ageatsurgery c.secondcode c.bmi i.sexcode c.doi_dos, ||
45 patientid:
46 // Calculated proportions for sgtscode
47 margins sgtscode
48 // Calculated proportions for sgtscode
49 margins sgtscode#timecode
50 // Calculate differences in proportions between TLS levels (Yes vs No)
51 margins ar.sgtscode
52 // Calculate proportions for sgtscode##timecode
53 margins ar.sgtscode@timecode
54
55 save "Filepath Analysis - First Pass.dta", replace
56 *Swap Filepath for local filepath
```