

16 **Supplemental Table 3A. Associations with Exacerbation Within 1 Year of Enrollment by Logistic Regression.***

Clinical Univariable	Estimate	Standard Error (SE)	<i>p</i>	95% CI
Age (years)	-0.028	0.016	0.09	-0.059 to 0.0043
Sex (Male=0, Female=1)	0.23	0.39	0.55	-0.53 to 1.00
FEV ₁ %	-0.022	0.0095	0.02	-0.041 to -0.0034
Number of Prior Pulmonary Exacerbations	0.64	0.17	<0.001	0.30 to 0.98
Diabetes Status	1.36	0.59	0.02	0.22 to 2.50
<i>Burkholderia cepacia</i> Complex Infection Status	16	1400	0.99	-2700 to 2700
Weight-for-age z-score	-0.24	0.20	0.23	-0.64 to 0.16
Methicillin Sensitive <i>Staphylococcus aureus</i> Infection Status	-0.18	0.39	0.64	-0.95 to 0.58
Pancreatic Sufficiency	-0.83	0.70	0.24	-2.20 to 0.54
Methicillin Resistant <i>S aureus</i> Infection Status	0.75	0.55	0.18	-0.34 to 1.80
<i>Pseudomonas aeruginosa</i> Infection Status	0.76	0.40	0.06	-0.026 to 1.50

17 *Univariable Logistic Regression. There was no missingness to consider among these clinical variables.

18

19

20 **Supplemental Table 3B. Associations with Exacerbation Within 1 Year of Enrollment by Proportional Hazards Modeling.***

Clinical Univariable	Estimate	SE	Hazard Ratio	<i>p</i>	95% CI of the Hazard Ratio
Age (years)	-0.023	0.012	0.98	0.044	0.96 to 1.00
Sex (Male=0, Female=1)	0.16	0.24	1.20	0.49	0.74 to 1.90
FEV ₁ %	-0.015	0.006	0.99	0.009	0.97 to 1.00
Number of Prior Pulmonary Exacerbations	0.32	0.071	1.40	<0.001	1.20 to 1.60
Diabetes Status	0.61	0.26	1.80	0.02	1.10 to 3.10
<i>Burkholderia cepacia</i> Complex Infection Status	0.50	0.59	1.70	0.39	0.52 to 5.30
Weight-for-age z-score	-0.30	0.14	0.74	0.034	0.57 to 0.98
Methicillin Sensitive <i>Staphylococcus aureus</i> Infection Status	-0.14	0.24	0.87	0.56	0.55 to 1.40
Pancreatic Sufficiency	-0.56	0.51	0.57	0.27	0.21 to 1.60
Methicillin Resistant <i>S aureus</i> Infection Status	0.39	0.28	1.50	0.17	0.85 to 2.60
<i>Pseudomonas aeruginosa</i> Infection Status	0.43	0.25	1.50	0.092	0.93 to 2.50

21 *Univariable Proportional Hazards Models. There was no missingness to consider among these clinical variables.

22

Supplemental Table 3C. Associations with Exacerbation Within Entire Study by Proportional Hazards Modeling.*

Clinical Univariable	Estimate	SE	Hazard Ratio	<i>p</i>	95% CI of the Hazard Ratio
Age (years)	-0.024	0.01	0.98	0.022	0.96 to 1.00
Sex (Male=0, Female=1)	0.22	0.21	1.20	0.29	0.82 to 1.90
FEV ₁ %	-0.012	0.005	0.99	0.013	0.98 to 1.00
Number of Prior Pulmonary Exacerbations	0.28	0.064	1.30	<0.001	1.20 to 1.50
Diabetes Status	0.66	0.24	1.90	0.0062	1.20 to 3.10
<i>Burkholderia cepacia</i> Complex Infection Status	0.50	0.59	1.70	0.39	0.52 to 5.30
Weight-for-age z-score	-0.31	0.13	0.74	0.016	0.57 to 0.94
Methicillin Sensitive <i>Staphylococcus aureus</i> Infection Status	-0.04	0.21	0.96	0.83	0.63 to 1.40
Pancreatic Sufficiency	-0.67	0.46	0.51	0.15	0.21 to 1.30
Methicillin Resistant <i>S aureus</i> Infection Status	0.31	0.26	1.40	0.24	0.81 to 2.30
<i>Pseudomonas aeruginosa</i> Infection Status	0.22	0.22	1.20	0.32	0.81 to 1.9

*Univariable Proportional Hazards Models. There was no missingness to consider among these clinical variables.