

Environmental impacts associated with UPF consumption: which food chain stages matter the most? Findings from a representative sample of French adults

Emmanuelle Kesse-Guyot¹, Benjamin Allès¹, Joséphine Brunin^{1,2}, Hélène Fouillet³, Alison Dussiot⁴, Florine Berthy¹, Elie Perraud³, Serge Hercberg^{1,4}, Chantal Julia^{1,4}, François Mariotti³, Mélanie Deschasaux-Tanguy¹, Bernard Srour¹, Denis Lairon⁵, Philippe Pointereau⁶, Julia Baudry^{1,7}, Mathilde Touvier^{1,7}

¹ Sorbonne Paris Nord University, Inserm, INRAE, Cnam, Nutritional Epidemiology Research Team (EREN), Epidemiology and Statistics Research Center – University Paris Cité (CRESS), 93017 Bobigny, France

² ADEME, (Agence de l'Environnement et de la Maîtrise de l'Energie), 49004 Angers, France

³ Paris-Saclay University, UMR PNCA, AgroParisTech, INRAE, 75005 Paris, France

⁴ Département de Santé Publique, Hôpital Avicenne, 93017 Bobigny, France

⁵ Aix Marseille Université, Inserm, INRAE, C2VN, 13005 Marseille, France

⁶ Solagro, 75, Voie TOEC, CS 27608, F-31076 Toulouse Cedex 3, France

⁷ Authors equally contributed

Correspondence: Email: emmanuelle.kesse-guyot@inrae.fr

Equipe de Recherche en Epidémiologie Nutritionnelle (EREN)

SMBH Université Sorbonne Paris Nord, 74 rue Marcel Cachin, 93017 Bobigny, France

Running title: Environmental impacts of ultra-processed food consumption

Number of tables: 4 / **Number of figures:** 2

Supplemental material: 3 methods, 4 tables and 2 figures

Data availability: Data described in the manuscript are available at : <https://www.data.gouv.fr/fr/datasets/donnees-de-consommations-et-habitudes-alimentaires-de-letude-inca-3/>.

Abstract

1 **Background** Scientific literature about environmental pressures associated with dietary patterns has
2 been considerably growing over the last decade. However, few studies have analyzed the
3 environmental impacts associated with the consumption of ultra-processed food (UPF) and which
4 steps of the food system that contribute most to environmental pressures. The objective of this study
5 was to investigate, in a representative sample of the adult French population, the environmental
6 pressures of diets according to UPF consumption.
7 **Methods** The study was conducted in 2,121 adults of the French nationally representative survey
8 INCA 3. Food intakes were analyzed to define the %UPF (in weight) in the diet according to NOVA
9 classification. Using detailed environmental data of foods of Agribalyse, we could assess the
10 contribution of UPF to 14 indicators of environmental pressure and details those related to the stage of
NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

11 the food consumed: production, processing, storage, packaging, transport and retailing at the food
12 level. The data was described according to quintiles of % of UPF in the diet and analyzed using crude
13 and energy-adjusted models.

14 **Results** Compared to low consumers of UPF (Q1, median UPF= 7%), high consumers (Q5, median
15 UPF= 35%) had a higher energy intake (+22%) which came along with different environmental
16 pressures (e.g. +15% greenhouse gas emissions (GHGe), +17% land use, -7% water use and +8%
17 cumulative energy demand). Higher pressures could be ascribed to higher energy. However, the
18 processing and packaging stages were significant contributor to energy demand.

19 In Q5, contributions of the UPF category to total pressure were 35%, 39%, 28% and 42% for GHGe,
20 water use land use, and energy demand, respectively, while ranging from 11 to 15% in Q1.

21 **Conclusions** Diets rich in UPF, compared to low, were overall associated with intensification in
22 GHGe, land use, and energy demand and with higher contribution of post-farm stages, in particular
23 processing regarding energy demand.

24 **Keywords:** Environmental footprint, Ultra-processed food, diet, food chain

Highlights

- Overall, higher UPF consumption was associated with higher environmental impact, in particular GHGE and land use
- Low UPF consumers had overall higher water footprint (due to their high fruit and vegetable intake)
- A large part of the higher pressures observed among participants with higher UPF consumption was explained by their higher dietary energy intake since the associations did not remain or were even reversed after energy adjustment
- Regardless of the % UPF in the diet, GHGe, land use and water use pressures mostly occurred at the stage of agricultural production, in contrast, packaging and processing stages were also important contributors to energy demand
- Contribution of the UPF category to total impact/pressure varied across indicators, with a high contribution of the UPF category to energy demand, due to the packaging and processing stages, but a low contribution to GHGe and land use, because higher consumers of UPF are lower consumers of animal products

Introduction

25 In the recent decades, food systems and food supplies have become largely globalized with changes
26 occurring at all steps of the food chain, from production to consumption (1). Urbanization and
27 modernization have profoundly changed eating habits (2). Diets first in Western industrialized
28 countries and now in many lower and middle-income countries are characterized by high consumption
29 of animal products but also salt, fat and sugar, and recognized as major risk factors for many chronic
30 diseases (3). Besides, the consumption of “ultra-processed foods” (UPF), has grown around the world
31 and now reaches more than half of daily energy intake in the UK and US (4) and is 31% in France (5).
32 These are foods that underwent extensive chemical or physical transformations and/or containing
33 cosmetic food additives or other industrials ingredients (e.g. hydrogenated oils, fructose syrup, etc.).
34 Although not systematically, they often contain on average higher amounts of saturated fat, salt, and
35 sugar and lower amounts of fibers, micronutrients and potentially healthy active compounds (6). In the
36 last decade, a growing body of studies suggesting a role of UPF consumption on health has emerged.
37 This literature is broadly growing, leading to the first reviews and meta-analyses summarizing findings
38 from prospective studies and consistently showing associations between UPF consumption and
39 increased risk of many non-communicable diseases (4,7–13). Most of the studies are based on the
40 NOVA classification distinguishing unprocessed, minimally processed, processed and ultra-processed
41 foods (6).

42 Beyond health issues, the current global food system greatly contributes to the degradation of the
43 environment by undermining natural resources, including water and forests, and jeopardizing climate
44 stability by increase of greenhouse gas emissions (GHGe) (14,15). However, even if the raw material
45 production stage is the most impacting for the environment, post-production steps (e.g. processing,
46 packaging, and transport) in the food industry are also resource-intensive (16,17) and are important
47 contributors of GHGe and energy demand. A recent time-series study conducted in Brazil showed that
48 the share of unprocessed foods in the diet decreased over a 30-year period, while the share of
49 processed and UPF increased, especially UPF based on animal products, doubling their contribution to
50 total diet-related environmental impacts over the same period (18).

51 The shortcomings of studies investigating environmental pressures associated with UPF have been
52 described in the literature and include, among other things, the failure to take into account some
53 ingredients and imprecise data on the processing and packaging stages (19). In addition, scientific
54 literature documenting environmental pressures related to UPF intakes is very scarce (20) and focused
55 on data from life cycle assessment (LCA) covering the entire chain, without individualizing its
56 different stages (production, processing, storage, packaging, transport and retailing).

57 However, some authors reported that consumption of discretionary foods (foods high in saturated fats,
58 sugars, salt and/or alcohol that can be eaten occasionally in small amounts, but are not a necessary part
59 of the diet) may contribute to an important part of diet-related environmental pressures (including
60 water use, energy use and GHGe). Energy density of such eating habits could be a strong determinant

61 of environmental pressures (21,22). In addition, the food processing sector is an important contributor
62 to total food loss and waste (along the entire food supply chain) generated by retailers and consumers
63 (23).

64 While the role of UPF on human health, based on the NOVA classification, is becoming well
65 documented, it is now important to estimate, using a systematic methodological approach, the impact
66 of UPF on planetary health, as part of a holistic approach to health.

67 In this context, the aim of the present study was to estimate environmental pressures of UPF
68 consumption, 1) overall, 2) adjusted for energy intake to account for level of consumption, 3)
69 differentiating the different stages of the whole food chain, and 4) by NOVA group, in a French
70 representative study that included a large set of environmental indicators.

Methods

Population

72 This study was based on the French nationally representative survey INCA 3 conducted in 2014-2015
73 by the French Agency for Food, Environmental and Occupational Health & Safety (ANSES) and
74 including 2,121 adult participants who provided valid dietary consumption data (24). The design of the
75 study as well as the methods have been detailed elsewhere (24).

76 Participants were selected according to a three-stage random sampling plan (geographical units,
77 dwellings then individuals) drawn at random by the National Institute of Statistics and Economic Studies
78 (INSEE), based on the annual population census in 2011. One individual per dwelling was then drawn
79 at random from among the eligible individuals at the time of contact with the household. Individual
80 weight was calculated according to INSEE method to improve representativeness according to region,
81 size of the urban area, occupation and socio-professional category of the household reference person,
82 size of the household, level of education, sex and age (25). The INCA 3 study protocol was authorised
83 by the National Commission on Informatics and Liberty, after a favourable opinion from the Advisory
84 Committee on Information Processing in Health Research (CCTIRS). The study also received a
85 favourable opinion from the Conseil National de l'Information Statistique (CNIS) on 15 June 2011
86 (n°121/D030) and was awarded the label of “general interest” and statistical quality by the INSEE Label
87 Committee (n°47/Label/D120). The data collected in the INCA 3 study are available on the website
88 [https://www.data.gouv.fr/fr/datasets/donnees-de-consommations-et-habitudes-alimentaires-de-letude-](https://www.data.gouv.fr/fr/datasets/donnees-de-consommations-et-habitudes-alimentaires-de-letude-inca-3/)
89 [inca-3/](https://www.data.gouv.fr/fr/datasets/donnees-de-consommations-et-habitudes-alimentaires-de-letude-inca-3/).

90 The data collected in the INCA 3 cross-sectional study included food and drink consumption and socio-
91 demographic and lifestyle characteristics.

Dietary data

93 Detailed consumption data were collected over 3 non-consecutive days (2 weekdays and 1 weekend
94 day) distributed over approximately 3 weeks, using the 24-hour recall method conducted by telephone
95 by trained interviewers using a standardised software (GloboDiet)(26). The quantification of portion
96 sizes was carried out using a picture booklet of food portions and household measurements.

97 Dietary intakes were calculated using the French nutritional composition data from the 2016 food
98 composition database published by the French Information Centre on Food Quality (27).

99 Mixed foods were decomposed using the standardized recipes validated by dietitians.

100 All food items were classified according to the NOVA classification (6,28) as previously extensively
101 described (29). At the individual level, the percentages (in weight) of the diet in NOVA1 (unprocessed
102 or minimally processed foods), NOVA2 (culinary ingredients), NOVA3 (processed foods), and
103 NOVA4 (UPF) were computed as described in the **Supplemental Material 1**.

104 The overall quality of the diet was assessed using two dietary scores, namely the sPNNS-GS2 (30) and
105 the PANDiet (31), which have extensively been described. Further details are presented in

106 **Supplemental Material 2**.

107 *Environmental indicators*

108 Diet-related environmental pressures were estimated using data from the French database Agribalyse®
109 3.0.1 developed by the French Agency for the Environment and Energy Management (ADEME).

110 Agribalyse® 3.0.1 contains environmental indicators for 2,497 foods consumed in France for which
111 nutritional contents is also available (32) using the same taxonomy. A total of 14 midpoint indicators
112 were available: GHGe, ozone depletion, particulate matters, ionizing radiation (effect on human
113 health), ecotoxicity, photochemical ozone formation (effect on human health), acidification, terrestrial
114 eutrophication, freshwater eutrophication, marine eutrophication, land use, water use, resource use,
115 minerals and metals and resource use, fossils and one endpoint ecological footprint (EF) calculated
116 according to the product environmental footprint (PEF) methodology (33).

117 Environmental indicator estimations were based on the method of LCA whose scope is "from field to
118 plate". The perimeter of the indicators covers each step of the value chain: agricultural production,
119 transport, processing, packaging, distribution and retailing, preparation at the consumer's level and
120 disposal of packaging. These different stages have been split into two phases 1) production and 2) post-
121 farm. The methodology has been extensively explained in *ad hoc* published reports (34,35) summarized
122 in **Supplemental Material 3**.

123 *Statistical analysis*

124 Participants were ranked and divided into weighted quintiles of %UPF. Socio-demographic and dietary
125 characteristics were described across weighted quintiles of %UPF using ANOVA, or ANCOVA models
126 when adjustment for energy intake was performed. Micronutrient and fiber intakes were adjusted for
127 energy intake using the residual method (36) and macronutrients were reported as % of total energy
128 intake.

129 Main food group contribution to %UPF by quintiles were also described.

130 In the main analysis, diet-related environmental footprints, according to quintiles of %UPF in the diet,
131 were first estimated overall, using crude and energy-adjusted ANOVA and ANCOVA models. For the
132 4 indicators that are well documented in the literature and therefore the most robust (GHGe, water use,
133 land use, and energy demand), we also examined the contribution of the different food system stages to

134 the environmental pressures. Finally, the respective contribution of the different NOVA food category
135 to the different environmental pressures was assessed.

136 Several sensitivity analyses were conducted: 1) models were adjusted for %NOVA1 in the diet as
137 NOVA1 consumption is inversely correlated to UPF, 2) UPF vs. %NOVA1 substitution was modeled
138 by adjustment for energy intake, %NOVA2 and %NOVA3 (36) for each indicator and farm and post-
139 farm steps, and 3) the main analysis described above was reperformed using the %UPF as % of total
140 energy intake instead of total weight. For this purpose, due to distribution of individual weightings,
141 quartiles were considered.

142 All tests were two-sided and a P-value <0.05 was considered significant. Statistical analyses were
143 performed using SAS Software (version 9.4, SAS Institute Inc, Cary, NC, USA) and figures were
144 performed developed using R version 3.6.

Results

145 The characteristics of the total study population and according to quintiles of %UPF are presented in
146 **Table 1**. The studied population included 2,121 participants (58% women), with a mean age of 47
147 years (SD=16). In this population, %UPF was 18.16 (SD=11.66) and 20.59 (SD=12.06) in women and
148 men, respectively. Participants with higher %UPF were more often male, younger, less educated, with
149 lower income and were more often unemployed or students. In addition, %UPF was inversely related
150 to proportion of NOVA1 in the diet (Q5 vs. Q1=-38%).

151 Dietary consumption greatly differed according to the %UPF in the diet (**Table 2 and Supplemental**
152 **Table 1**). In energy-adjusted models (**Table 2**), participants with high %UPF (Q5), compared to low
153 (Q1), had higher consumption of sweetened beverages, fruit juices, legumes, soup, and prepared dishes,
154 and lower consumption of non-sweetened beverages, whole grain and refined cereals, fruits, vegetables,
155 animal and vegetable fat, eggs, fish, and red meat. No linear relationship was observed for alcoholic
156 beverages, sweet and fat foods, condiments, potatoes, dairy products, processed meat, and poultry.

157 Energy intake increased across quintiles (Q5 vs. Q1=+22%) (**Table 3**). Higher %UPF was associated
158 with higher intake of carbohydrates and sugar, and lower intake of protein, fibers and most
159 micronutrients (total and non-heme iron, copper, magnesium, potassium, selenium, vitamins C and B12,
160 and zinc). Of note, for saturated fatty acids and lipids no linear trends were observed.

161 Overall, quality of the diet was lower in Q5 compared to Q1, with a decrease in sPNNS-GS2 (-77.5%)
162 and PANDiet (-4.6%) across quintiles (**Table 3**).

163 Overall, diet-related environmental pressures greatly differed depending on whether energy adjustment
164 was applied or not (**Table 4**).

165 In the unadjusted model, most indicators were higher among participants with higher %UPF, including
166 GHGe (Q5 vs. Q1: +15%), land use (+17%), fossils resource use (8%), marine and terrestrial
167 eutrophication (+13% and 15%), particulate matter (+13%), ionizing radiation (+16%) and the overall
168 endpoint ecological footprint (EF) score (+11%). On the contrary, water use was inversely associated
169 with %UPF with a lower mean value found in Q5 compared to Q1 (-7%).

170 When adjustment for energy intake was applied, some associations were no longer significant, including
171 GHGe while the negative association with regard to water use was slightly strengthened (Q5 vs. Q1
172 %UPF=-17%). In addition, associations regarding resource use, freshwater and marine eutrophication,
173 and ozone formation were reversed as well as the association concerning the global EF score (-6%).
174 GHGe, land use, energy demand and water use, according to quintiles of %UPF are presented by food
175 supply chain stages in **Figure 1** and **Supplemental Table 2** (energy-adjusted models). Overall, whatever
176 the quintile, agricultural production of raw material was the main driver for GHGe, land use and water
177 use, while packaging and processing stages were also important contributors to the energy demand
178 pressure. The substantial differences between %UPF quintiles (Q5 vs. Q1) were higher impacts of the
179 processing stage on water use (+53%), GHGe (+42%) and energy demand (+37%).
180 Concerning results by NOVA class (**Figure 2** and **Supplemental Table 3**), NOVA1 (unprocessed)
181 food consumption greatly contributed to each environmental indicator in all quintiles. However, the
182 contribution of UPF consumption to environmental pressures greatly differed according to quintile
183 and indicators. In Q1, GHGe, land use, energy demand and water use related to consumption of UPF
184 contributed to 10% to 13% while in Q5, contributions of NOVA4 consumption were 35%, 39%,
185 28% and 42% to total GHGe, water use, land use, and energy demand, respectively.
186 In sensitivity analyses, further adjustment for %NOVA1 in addition to energy intake (**Supplemental**
187 **Table 4**), yielded strengthened associations, with notably increases in GHGe (Q5 vs. Q1=+32%) and
188 land use (Q5 vs. Q1=+29%).
189 With the substitution model from NOVA1 (unprocessed) to UPF consumption (modeled as a continuous
190 variable) produced similar trends to the models adjusted for energy intake (**Figure 3**); with a decrease
191 in water use and an increase in GHGe and land use from production and high increase in energy demand
192 from processing.
193 Finally, results regarding % UPF as calorie are presented in **Supplemental Table 5**. Overall, the results
194 were similar but trend across quartiles was less evident. Notably, the association between GHGe and
195 level in UPF in the diet remained in the energy-adjusted model (+11%).

Discussion

196 In the present study conducted in a representative French population-based survey, we observed that
197 most of the studied environmental pressure indicators were higher among participants with higher
198 %UPF in their diet, in particular GHGe, energy demand and land use. A large part of the higher
199 pressures observed was explained by higher energy intake. Most environmental pressures occurred at
200 the stage of agricultural production, apart from energy demand for which the processing and
201 packaging stages was also an important contributor. NOVA1 food consumption highly contributed to
202 land use and GHGe in all quintiles. NOVA4 food consumption greatly contributed to energy demand
203 and to a lesser extent to water use with a strong gradient across quintiles.

204

205 To our knowledge, only one study previously investigated the environmental pressure of diet
206 according to the degree of food processing evaluated through the NOVA classification (20).
207 Comparison of our results with those of this study is not straightforward, since it has been conducted
208 in a Brazilian population, which may have exhibited some dissimilarities as regards dietary habits and
209 environmental pressures associated to food production and processing. As found herein, this study
210 reported difference in energy intake and GHGe according to level of UPF in the diet. Also, in our
211 study, the positive relationship between GHGe and %UPF did not remain significant after adjustment
212 for energy. In the study by Garzillo *et al* (20), the singular role of energy intake was not available as
213 the authors also adjusted for sociodemographic factors, which may be considered questionable
214 inasmuch as this is not supposed to confuse the relation when focusing on the link between dietary
215 patterns and environmental pressures. Moreover, when %UPF was expressed in kcal, as performed in
216 the study of Garzillo *et al.*, the associations between GHGe and quintiles of %UPF remained
217 statistically significant even in the energy-adjusted model. Besides, our findings related to water
218 footprint were not similar to those observed in the Brazilian study, since they documented a higher
219 food-related water use in participants with higher levels of UPF in their diet, which did not remain
220 after energy intake adjustment. In contrast, in our study, a higher water footprint from diet of
221 participants was observed among low %UPF participants, the association was even strengthened after
222 adjustment for energy intake. This latter result was attributable primarily to the fact that participants
223 who consumed lower amounts of UPF and, thus, in proportion, more NOVA1 food, tended to have
224 higher fruit and vegetable intakes and non-alcoholic beverages. This is also clearly observed in our
225 substitution model modelled with %UPF as a continuous variable. Thus, our finding is consistent with
226 the literature documenting a higher water footprint in plant-rich diets (37). Similarly, at constant
227 energy intake, the fact that the association between %UPF in the diet and GHGe and land use
228 disappeared, is related to the somewhat higher red meat, dairy products, fruit, vegetables, and fish
229 consumption (all classified as NOVA1) among low consumers of UPF compared to high consumers,
230 as previously reported (5). Of note meat, in particular ruminant meat, is the strongest contributor to
231 GHGe (17,38). These findings were consistent with the strong increase in environmental pressure with
232 %UPF when adjustment for NOVA1 was performed.

233 The present work is the first to explore the pressures associated with the production stage and all post-
234 farm stages including processing, packaging, transport and supply, according to the degree of
235 processing in the diet. Our results showed an important role of post-farm stages in energy demand
236 whatever the %UPF in the diet. The agricultural production stage is the main contributor to diet-
237 related GHGe, land and water use while with regard to energy demand, this is less evident since post-
238 farm stages play also an important role in the total pressure.

239 Our findings indicate that environmental impacts of UPF consumption could be linked to at least two
240 main factors. First, the higher energy intake of high consumers of UPF is a major determinant of their
241 diet-related impacts (39,40). Second, the higher number of post-farming stages for UPF production

242 can lead to an increase in energy demand and various environmental pressures. Indeed, UPF are, by
243 definition, related to more industrial processing, more packaging (for instance, they represent 2/3 and
244 more than 70% of packaged foods in France and in the US, respectively), and longer transport which
245 may substantially contribute to environmental impact of food (41). Consistently with the scientific
246 literature (16,17), even if the production stage is the most impacting stage for the environmental
247 resources depletion, it appeared that the food processing consumes a large amount of fossil resources.
248 On the contrary, it has been suggested that UPF (on a 100kcal basis) are less GHG emitting and
249 environmentally harmful (water and land use) than some minimally/unprocessed foods, in particular if
250 they contain small amount of animal ingredients (14,17,42). This last aspect is consistent with 1) our
251 results when dietary energy intake was accounted for, as most the pressures decreased across the
252 quintiles, and 2) with a recent British study analyzing foods properties that documented that processed
253 foods have lower nutritional quality, but also lower GHGe and were less expensive than minimally
254 processed foods, regardless of their total fat, salt and/or sugar content (42).

255 Studies that have explored the environmental pressures of UPF in details are very limited and none of
256 them considered the details of the post-agricultural stages, even though this element is essential to
257 accurately assess environmental impacts of such foods. In addition to potential human health benefits
258 (4,7–13), the reduction of high UPF consumption, associated with a greater overall consumption,
259 could be a driver in the transition towards a more sustainable food system by contributing to the
260 reduction of GHGe, energy demand, land use, soil and water degradation, and pollution. In addition, it
261 has been documented that production of UPF is also associated with the use of fertilizers and
262 pesticides, deforestation and biodiversity loss as well as packaging (2,19,43,44). With respect to our
263 findings as regards water use, the benefit is less convincing since, as previously emphasized (37),
264 healthy diets rich in fruits, vegetables and nuts are water consuming. Thus, some discrepancies exist in
265 alignment of foods as regards environmental sustainability and health impacts. For instance, it is now
266 well known that sugar, salt and food staples can have lower environmental impacts per calorie than
267 fruits, vegetables, and animal-based foods (45,46). The main lever to achieve food sustainability
268 remains the reduction of red meat and processed meat intake, as this would benefit for both human
269 health and the environment (14).

270 Our study exhibits some limitations and strengths. First, the study was based on diets of French adults
271 in a relatively small – though representative – sample, limiting the diversity of dietary patterns such as
272 vegetarian diets. For example, participants in INCA 3 had an average age of 47 years, whereas
273 consumers of UPF have been shown to be young (47). Second, limitations were somewhat inherent to
274 the matching between Agribalyse 3.01 and INCA 3, as food databases were independently developed,
275 and environmental indicators were not available for certain foods (for instance, culinary aids) or not
276 detailed (for instance, type of mushrooms). Third, even though the Agribalyse database is very rich
277 and accurate, some elements have been prioritized. For example, regarding packaging, only B2C
278 (business-to-consumer) packaging was considered and not B2B (business-to-business). Regarding

279 processing, the focus was driven by stages related to mass and yield changes and some other lacking
280 have been underlined (35). As regards transport, average values along the value chain were considered
281 but transport from the supply point to the household was not considered. Finally, the data were based
282 on LCA according to the standardized guidelines and methodologies but did not consider the type of
283 farming system (organic or conventional), limiting the consideration of the variety of practices along
284 the food chain. In addition, some indicators such as biodiversity loss were not available. As to the
285 strengths of our study, dietary data was collected in a nationally representative sample of the adult
286 population of France in 2015-2016. Furthermore, the consumption and pressures data were collected
287 using standardized methodologies. In addition, environmental data were validated by several expert
288 entities (35). Finally, the detail of pressures by stage of the value chain allowed us to consider for the
289 first time the contribution of UPF while taking into the different stages.

Conclusion

290 This study is the first to explore the contribution of UPF consumption to different environmental
291 pressures, while detailing the different stages of the food chain. As high consumers of ultra-processed
292 foods generally have higher energy intake, overall their diets were associated with a higher footprint.
293 In addition, the consumption of UPF had a substantial role on some indicators, particularly on energy
294 demand through the processing stage. Such investigations could be considered in the development of
295 sustainable dietary guidelines in light of the previously documented links between UPF and human
296 health.

The authors' contributions

298 EKG conducted the research, implemented the databases, conducted the analyses and wrote the
299 manuscript.

300 All authors critically helped in the interpretation of results, revised the manuscript and provided relevant
301 intellectual input. They all read and approved the final manuscript.

302 EKG had primary responsibility for the final content, she is the guarantor.

Conflict of Interest

303 FM was the scientific leader of a PhD project funded until October 2021 in part by a research contract
304 with Terres Univia, the French Interbranch organization for plant oils and proteins. The other authors
305 declared no conflict of interest.

Acknowledgment

306 The authors are indebted to Vincent Colomb and Mélissa Cornélius (ADEME) for their support in the
307 use of the Agribalyse database ®.

References

- 308 1. Lindgren E, Harris F, Dangour AD, Gasparatos A, Hiramatsu M, Javadi F, et al. Sustainable
309 food systems-a health perspective. *Sustain Sci*. 2018;13(6):1505–17.
- 310 2. Baker P, Machado P, Santos T, Sievert K, Backholer K, Hadjidakou M, et al. Ultra-processed
311 foods and the nutrition transition: Global, regional and national trends, food systems
312 transformations and political economy drivers. *Obes Rev*. 2020 Dec;21(12):e13126.
- 313 3. Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the
314 Anthropocene: the EAT-Lancet Commission on healthy diets from sustainable food systems.
315 *Lancet*. 2019 Jan 16;393(10170):447–92.
- 316 4. Srour B, Touvier M. Ultra-processed foods and human health: What do we already know and
317 what will further research tell us? *EClinicalMedicine*. 2021 Feb;32:100747.
- 318 5. Salomé M, Arrazat L, Wang J, Dufour A, Dubuisson C, Volatier JL, et al. Contrary to ultra-
319 processed foods, the consumption of unprocessed or minimally processed foods is associated
320 with favorable patterns of protein intake, diet quality and lower cardiometabolic risk in French
321 adults (INCA3). *Eur J Nutr*. 2021 Oct;60(7):4055–67.
- 322 6. Monteiro CA, Cannon G, Moubarac JC, Levy RB, Louzada MLC, Jaime PC. The UN Decade of
323 Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health*
324 *Nutr*. 2018 Jan;21(1):5–17.
- 325 7. Taneri PE, Wehrli F, Roa Diaz ZM, Itodo OA, Salvador D, Raeisi-Dehkordi H, et al. Association
326 Between Ultra-Processed Food Intake and All-Cause Mortality: A Systematic Review and Meta-
327 Analysis. *Am J Epidemiol*. 2022 Mar 1;kwac039.
- 328 8. Suksatan W, Moradi S, Naeini F, Bagheri R, Mohammadi H, Talebi S, et al. Ultra-Processed
329 Food Consumption and Adult Mortality Risk: A Systematic Review and Dose-Response Meta-
330 Analysis of 207,291 Participants. *Nutrients*. 2021 Dec 30;14(1):174.
- 331 9. Jardim MZ, Costa BV de L, Pessoa MC, Duarte CK. Ultra-processed foods increase
332 noncommunicable chronic disease risk. *Nutr Res*. 2021 Nov;95:19–34.
- 333 10. Delpino FM, Figueiredo LM, Bielemann RM, da Silva BGC, Dos Santos FS, Mintem GC, et al.
334 Ultra-processed food and risk of type 2 diabetes: a systematic review and meta-analysis of
335 longitudinal studies. *Int J Epidemiol*. 2021 Dec 14;dyab247.
- 336 11. Moradi S, Hojjati Kermani MA, Bagheri R, Mohammadi H, Jayedi A, Lane MM, et al. Ultra-
337 Processed Food Consumption and Adult Diabetes Risk: A Systematic Review and Dose-
338 Response Meta-Analysis. *Nutrients*. 2021 Dec 9;13(12):4410.
- 339 12. Dicken SJ, Batterham RL. The Role of Diet Quality in Mediating the Association between Ultra-
340 Processed Food Intake, Obesity and Health-Related Outcomes: A Review of Prospective Cohort
341 Studies. *Nutrients*. 2021 Dec 22;14(1):23.
- 342 13. Chen X, Zhang Z, Yang H, Qiu P, Wang H, Wang F, et al. Consumption of ultra-processed
343 foods and health outcomes: a systematic review of epidemiological studies. *Nutr J*. 2020 Aug
344 20;19(1):86.
- 345 14. Clark MA, Springmann M, Hill J, Tilman D. Multiple health and environmental impacts of
346 foods. *PNAS* [Internet]. 2019 Nov 12 [cited 2020 Mar 27];116(46):23357–62. Available from:
347 <https://www.pnas.org/content/116/46/23357>

- 348 15. HLPE. Nutrition and food systems. A report by the High Level Panel of Experts on Food
349 Security and Nutrition of the Committee on World Food Security,. Rome, Italy; 2017 p. 152.
350 Report No.: 12.
- 351 16. Crippa M, Solazzo E, Guizzardi D, Monforti-Ferrario F, Tubiello FN, Leip A. Food systems are
352 responsible for a third of global anthropogenic GHG emissions. *Nat Food* [Internet]. 2021 Mar
353 [cited 2021 Sep 28];2(3):198–209. Available from: [https://www.nature.com/articles/s43016-021-](https://www.nature.com/articles/s43016-021-00225-9)
354 [00225-9](https://www.nature.com/articles/s43016-021-00225-9)
- 355 17. Poore J, Nemecek T. Reducing food’s environmental impacts through producers and consumers.
356 *Science*. 2018 Jun 1;360(6392):987–92.
- 357 18. da Silva JT, Garzillo JMF, Rauber F, Kluczkovski A, Rivera XS, da Cruz GL, et al. Greenhouse
358 gas emissions, water footprint, and ecological footprint of food purchases according to their
359 degree of processing in Brazilian metropolitan areas: a time-series study from 1987 to 2018.
360 *Lancet Planet Health*. 2021 Nov;5(11):e775–85.
- 361 19. Seferidi P, Scrinis G, Huybrechts I, Woods J, Vineis P, Millett C. The neglected environmental
362 impacts of ultra-processed foods. *Lancet Planet Health*. 2020 Oct;4(10):e437–8.
- 363 20. Garzillo JMF, Poli VFS, Leite FHM, Steele EM, Machado PP, Louzada ML da C, et al. Ultra-
364 processed food intake and diet carbon and water footprints: a national study in Brazil. *Rev Saude*
365 *Publica* [Internet]. [cited 2022 Apr 1];56:6. Available from:
366 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8859933/>
- 367 21. Barré T, Perignon M, Gazan R, Vieux F, Micard V, Amiot MJ, et al. Integrating nutrient
368 bioavailability and co-production links when identifying sustainable diets: How low should we
369 reduce meat consumption? *PLoS ONE*. 2018;13(2):e0191767.
- 370 22. Hadjikakou M. Trimming the excess: environmental impacts of discretionary food consumption
371 in Australia. *Ecological Economics* [Internet]. 2017 [cited 2022 Apr 1];131(C):119–28.
372 Available from: <https://ideas.repec.org/a/eee/ecolec/v131y2017icp119-128.html>
- 373 23. Lemaire A, Limbourg S. How can food loss and waste management achieve sustainable
374 development goals? *Journal of Cleaner Production* [Internet]. 2019 Oct 10 [cited 2022 Apr
375 1];234:1221–34. Available from:
376 <https://www.sciencedirect.com/science/article/pii/S0959652619321912>
- 377 24. Dubuisson C, Dufour A, Carrillo S, Drouillet-Pinard P, Havard S, Volatier JL. The Third French
378 Individual and National Food Consumption (INCA3) Survey 2014-2015: method, design and
379 participation rate in the framework of a European harmonization process. *Public Health Nutr*.
380 2019 Mar;22(4):584–600.
- 381 25. Sautory O. INSEE: La macro CALMAR-Redressement d’un échantillon par calage sur marges.
382 <http://www.insee.fr/fr/methodes/outils/calmar/doccalmar.pdf>. 1993.
- 383 26. Aglago EK, Landais E, Nicolas G, Margetts B, Leclercq C, Allemand P, et al. Evaluation of the
384 international standardized 24-h dietary recall methodology (GloboDiet) for potential application
385 in research and surveillance within African settings. *Globalization and Health* [Internet]. 2017
386 Jun 19 [cited 2021 Nov 9];13(1):35. Available from: <https://doi.org/10.1186/s12992-017-0260-6>
- 387 27. Agence nationale de sécurité sanitaire de l’alimentation de l’environnement et du travail
388 (ANSES). French food composition table (CIQUAL) [Internet]. ANSES. 2012. Available from:
389 <http://www.afssa.fr/TableCIQUAL> (accessed 29 september 2012)

- 390 28. Monteiro CA, Cannon G, Levy RB, Moubarac JC, Louzada ML, Rauber F, et al. Ultra-processed
391 foods: what they are and how to identify them. *Public Health Nutr.* 2019;22(5):936–41.
- 392 29. Srouf B, Fezeu LK, Kesse-Guyot E, Allès B, Debras C, Druesne-Pecollo N, et al. Ultraprocessed
393 Food Consumption and Risk of Type 2 Diabetes Among Participants of the NutriNet-Santé
394 Prospective Cohort. *JAMA Intern Med.* 2019 Dec 16;
- 395 30. Chaltiel D, Adjibade M, Deschamps V, Touvier M, Hercberg S, Julia C, et al. Programme
396 National Nutrition Santé – guidelines score 2 (PNNS-GS2): development and validation of a diet
397 quality score reflecting the 2017 French dietary guidelines. *British Journal of Nutrition*
398 [Internet]. 2019 Aug [cited 2019 Sep 2];122(3):331–42. Available from:
399 [https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/programme-](https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/programme-national-nutrition-sante-guidelines-score-2-pnns-gs2-development-and-validation-of-a-diet-quality-score-reflecting-the-2017-french-dietary-guidelines/BA866BAC50F0C9154BF8FF235B5A99DD)
400 [national-nutrition-sante-guidelines-score-2-pnns-gs2-development-and-validation-of-a-diet-](https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/programme-national-nutrition-sante-guidelines-score-2-pnns-gs2-development-and-validation-of-a-diet-quality-score-reflecting-the-2017-french-dietary-guidelines/BA866BAC50F0C9154BF8FF235B5A99DD)
401 [quality-score-reflecting-the-2017-french-dietary-](https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/programme-national-nutrition-sante-guidelines-score-2-pnns-gs2-development-and-validation-of-a-diet-quality-score-reflecting-the-2017-french-dietary-guidelines/BA866BAC50F0C9154BF8FF235B5A99DD)
402 [guidelines/BA866BAC50F0C9154BF8FF235B5A99DD](https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/programme-national-nutrition-sante-guidelines-score-2-pnns-gs2-development-and-validation-of-a-diet-quality-score-reflecting-the-2017-french-dietary-guidelines/BA866BAC50F0C9154BF8FF235B5A99DD)
- 403 31. Gavelle E de, Huneau JF, Mariotti F. Patterns of Protein Food Intake Are Associated with
404 Nutrient Adequacy in the General French Adult Population. *Nutrients.* 2018 Feb 17;10(2).
- 405 32. Colomb V, A. Colsaet, S. Ait-Amar, C. Basset-Mens, G. Mevel, V. To, et al. AGRIBALYSE:
406 the French public LCI database for agricultural products. 2015 [cited 2017 Sep 28]; Available
407 from: <http://rgdoi.net/10.13140/RG.2.1.2586.0240>
- 408 33. European Commission. Guidance for the development of Product Environmental Footprint
409 Category Rules (PEFCRs) - version 6.3. [Internet]. 2018. Available from:
410 https://ec.europa.eu/environment/eussd/smgp/pdf/PEFCR_guidance_v6.3.pdf
- 411 34. Koch P, Salou T. AGRIBALYSE®: Rapport Méthodologique- Volet Agriculture- Version 3.0
412 [Internet]. Angers, France: ADEME; 2020 p. 384. Available from: Available at:
413 www.ademe.fr/agribalyse-en.
- 414 35. AGRIBALYSE 3.0, the French agricultural and food LCI database, methodology for food
415 products [Internet]. ADEME; 2020 May [cited 2022 Oct 3] p. 85. Available from:
416 <https://doc.agribalyse.fr/documentation-en/agribalyse-data/documentation>
- 417 36. Willett WC. *Nutritional Epidemiology*. Third Edition, New to this Edition: Oxford, New York:
418 Oxford University Press; 2012. 552 p. (Monographs in Epidemiology and Biostatistics).
- 419 37. Fresán U, Sabaté J. Vegetarian Diets: Planetary Health and Its Alignment with Human Health.
420 *Adv Nutr* [Internet]. 2019 Nov 1 [cited 2020 Jun 5];10(Supplement_4):S380–8. Available from:
421 https://academic.oup.com/advances/article/10/Supplement_4/S380/5624053
- 422 38. Clark M, Hill J, Tilman D. The Diet, Health, and Environment Trilemma. *Annual Review of*
423 *Environment and Resources* [Internet]. 2018 [cited 2020 Mar 26];43(1):109–34. Available from:
424 <https://doi.org/10.1146/annurev-environ-102017-025957>
- 425 39. Vieux F, Darmon N, Touazi D, Soler L, Soler LG. Greenhouse gas emissions of self-selected
426 individual diets in France: Changing the diet structure or consuming less? *EcolEcon.*
427 2012;75:91–101.
- 428 40. Hendrie GA, Baird D, Ridoutt B, Hadjikakou M, Noakes M. Overconsumption of Energy and
429 Excessive Discretionary Food Intake Inflates Dietary Greenhouse Gas Emissions in Australia.
430 *Nutrients.* 2016 Oct 31;8(11):E690.

- 431 41. Fardet A, Rock E. Ultra-Processed Foods and Food System Sustainability: What Are the Links?
432 Sustainability [Internet]. 2020 Jan [cited 2022 Apr 14];12(15):6280. Available from:
433 <https://www.mdpi.com/2071-1050/12/15/6280>
- 434 42. Aceves-Martins M, Bates RL, Craig LCA, Chalmers N, Horgan G, Boskamp B, et al. Nutritional
435 Quality, Environmental Impact and Cost of Ultra-Processed Foods: A UK Food-Based Analysis.
436 International Journal of Environmental Research and Public Health [Internet]. 2022 Jan [cited
437 2022 Apr 1];19(6):3191. Available from: <https://www.mdpi.com/1660-4601/19/6/3191>
- 438 43. Li D, Wang X, Chan HK, Manzini R. Sustainable food supply chain management. International
439 Journal of Production Economics [Internet]. 2014 Jun 1 [cited 2022 Apr 14];152:1–8. Available
440 from: <https://www.sciencedirect.com/science/article/pii/S0925527314001133>
- 441 44. Leite FHM, Khandpur N, Andrade GC, Anastasiou K, Baker P, Lawrence M, et al. Ultra-
442 processed foods should be central to global food systems dialogue and action on biodiversity.
443 BMJ Global Health [Internet]. 2022 Mar 1 [cited 2022 Apr 14];7(3):e008269. Available from:
444 <https://gh.bmj.com/content/7/3/e008269>
- 445 45. Vieux F, Soler LG, Touazi D, Darmon N. High nutritional quality is not associated with low
446 greenhouse gas emissions in self-selected diets of French adults. *AmJ Clin Nutr*. 2013
447 Mar;97(3):569–83.
- 448 46. Tilman D, Clark M. Global diets link environmental sustainability and human health. *Nature*.
449 2014 Nov 27;515(1476-4687 (Electronic)):518–22.
- 450 47. Gehring J, Touvier M, Baudry J, Julia C, Buscail C, Srour B, et al. Consumption of Ultra-
451 Processed Foods by Pesco-Vegetarians, Vegetarians, and Vegans: Associations with Duration
452 and Age at Diet Initiation. *J Nutr* [Internet]. [cited 2020 Aug 24]; Available from:
453 <https://academic.oup.com/jn/advance-article/doi/10.1093/jn/nxaa196/5874423>
- 454
- 455

Table 1: Characteristics of study participants according to quintiles of %UPF, (INCA 3, n=2,121)¹

	Total sample %	%UPF	Q1	Q2	Q3	Q4	Q5	P for trend ²
%UPF cut-off³	-		0-<9.95	9.95-<14.68	14.68-<20.25	20.25-<28.44	>28.44	
%NOVA1	69.83 (10.42)		82.67 (6.29)	76.4 (6.54)	70.17 (7.30)	65.86 (6.33)	51.46 (10.42)	<.0001
%NOVA2	1.50 (0.01)		1.50 (1.06)	1.52 (1.00)	1.49 (1.15)	1.52 (1.22)	1.47 (1.09)	<.0001
%NOVA3	9.43 (0.43)		9.37 (5.40)	9.66 (6.06)	10.97 (7.00)	8.61 (5.80)	8.39 (5.84)	0.84
%UPF	19.29 (10.86)		6.56 (2.29)	12.43 (1.3)	17.4 (1.51)	24.02 (2.47)	38.78 (10.38)	<.0001
N	2,121	-	471	420	429	393	408	
Weighted N	2,121	-	421.36	425.13	425.86	422.35	426.30	
Sex								<.0001
Men	41.82	20.59 (12.06)	38.96	45.53	51.02	53.14	53.71	
Women	58.18	18.16 (11.66)	61.04	54.47	48.98	46.86	46.29	
Age (y)	46.9 (16.3)	-	54.5 (13.6)	51.2 (15.0)	50.0 (15.1)	49.2 (17)	47.29 (18.1)	<.0001
Geographic area								0.09
Ile-de-France	12.35	18.18 (11.55)	21.31	19.49	20.13	11	10.07	
North-West	21.97	19.42 (11.76)	19.98	17.54	20.24	19.21	22.03	
North-East	24.38	21.36 (13.14)	21.17	25.71	22.2	29.16	38.95	
South-East	22.30	18.15 (11.23)	18.76	20.53	18.49	19.23	16.15	
South-West	19.00	17.95 (10.91)	18.77	16.74	18.94	21.41	12.8	
Professional position								0.45
Employees	15.79	19.96 (12.89)	16.22	14.18	15.96	16.64	17.41	
Workers	9.29	21.19 (13.70)	14.4	11.58	13.42	14.18	16.47	
Farmers	0.75	19.47 (10.68)	0.58	0.73	1.00	0.67	0.24	
Craftsmen and company managers	2.64	19.38 (10.99)	4.58	1.73	4.64	3.39	1.23	
Intermediate professions	19.00	18.62 (11.18)	19.6	24.44	18.22	18.67	14.61	
Executive, liberal professions	13.06	17.34 (9.86)	11.74	14.42	13.18	12.31	7.98	
Retired individuals	30.60	18.42 (11.62)	26.96	22.84	22.87	21.74	21.07	
Other inactive individuals	8.86	22.12 (12.70)	5.92	10.07	10.71	12.38	20.98	
Education								0.09
Primary+College	37.86	19.90 (12.88)	48.72	50.07	46.77	47.09	47.14	
High school	20.79	20.77 (13.03)	16.8	14.94	14.1	21.4	24.6	
Undergraduate level	21.12	18.83 (10.51)	16.33	16.18	17.46	16.57	18.12	
Postgraduate level	20.18	16.56 (9.41)	18.16	18.41	21.67	14.94	10.14	
No information	0.05	13.27 (.)	0	0.41	0	0	0	
Employment status								0.01
Employed	53.84	18.80 (11.43)	62.22	59.28	58.94	51.62	47.5	
Unemployed	6.84	21.76 (14.41)	5.63	7.11	7.47	14.95	10.45	
Student	3.68	25.25 (11.44)	0.85	3.63	4.59	7.32	11.45	
Retired (former employee)	28.85	18.34 (11.56)	25.9	20.79	21.25	20.09	20.26	
Retired from business	1.74	19.60 (12.64)	1.06	2.06	1.62	1.65	0.81	
Housewife or man	3.25	20.12 (14.19)	3.43	4.88	4.45	2.51	7.53	
Other inactive	1.74	19.66 (11.48)	0.92	1.51	1.67	1.86	2.01	
No information	0.05	12.11 (.)	0	0.75	0	0	0	
Marital status								0.41
Single	15.61	18.08 (10.49)	11.01	14.43	16.3	11.59	12.92	
Married	37.01	19.14 (11.92)	38.39	30.84	38.19	32.58	37.39	
Unmarried couple	10.94	18.81 (11.57)	10.35	10.37	10.57	8.11	13.26	
Widowed	28.76	19.67 (12.31)	29.03	33.58	27.03	34.85	29.29	
Separated/Divorced	6.36	20.13 (12.73)	7.98	7.87	7.9	12.56	4.93	
No information	1.27	20.25 (14.92)	3.24	2.92	0	0.31	2.05	
Living area								0.23
Rural	26.97	19.60 (12.18)	26.64	24.44	22.87	28.53	26.07	
2,000-19,999 inhab.	19.42	19.57 (13.08)	18.18	18.78	17.9	17.75	16.88	
20,000-99,999 inhab.	13.81	18.83 (10.53)	12.11	10.03	11.7	13.42	13.68	
≥100,000 inhab.	28.95	19.01 (11.70)	28.68	31.38	31.66	31.76	34.69	
Paris area	10.84	18.30 (11.02)	14.38	15.38	15.86	8.54	8.68	
Income (euros/month)								0.69
<1300	16.22	20.60 (13.15)	17.42	17.47	18.87	21.06	23.49	
1300-<1900	17.26	19.01 (11.80)	12.65	17.64	16.35	21.87	13.73	

1900-<2500	30.55	19.19 (11.83)	20.79	17.69	17.18	16.09	17.38	
2500-<4600	18.81	19.05 (11.37)	31.71	30.39	30.02	29.91	28.52	
>4600	9.48	16.40 (10.18)	11.77	11	11.82	5.5	5.79	
No information	7.69	19.96 (12.43)	5.66	5.8	5.75	5.58	11.09	
BMI (kg.m²)	25.86 (4.82)	-	25.96 (4.67)	25.64 (4.59)	25.8 (4.57)	25.75 (5.02)	25.65 (5.25)	0.05

456 Abbreviations: BMI, body mass index; inhab: inhabitants

457 ¹Values are n, means (SD) or % as appropriate, all data are weighted

458 ²P-value for linear trend is estimated using linear contrast

459 ³Values are ranges of %UPF (in weight)

Table 2: Consumption of food groups (g/d) according to %UPF quintiles, (INCA 3, N=2,121)¹

	<i>Total sample</i>	Q1	Q2	Q3	Q4	Q5	<i>ΔQ5 vs. Q1²</i>	<i>P for trend³</i>
Alcoholic beverages	122.0 (92.7)	106.6 (11.1)	131.4 (11.0)	122.3 (11.0)	109.9 (11.0)	141.2 (11.0)	32	0.18
Sweetened beverages	105.4 (94.7)	29.7 (10.9)	41.3 (10.8)	44.7 (10.8)	117.8 (10.9)	315.8 (10.8)	963	<.0001
Non-sweetened beverages	1,394.2 (320.0)	1,797.4 (30.8)	1,583.4 (30.4)	1,385.4 (30.6)	1,229.0 (30.6)	898.8 (30.6)	-50	<.0001
Sweet and fat foods	109.5 (50.5)	97.2 (3.5)	112.0 (3.4)	103.7 (3.5)	119.8 (3.5)	116.3 (3.5)	20	0.10
Condiments	26.5 (4.7)	24.4 (1.5)	25.3 (1.5)	27.6 (1.5)	30.2 (1.5)	25.2 (1.5)	4	0.16
Whole grains	6.5 (1.3)	7.4 (1.0)	7.6 (1.0)	7.4 (1.0)	5.6 (1.0)	4.2 (1.0)	-43	0.01
Refined cereals	181.3 (62.8)	206.8 (5.0)	183.3 (4.9)	196.3 (4.9)	159.1 (4.9)	156.8 (4.9)	-24	<.0001
Fruits	144.1 (33.0)	186.3 (6.6)	161.3 (6.5)	147.7 (6.5)	116.9 (6.6)	100.5 (6.6)	-46	<.0001
Fruit juice	62.6 (16.1)	47.8 (4.8)	57.7 (4.8)	66.2 (4.8)	65.4 (4.8)	78.3 (4.8)	64	<.0001
Vegetables	141.6 (32.4)	186.2 (5.4)	157.9 (5.3)	139.1 (5.4)	121.9 (5.4)	94.9 (5.4)	-49	<.0001
Legumes	7.4 (5.0)	6.2 (1.2)	7.0 (1.2)	5.1 (1.2)	10.7 (1.2)	8.5 (1.2)	39	0.03
Soup	98.5 (46.7)	26.8 (8.3)	60.1 (8.2)	117.8 (8.3)	154.0 (8.3)	146.3 (8.3)	447	<.0001
Potatoes	45.9 (19.1)	47.6 (3.5)	41.5 (3.4)	47.5 (3.4)	42.3 (3.4)	50.7 (3.4)	7	0.52
Nuts	3.2 (1.3)	4.1 (0.4)	3.2 (0.4)	3.2 (0.4)	2.9 (0.4)	2.5 (0.4)	-38	0.01
Animal fat	9.4 (3.3)	10.7 (0.7)	9.0 (0.6)	9.7 (0.6)	9.6 (0.6)	7.8 (0.6)	-27	0.01
Vegetable fat	8.2 (1.5)	10.1 (0.5)	8.4 (0.5)	6.8 (0.5)	7.5 (0.5)	8.4 (0.5)	-17	<.0001
Prepared dishes	114.2 (26.8)	109.3 (5.4)	110.7 (5.3)	104.5 (5.4)	122.0 (5.4)	125.9 (5.4)	15	0.01
Dairy products	185.2 (30.9)	183.8 (8.4)	187.3 (8.3)	186.6 (8.3)	205.8 (8.3)	161.8 (8.3)	-12	0.34
Eggs	13.9 (1.7)	17.6 (1.3)	14.8 (1.3)	10.9 (1.3)	13.8 (1.3)	11.8 (1.3)	-33	<.0001
Fish	28.3 (3.9)	35.5 (2.0)	29.8 (1.9)	25.1 (1.9)	24.4 (1.9)	25.6 (1.9)	-28	<.0001
Red meat	51.0 (16.0)	55.5 (2.7)	52.9 (2.6)	49.7 (2.7)	49.6 (2.7)	46.4 (2.7)	-17	0.01
Processed meat	30.3 (14.4)	27.8 (1.8)	29.2 (1.7)	32.4 (1.7)	31.2 (1.8)	31.4 (1.8)	13	0.10
Poultry	26.9 (3.7)	28.3 (1.8)	26.4 (1.8)	29.2 (1.8)	22.6 (1.8)	27.7 (1.8)	-2	0.37

460 ¹Values are means (standard errors of the means) adjusted for total energy intake

461 ²Relative difference

462 ³P-value for linear trend is estimated using linear contrast

Table 3: Nutrient intakes and dietary scores according to %UPF quintiles, (INCA 3, n=2,121)¹

	Q1	Q2	Q3	Q4	Q5	P for trend ⁴
Alcohol-free energy intake (kcal/d) ²	1,782.70 (34.81)	1,949.69 (34.66)	2,133.58 (34.63)	2,110.19 (34.77)	2,178.44 (34.61)	<.0001
Lipids ³ (%EI/d)	33.40 (0.31)	32.01 (0.30)	31.74 (0.31)	32.83 (0.31)	32.32 (0.31)	0.17
Proteins ³ (%EI/d)	38.26 (0.39)	36.79 (0.38)	36.50 (0.38)	36.22 (0.38)	35.10 (0.38)	<.0001
Carbohydrates ³ (%EI/d)	97.95 (0.82)	101.38 (0.80)	101.91 (0.81)	102.86 (0.81)	105.36 (0.81)	<.0001
Sugar ³ (%EI/d)	16.12 (0.27)	16.99 (0.27)	17.10 (0.27)	18.47 (0.27)	20.03 (0.27)	<.0001
Animal protein ³ (%EI/d)	25.94 (0.40)	24.23 (0.40)	23.60 (0.40)	23.44 (0.40)	22.75 (0.40)	<.0001
Plant protein ³ (%EI/d)	12.90 (0.15)	13.27 (0.15)	13.44 (0.15)	13.38 (0.15)	13.00 (0.15)	0.51
SFA (g/d)	31.47 (0.38)	31.44 (0.38)	31.62 (0.38)	32.68 (0.38)	30.83 (0.38)	0.96
Atherogenic SFA ⁵ (g/d)	20.25 (0.25)	20.26 (0.25)	20.00 (0.25)	20.78 (0.25)	19.74 (0.25)	0.54
EPA+DHA (g/d)	0.31 (0.02)	0.34 (0.02)	0.25 (0.02)	0.25 (0.02)	0.28 (0.02)	0.03
Linoleic FA (g/d)	7.11 (0.13)	6.81 (0.13)	6.57 (0.13)	6.89 (0.13)	6.91 (0.13)	0.45
Alpha-linolenic FA (g/d)	1.03 (0.03)	1.01 (0.02)	0.96 (0.02)	0.92 (0.02)	0.93 (0.02)	<.0001
PUFA (g/d)	10.08 (0.17)	9.54 (0.17)	9.20 (0.17)	9.41 (0.17)	9.75 (0.17)	0.15
Beta-carotene (µg/d)	2737 (129)	2563 (127)	2890 (128)	2715 (121)	2445 (128)	0.29
Calcium (mg/d)	921.50 (15.06)	909.39 (14.86)	913.67 (14.92)	948.43 (14.94)	845.54 (14.93)	0.02
Cholesterol (mg/d)	340.33 (6.56)	329.32 (6.47)	315.25 (6.50)	323.13 (6.51)	293.95 (6.51)	<.0001
Copper (mg/d)	1.85 (0.04)	1.76 (0.04)	1.81 (0.04)	1.53 (0.04)	1.44 (0.04)	<.0001
Iron (mg/d)	10.47 (0.13)	10.78 (0.13)	10.58 (0.13)	10.10 (0.13)	10.02 (0.13)	<.0001
Non-heme iron (mg/d)	9.26 (0.11)	9.55 (0.11)	9.29 (0.11)	8.85 (0.11)	8.82 (0.11)	<.0001
Zinc (mg/d)	9.70 (0.15)	9.76 (0.15)	9.47 (0.15)	9.41 (0.15)	8.64 (0.15)	<.0001
Fiber (g/d)	20.42 (0.25)	20.04 (0.24)	20.12 (0.24)	19.58 (0.24)	18.23 (0.24)	<.0001
Iodine (µg/d)	147.58 (2.72)	153.14 (2.68)	148.20 (2.70)	156.15 (2.70)	143.66 (2.70)	0.58
Magnesium (mg/d)	354.69 (4.26)	355.80 (4.21)	336.18 (4.22)	331.81 (4.23)	316.67 (4.23)	<.0001
Manganese (mg/d)	3.48 (0.05)	3.22 (0.05)	3.18 (0.05)	2.76 (0.05)	2.66 (0.05)	<.0001
Phosphorus (mg/d)	1,221 (11)	1,206 (11)	1,185 (11)	1,179 (11)	1,136 (11)	<.0001
Phytates (µg/d)	618.87 (9.67)	612.97 (9.54)	594.88 (9.58)	579.84 (9.59)	569.60 (9.59)	<.0001
Polyols (mg/d)	1.23 (0.06)	1.26 (0.06)	1.19 (0.06)	0.97 (0.06)	0.94 (0.06)	<.0001
Potassium (mg/d)	3,226 (35)	3,188 (34)	3,054 (34)	3,028 (34)	2,873 (34)	<.0001
Retinol (µg/d)	474.30 (33.43)	491.88 (32.98)	500.93 (33.11)	435.60 (33.16)	377.45 (33.15)	0.02
Selenium (µg/d)	130.67 (2.04)	123.18 (2.01)	125.71 (2.02)	122.92 (2.02)	113.86 (2.02)	<.0001
Sodium (mg/d)	2,978 (43)	3,033 (43)	3,263 (43)	3,273 (43)	3,059 (43)	<.0001
Vitamin B1 (mg/d)	1.20 (0.02)	1.20 (0.02)	1.21 (0.02)	1.23 (0.02)	1.18 (0.02)	0.87
Vitamin B2 (mg/d)	1.80 (0.03)	1.80 (0.03)	1.74 (0.03)	1.83 (0.03)	1.75 (0.03)	0.52
Vitamin B3 (mg/d)	20.45 (0.35)	20.56 (0.35)	19.70 (0.35)	19.46 (0.35)	20.14 (0.35)	0.13
Vitamin B5 (mg/d)	5.58 (0.07)	5.68 (0.07)	5.52 (0.07)	5.60 (0.07)	5.46 (0.07)	0.16
Vitamin B6 (mg/d)	1.71 (0.02)	1.71 (0.02)	1.70 (0.02)	1.67 (0.02)	1.70 (0.02)	0.41
Vitamin B9 (µg/d)	308.39 (4.24)	306.96 (4.18)	300.22 (4.20)	293.86 (4.21)	279.98 (4.20)	<.0001
Vitamin B12 (µg/d)	5.58 (0.20)	5.64 (0.20)	5.32 (0.20)	4.79 (0.20)	4.62 (0.20)	<.0001
Vitamin C (mg/d)	95.89 (2.65)	89.08 (2.61)	93.03 (2.62)	85.96 (2.63)	81.68 (2.62)	<.0001
Vitamin E (mg/d)	10.37 (0.18)	9.71 (0.17)	9.14 (0.17)	9.31 (0.17)	9.59 (0.17)	<.0001
PANDiet (/100)	59.64 (0.28)	59.59 (0.28)	59.08 (0.28)	57.99 (0.28)	56.89 (0.28)	<.0001
sPNNS-GS2 (/14.25)	1.42 (0.13)	1.07 (0.12)	0.55 (0.13)	0.36 (0.13)	0.32 (0.13)	<.0001

463 Abbreviations: EI, energy intake; DHA, docosahexaenoic acid; EPA, eicosapentaenoic acid; FA, fatty acids;

464 PANDiet, Diet Quality Index Based on the Probability of Adequate Nutrient Intake; PUFA, polyunsaturated

465 fatty acids; SFA, saturated fatty acids; sPNNS-GS2, simplified Programme National Nutrition Santé-Guidelines

466 Score 2

467 ¹Values are means (standard errors of the means) adjusted for energy intake using the residual method or

468 standard adjustment for contribution of macronutrient to energy intake, PANDiet, and sPNNS-GS2.

469 ²Values are crude means (SD)

470 ³Values are expressed as % of total energy intake

471 ⁴P-value for linear trend is estimated using linear contrast

472 ⁵Atherogenic fatty acids include lauric and myristic and palmitic acids

Table 41: Daily diet-related environmental indicators according to %UPF quintiles, (INCA 3, n=2,121)¹


Crude	Q1	Q2	Q3	Q4	Q5	P-trend²
Climate change (GHGe) (kg CO ₂ eq)	5.42 (5.15-.69)	5.81 (5.54-6.08)	5.96 (5.69-6.23)	6.19 (5.92-6.45)	6.24 (5.98-6.51)	0.003
Water use (m ³ world eq)	6.90 (6.62-7.18)	6.55 (6.27-6.83)	6.64 (6.36-6.92)	6.15 (5.87-6.43)	6.44 (6.16-6.72)	0.01
Land use (pt)	255.41 (240.34-270.48)	272.07 (257.07-287.08)	281.54 (266.55-296.53)	290.87 (275.82- 305.92)	299.58 (284.59-314.56)	0.01
Energy demand (MJ)	58.81 (56.94-60.67)	60.82 (58.96-62.68)	63.34 (61.49-65.20)	61.68 (59.82-63.55)	63.23 (61.37-65.08)	0.01
Acidification (mol H ⁺ eq)	0.066 (0.002)	0.069 (0.002)	0.071 (0.002)	0.073 (0.002)	0.075 (0.002)	0.01
Resource use, minerals and metals (kg Sb eq)	9.73 (9.38-10.07)	10.04 (9.70-10.38)	10.00 (9.66-10.34)	9.58 (9.24-9.92)	9.63 (9.29-9.97)	0.25
Eutrophication, freshwater (kg P eq)	1.02 (0.97-1.07)	1.03 (0.98-1.08)	1.01 (0.96-1.06)	0.98 (0.93-1.03)	1.02 (0.97-1.06)	0.25
Eutrophication, marine (kg N eq)	20.96 (19.99-21.93)	22.57 (21.61-23.54)	23.53 (22.57-24.50)	22.88 (21.91-23.85)	23.61 (22.65-24.58)	0.06
Eutrophication, terrestrial (mol N eq)	0.26 (0.25-0.28)	0.28 (0.26-0.29)	0.28 (0.27-0.30)	0.30 (0.28-0.31)	0.30 (0.29-0.31)	0.01
Photochemical ozone formation (kg NMVOC eq)	16.19 (15.40-16.97)	16.69 (15.91-17.47)	16.52 (15.74-17.30)	16.87 (16.09-17.66)	17.00 (16.23-17.78)	0.37
Ozone depletion (kg CFC-11eq)	0.53 (0.45-0.62)	0.56 (0.48-0.64)	0.51 (0.43-0.59)	0.66 (0.58-0.74)	0.51 (0.43-0.59)	0.80
Particulate matter (disease incidence)	0.48 (0.46-0.50)	0.51 (0.48-0.53)	0.52 (0.49-0.54)	0.53 (0.51-0.55)	0.54 (0.52-0.56)	0.01
Ionising radiation (kBq U235 eq)	1.27 (1.22-1.31)	1.34 (1.30-1.38)	1.44 (1.40-1.49)	1.42 (1.37-1.46)	1.47 (1.43-1.52)	<.0001
EF score ³	0.66 (0.63-0.69)	0.69 (0.67-.72)	0.71 (0.68-0.74)	0.71 (0.69-0.74)	0.73 (0.70-0.75)	0.02
Adjusted for dietary energy intake	Q1	Q2	Q3	Q4	Q5	P-trend²
Climate change (GHGe) (kg CO ₂ eq)	5.93 (5.71-6.15)	5.84 (5.62-6.05)	5.59 (5.37-5.80)	5.93 (5.71-6.14)	5.82 (5.60-6.03)	0.19
Water use (m ³ world eq)	7.34 (7.09-7.59)	6.58 (6.34-6.82)	6.32 (6.07-6.56)	5.92 (5.68-6.17)	6.07 (5.82-6.31)	<.0001
Land use (pt)	280.28 (267.28-293.28)	273.46 (260.63-286.28)	263.24 (250.36-276.12)	278.17 (265.27-291.07)	278.65 (265.76-291.54)	0.31
Energy demand (MJ)	63.38 (62.17-64.58)	61.07 (59.89-62.26)	59.98 (58.79-61.17)	59.35 (58.16-60.54)	59.38 (58.19-60.57)	<.0001
Acidification (mol H ⁺ eq)	0.072 (0.001)	0.070 (0.001)	0.067 (0.001)	0.070 (0.001)	0.070 (0.001)	0.11
Resource use, minerals and metals (kg Sb eq)	10.49 (10.25-10.74)	10.08 (9.84-10.33)	9.44 (9.19-9.68)	9.19 (8.94-9.43)	8.99 (8.74-9.23)	<.0001
Eutrophication, freshwater (kg P eq)	1.09 (1.05-1.13)	1.04 (0.99-1.08)	0.96 (0.92-1.00)	0.94 (0.90-0.99)	0.96 (0.91-1.00)	<.0001
Eutrophication, marine (kg N eq)	22.94 (22.19-3.69)	22.68 (21.94-3.42)	22.07 (21.33-2.82)	21.87 (21.12-22.61)	21.95 (21.20-22.69)	0.001
Eutrophication, terrestrial (mol N eq)	0.29 (0.27-0.30)	0.28 (0.27-0.29)	0.27 (0.26-0.28)	0.28 (0.27-0.29)	0.28 (0.27-0.29)	0.20
Photochemical ozone formation (kg NMVOC eq)	17.38 (16.69-8.07)	16.75 (16.07-7.44)	15.64 (14.96-16.33)	16.26 (15.57-6.95)	16.00 (15.31-6.69)	0.01
Ozone depletion (kg CFC-11eq)	0.58 (0.49-0.66)	0.56 (0.48-0.64)	0.48 (0.40-0.56)	0.64 (0.55-0.72)	0.47 (0.39-0.55)	0.39
Particulate matter (disease incidence)	0.52 (0.51-0.54)	0.51 (0.49-0.53)	0.49 (0.47-0.50)	0.51 (0.49-0.53)	0.50 (0.49-0.52)	0.06
Ionising radiation (kBq U235 eq)	1.38 (1.35-1.41)	1.35 (1.32-1.37)	1.36 (1.34-1.39)	1.36 (1.33-1.39)	1.38 (1.35-1.41)	0.53
EF score ³	0.72 (0.70-0.74)	0.69 (0.68-0.71)	0.67 (0.65-0.69)	0.68 (0.66-0.70)	0.68 (0.66-0.70)	0.002

473 Abbreviation: EF, ecological footprint; GHGe, greenhouse gas emissions. Units are as follows: kg CO₂ eq, carbon dioxide equivalent; m³ world eq, water use in cubic meters
474 of water; land use is estimated as loss of soil organic matter content in kilograms of carbon deficit (kg C deficit) dimensionless and expressed as Points (Pt); MJ, megajoule;
475 mol H⁺ eq, equivalent of moles hydron; kg Sb eq, equivalent of kilograms of antimony; kg P eq, equivalent of kilograms of phosphorus, kg N eq, equivalent of kilograms of
476 nitrogen; mol N eq, equivalent of moles of nitrogen; kg NMVOC eq, equivalent of kilograms of non-methane volatile organic compounds; kg CFC-11eq, equivalent of
477 kilograms of trichlorofluoromethane (Freon-11); Emission of particulate matter in change in mortality due to particulate matter emissions; kg U235 eq, equivalent of
478 kilobecquerels of Uranium 235
479 ¹Values are means (95% CI) (crude or energy-adjusted)

480 ²P-value for linear trend is estimated using contrast


481 ³For the EF score, the higher it is, the more impactful it is

Figure 1: Farm and post-farm stages contribution to daily diet-related environmental indicators according to %UPF quintiles (INCA 3, n=2,121), adjusted for energy intake^{1,2}



482
483 Panel A corresponds to the pressures, in absolute values, of each stage. Panel B corresponds to the pressures, in
484 relative value (%), of each stage
485 ¹Climate change (Greenhouse gas emissions), energy demand, land use and water use are expressed in kg
486 CO₂eq, m³ world eq, pt, MJ, respectively
487 ²Energy-adjusted values

Figure 2: Contribution of NOVA class consumption to daily diet-related environmental indicators according to %UPF quintiles (INCA 3, n=2,121), adjusted for energy




488

489 Panel A corresponds to the pressures, in absolute value, related to %NOVA class in the diet. Consumption,
490 climate change (Greenhouse gas emissions), energy demand, land use and water use are expressed in g/d, kg
491 CO₂eq, m³ world eq, pt, MJ, respectively. Panel B corresponds to same data but in relative value (%).

492

Figure 3: Daily diet-related environmental indicators for each step for substitution model of UPF (INCA 3, n=2,121)^{1,2}



493

494 ¹ Climate change (Greenhouse gas emissions), energy demand, land use and water use are initially expressed in
495 g/d, kg CO₂eq, m³ world eq, pt, MJ, respectively

496 ² Model is substitution of NOVA1 (unprocessed) by UPF adjusted for energy intake