

Tab. S1: Number of diseases that are significantly diagnosed first or last from each category. From a total of n such diseases (first or last), x are observed from each category among the m diseases of that category and $N - m$ diseases not from that category in the HUNT sub-PheNet of N diseases. A hypergeometric test with variables $x, m, N - m$ and n are used to test if the observed values are greater than expected, $P(X \geq x)$, or less than expected, $P(X \leq x)$, where the p -values are Bonferroni adjusted with 16 tests, where p -values less than 0.025 are marked with a star.

Category	Diseases diagnosed first					Diseases diagnosed last				
	x	m	$N - m$	$P(X \geq x)$	$P(X \leq x)$	x	m	$N - m$	$P(X \geq x)$	$P(X \leq x)$
circulatory system	18	54	305	0.0214*	1	20	54	305	0.0355	1
congenital anomalies	1	1	358	1	1	1	1	358	1	1
dermatologic	2	15	344	1	1	1	15	344	1	1
digestive	3	35	324	1	1	4	35	324	1	1
endocrine/metabolic	7	30	329	1	1	10	30	329	1	1
genitourinary	4	33	326	1	1	9	33	326	1	1
hematopoietic	1	21	338	1	1	2	21	338	1	1
infectious diseases	0	2	357	1	1	0	2	357	1	1
injuries & poisonings	0	3	356	1	1	1	3	356	1	1
mental disorders	3	22	337	1	1	3	22	337	1	1
musculoskeletal	9	28	331	0.5521	1	9	28	331	1	1
neoplasms	5	55	304	1	0.8688	3	55	304	1	0.0122*
neurological	3	15	344	1	1	2	15	344	1	1
respiratory	4	17	342	1	1	4	17	342	1	1
sense organs	2	17	342	1	1	1	17	342	1	1
symptoms	0	11	348	1	1	5	11	348	0.9061	1
n	62					75				
N	359					359				