

# Testing, Tracing, and Vaccination Targets for Containment of the US Monkeypox Outbreak: A Modeling Study

Running title: “Testing, Tracing, and Vaccinating for Monkeypox Containment”

Melanie H. Chitwood<sup>1,2</sup>, Jiye Kwon<sup>1,2</sup>, Alexandra Savinkina<sup>1,2</sup>, Jo Walker<sup>1,2</sup>, Alyssa Bilinski<sup>3,4</sup>, Gregg Gonsalves<sup>1,2</sup>

1. Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT
2. Public Health Modeling Unit, Yale School of Public Health, New Haven, CT
3. Department of Health Services, Policy, and Practice, Brown School of Public Health, Providence, RI
4. Department of Biostatistics, Brown School of Public Health, Providence, RI

Word Count: 686

**Abstract:** We estimate the levels of community testing, contact tracing, and vaccination required to reduce the effective reproduction number ( $R_t$ ) of Monkeypox Virus (MPXV) below 1 for high risk men who have sex with men. We found that the critical threshold to vaccinate depends on the basic reproduction number of MPXV and on other public health measures. This analysis provides a framework for quantitative targets toward MPXV containment.

## Background

As of July 31 2022, there have been over 5,000 confirmed cases of monkeypox across 48 U.S. states, predominantly among men who have sex with men (MSM)<sup>1</sup>. Reducing transmission of monkeypox virus (MPXV) is crucial to protect the health of the MSM population and reduce the risk of transmission in the general US population. Public health measures – testing, contact tracing, vaccination – can slow the spread of MPXV and officials have called for rapid scale-up of these efforts to support containment<sup>2</sup>. However, target levels of these measures have not been established.

## Objective

We estimate the levels of community testing, contact tracing, and vaccination required to reduce the effective reproduction number ( $R_t$ ) of MPXV below 1 for high risk MSM (HR-MSM).

## Methods and Findings

We adapted a deterministic branching model<sup>3</sup> to describe the transmission of MPXV in HR-MSM (Supplement). Infectious individuals are detected with a probability that varies based on community detection rate and whether their source case underwent contact tracing (Figure 1). We assume that community detection of a case reduces the number of secondary infections by 50%, identifying a case early through contact tracing reduces the number of secondary infections by 90%, and all detected cases have the same probability of having their contacts traced. We used Byrd optimization to solve for the level of vaccine coverage (assuming 100% efficacy) for HR-MSM that would reduce  $R_t$  below 1 at different estimates of the basic reproduction number ( $R_0$ ) in HR-MSM<sup>4</sup> and rates of contact tracing and community detection. Previous work indicates that  $R_0$  remains below 1 in the general population; to reflect uncertainty in HR-MSM, we vary  $R_0$  from 1.2-2.0.

We estimate there are 498,000 HR-MSM in the US (Supplement). We found that testing and contact tracing without vaccination can reduce  $R_t$  below 1 in this population if  $R_0 \leq 1.4$ , at least 40% of cases are detected through community testing, and at least 50% of case contacts are traced. In the absence of a robust public health response (10% community detection; no contact tracing), we estimate that the critical threshold to vaccinate ranges from 12% ( $R_0 = 1.2$ ) to 47% ( $R_0 = 2.0$ ). With a moderate response, (at least 20% community detection; at least 25% of cases contact traced) we estimate the critical threshold to vaccinate ranges from 5% ( $R_0 = 1.2$ ) to 43% ( $R_0 = 2$ ) of the HR-MSM population (Figure 2). In this scenario, 49,800 – 428,080 doses of the Jynneos vaccine would need to be administered to 24,900 – 214,040 HR-MSM.

The critical threshold to vaccinate was not sensitive to assumptions about the fraction of secondary cases detected through contact tracing (Figure S1), but was sensitive to assumptions about how much less infectious detected cases are relative to undetected infections. With 20% community detection, 25% of case contacts traced, and  $R_0 = 1.6$ , a 25% reduction in secondary infections suggests a 32% threshold, while a 75% reduction suggests a 25% threshold (Figure S2).

## Discussion

We project the impact of public health measures to reduce the spread of MPXV in the US. We found that the critical threshold to vaccinate depends on the basic reproduction number of MPXV and on other public health measures. Nevertheless, rapid distribution of vaccinations to at least one third of the HR-MSM (at least 329,000 doses) with increased testing and contact tracing, can support containment efforts in most modeled scenarios. As of July 29, the US government has shipped 336,710 doses, although the number of administered doses and the specific profile of recipients are unclear<sup>5</sup>.

This analysis has a number of limitations. We estimate vaccine needs based on the assumption that spread outside the HR-MSM population is negligible. If there is sustained transmission of MPXV in other populations at risk (e.g. low-risk MSM; in congregate settings), increased vaccination efforts may be required. We further assume that vaccines have 100% efficacy and are efficiently targeted to HR-MSM. Last, key parameters remain uncertain, although we illustrate sensitivity of our results to variations in these, and can adjust estimates as more information becomes available. As a result, we believe this analysis provides a useful framework for quantitative targets toward MPXV containment.

## References

- (1) CDC. 2022 U.S. Map & Case count [Internet]. Centers for Disease Control and Prevention. Centers for Disease Control and Prevention; 2022 [cited 2022Jul27]. Available from: <https://www.cdc.gov/poxvirus/monkeypox/response/2022/us-map.html>. Archived at: <https://archive.ph/ySIpM>.
- (2) Nirappil, F. San Francisco, New York state declare monkeypox emergencies as outbreak grows. The Washington Post. 2022 July 28. [cited 2022Jul30]. Available from: <https://www.washingtonpost.com/health/2022/07/28/monkeypox-emergency-san-francisco-new-york/>. Archived at: <https://archive.ph/n8ryd>.
- (3) Bilinski A, Mostashari F, Salomon JA. Modeling Contact Tracing Strategies for COVID-19 in the Context of Relaxed Physical Distancing Measures. JAMA Network Open. 2020 Aug 21;3(8):e2019217–e2019217.
- (4) WHO. Second meeting of the International Health Regulations (2005) (IHR) emergency committee regarding the multi-country outbreak of Monkeypox [Internet]. World Health Organization; 2022 [cited 2022Jul28]. Available from: [https://www.who.int/news/item/23-07-2022-second-meeting-of-the-international-health-regulations-\(2005\)-\(ihr\)-emergency-committee-regarding-the-multi-country-outbreak-of-monkeypox](https://www.who.int/news/item/23-07-2022-second-meeting-of-the-international-health-regulations-(2005)-(ihr)-emergency-committee-regarding-the-multi-country-outbreak-of-monkeypox). Archived at: <https://archive.ph/gPeC4>.
- (5) US Department of Health and Human Services, Administration for Strategic Preparedness and Response. JYNNEOS Monkeypox Vaccine Distribution by Jurisdiction. [cited 2022Jul29]. Available from: <https://aspr.hhs.gov/SNS/Pages/JYNNEOS-Distribution.aspx>. Archived at: <https://archive.ph/1OeH8>.

## Supplement


Table S1. Model parameter values

Table S2. Estimation of secondary infections.

Figure S1. Vaccination and contact tracing efforts required to reduce  $R_t$  below 1, by  $R_0$  and community detection rates.

Figure S2. Vaccination and contact tracing efforts required to reduce  $R_t$  below 1 by percentage of contacts traced.

**Figure 1. Model structure.**



## Figure 2. Frontier of Contact Tracing, Community Testing, and Vaccination Coverage to Reduce the Effective Reproduction Number Below 1.

The x-axis displays the fraction of cases traced and the y-axis vaccine coverage. The community detection rate is varied across facets. Each line shows the efficient frontier for a given value of  $R_0$ . Each point on the line can be interpreted to mean the fraction of contacts traced and the vaccination rate at which  $R_t$  remains below 1.

