

Supplemental information

Extended BNT162b2 booster dosing generates a broader neutralizing antibody response against SARS-CoV-2 variants of concern

Carl Graham^{1*}, Thomas Lechmere^{1*}, Aisha Rehman¹, Jeffrey Seow¹, Ashwini Kurshan¹, Isabella Huettner¹, Thomas J.A. Maguire¹, Jerry Tam¹, Daniel Cox¹, Christopher Ward¹, Mariusz Racz², Anele Waters², Christine Mant³, Michael H. Malim¹, Julie Fox¹, Katie J. Doores^{1#}

¹ Department of Infectious Diseases, School of Immunology & Microbial Sciences, King's College London, London, UK.

² Harrison Wing, Guys and St Thomas' NHS Trust, London, UK.

³ Infectious Diseases Biobank, Department of Infectious Diseases, School of Immunology and Microbial Sciences, King's College London, London, UK.

* These authors contributed equally


Corresponding author: katie.doores@kcl.ac.uk

Supplementary Figure S1: SARS-CoV-2 Spike-reactive memory B cells measured


before and after vaccination. A) Example FACS gating for pre-vaccination sample (visit 1) and from post-vaccination sample (visit 3) from SARS-CoV-2 naïve individual. **B)** Frequency of memory B cells in the total B cell population at visit 1 and visit 3 for matched donors.

a

VA005 visit 1



VA005 visit 3



b

