

Finding a Needle in the Haystack: Design and Implementation of a Digital Site-less Clinical Study of Serial Rapid Antigen Testing to Identify Asymptomatic SARS-CoV-2 Infection

Authors: Apurv Soni MD PhD^{1,2,3}, Carly Herbert BA¹, Caitlin Pretz MS¹, Pamela Stamegna MD¹, Andreas Filippaios MD¹, Qiming Shi MS^{1,3,4}, Thejas Suvarna BBA BS⁵, Emma Harman MPH⁵, Summer Schrader BA⁵, Chris Nowak BA⁵, Eric Schramm PharmD⁵, Vik Kheterpal MD⁵, Stephanie Behar BA¹, Seanan Tarrant BA¹, Julia Ferranto BS¹, Nathaniel Hafer PhD⁴, Matthew Robinson MD⁶, Chad Achenbach MD⁷, Rob Murphy MD⁷, Yuka Manabe MD⁶, Bruce Barton PhD², Laurel O'Connor MD⁸, Nisha Fahey DO ScM⁹, Elizabeth Orvek MS², Peter Lazar BS², Didem Ayturk MS², Steven Wong BS², Adrian Zai MD PhD MPH², Lisa Cashman MPH¹⁰, Lokinendi V Rao PhD¹⁰, Katherine Luzuriaga MD⁴, Stephenie Lemon PhD², Allison Blodgett PhD⁴, William Heetderks PhD¹¹, John Broach MD MPH MBA⁸, David McManus MD ScM^{1,3,12}

AS and CH contributed equally to this work

Affiliations:

¹Program in Digital Medicine, Department of Medicine, University of Massachusetts Chan Medical School, Worcester, MA, USA

²Department of Population and Quantitative Health Sciences, University of Massachusetts Chan Medical School, Worcester, MA, USA

³Division of Clinical Informatics, Department of Medicine, University of Massachusetts Chan Medical School, Worcester, MA, USA

⁴University of Massachusetts Center for Clinical and Translational Science, University of Massachusetts Chan Medical School, Worcester, MA, USA

⁵CareEvolution, LLC, Ann Arbor, MI, USA

⁶Division of Infectious Disease, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA

⁷Division of Infectious Disease, Department of Medicine, Havey Institute for Global Health, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

⁸Department of Emergency Medicine, University of Massachusetts Chan Medical School, Worcester, MA, USA

⁹Department of Pediatrics, University of Massachusetts Chan Medical School, Worcester, MA, USA

¹⁰Quest Diagnostics, Marlborough, MA, USA

¹¹National Institute of Biomedical Imaging and Bioengineering, NIH, via contract with Kelly Services, Bethesda, MD, USA

¹²Division of Cardiology, Department of Medicine, University of Massachusetts Chan Medical School, Worcester, MA, USA

Corresponding Author:

Apurv Soni, M.D., Ph.D.

38 Program in Digital Medicine, Department of Medicine
39 S6-757, Medical School Building
40 55 Lake Avenue N
41 Worcester, MA, 01605
42 Apurv.soni@umassmed.edu
43

Abstract:

Background: Over-the-counter rapid antigen tests for SARS-CoV-2 with an Emergency Use Authorization (EUA) in the United States generally include a condition of authorization to evaluate the test's performance in asymptomatic individuals when used serially. A goal of this study was to investigate the performance of SARS-CoV-2 antigen serial testing and generate data to support regulatory decisions.

Objective: To describe a novel study design to evaluate serial use of rapid antigen tests in detecting SARS-CoV-2 virus among asymptomatic individuals.

Design: Prospective cohort study using a decentralized approach. Eligible participants from across the U.S. could enroll and complete this study from their home environment through a study app. Participant enrollment was prioritized based on regional 7-day case rates, participants' vaccination status, and sociodemographic characteristics prior to enrollment. Prioritization criteria were adjusted on a daily or weekly basis. Enrolled participants were mailed rapid antigen tests and molecular comparator collection kits and asked to test every 48 hours for 15 days. Three companies' rapid antigen tests were used in the study; assignment of participant to a test was criteria-based and non-random, precluding head-to-head comparison between the tests.

Participants: Mainland United States residents over 2 years old with no reported COVID-19 symptoms in the 14 days prior to study enrollment.

Main Measures: Participant demographics, COVID-19 vaccination status, and geographic distribution were used to understand the impact of the site-less recruitment and enrollment strategy.

Key Results: A total of 7,361 participants enrolled in the study between October 18, 2021 and February 15, 2022. Throughout the study, 369 participants tested positive for SARS-CoV-2,

including 167 who were asymptomatic and tested negative on SARS-CoV-2 molecular assays to start the study. This exceeded the initial enrollment goals of 60 positive participants. We enrolled participants from 44 of the 48 mainland U.S. states, and geographic distribution of participants shifted in accordance with the changing COVID-19 prevalence nationwide.

Conclusions: The novel, digital site-less approach employed in the ‘Test Us At Home’ study enabled rapid, efficient, and rigorous evaluation of rapid diagnostics for COVID-19, and can be adapted across research disciplines to optimize study enrollment and accessibility.

INTRODUCTION

Over-the-counter (OTC) rapid antigen SARS-CoV-2 tests present unique opportunities for widespread COVID-19 testing. However, the authorization of these tests under the Food and Drug Administration's (FDA) Emergency Use Authorization (EUA) generally imposes a requirement on companies to evaluate their rapid antigen test's performance for detecting asymptomatic infections with serial screening. To accomplish this, it is necessary to conduct a longitudinal study to observe the onset of SARS-CoV-2 infection and performance of rapid antigen tests across the course of infection.⁵

The challenge of conducting this study is further compounded by fluctuations in the daily incidence of SARS-CoV-2 across different geographical regions. For instance, on September 1, 2021, Massachusetts reported 24.2 cases of SARS-CoV-2 per 100k residents; however, on that same date, the state of Georgia reported 85.9 cases per 100k residents.⁶ Further, asymptomatic positivity rate from screening studies has been found to be between 1.9% and 3%.⁷ A study to identify new-onset SARS-CoV-2 infection in this setting presents limited viable options, especially when considering the financial resources needed to set up multiple in-person recruitment centers to facilitate recruitment from different areas of the country where case-rate may be high at any given time. However, recruitment and enrollment methodologies often fall short, with 11% of sites in multi-center clinical trials failing to enroll any participants, 40% unable to meet recruitment goals, and nearly half of enrolled participants dropping out of trials prior to study completion.^{8,9} Additionally, despite efforts to increase diversity in clinical trials, clinical trial participants are still predominantly white and English-speaking, posing ethical and scientific issues.¹⁰⁻¹² Digital site-less studies pose great opportunities to enable community-based recruitment and retention across geographic regions, and have been facilitated by increased acceptance of digital medicine and mobile technologies in recent years.¹³

This manuscript describes the design, implementation, and completion of a novel, digital, site-less study called ‘Test Us at Home’ funded by the NIH Rapid Acceleration Diagnostics (RADx) Program. This study is a result of the collaboration between UMass Chan Medical School, the FDA, and the National Institutes of Health (NIH). In this manuscript, we outline the rationale for key decision-making and describe findings of the operational effort needed to successfully collect data under this novel paradigm. We describe the challenges faced by our study team and how we overcame those challenges to provide rapid, rigorous evaluation of OTC diagnostics across diverse participants of all ages in community-based settings.

METHODS

Study Design:

This prospective cohort study recruited participants over two-years old who were asymptomatic and self-reported absence of SARS-CoV-2 infection in the three months prior to enrollment. Participants were asked to test via rapid antigen test and molecular comparator every 48-hours for 15 days. Primary analysis was restricted to participants that started the study with a negative result on SARS-CoV-2 molecular assays.

Study Objectives:

The primary objective of this collaboration between the FDA, NIH, and the RADx Tech Clinical Studies Core at UMass Chan was to generate right-of-reference data that can be used towards FDA EUA. To accomplish this, we aimed to evaluate the performance of serial use of three different rapid antigen test to detect a new-onset SARS-CoV-2 infection, as determined by a laboratory-based molecular test. Secondary objectives of this study were to understand the different factors associated with detection of asymptomatic positive patients using OTC devices, including the optimal testing frequency and duration for individuals after exposure to COVID-19. Lastly, this study aimed to identify whether there are certain subpopulations that are more likely to test positive asymptotically.

Study Population and Recruitment: To approximate the performance of rapid antigen tests in the general public, the FDA suggested against preferentially enrolling participants with a known history of SARS-CoV-2 exposure (oral communication, August 3, 2021). Based on the FDA guidance, the goal of the study was to recruit at least 60 participants who started the study without any symptoms and with a negative SARS-CoV-2 molecular test and who subsequently tested positive during the study period. Due to the fluctuating rates of COVID-19 positivity and the importance of recruiting asymptomatic participants, we employed a decentralized study design whereby eligible participants from anywhere in the mainland United States except for Arizona could participate. Residents from the state of Arizona were excluded because Quest Diagnostics, the laboratory performing molecular tests for this study, does not provide direct-to-consumer tests in that state. Additional inclusion and exclusion criteria are listed in Table 1. Recruitment relied on the engagement of community stakeholders, including local public health officials and community organizers, to advertise the study and reach communities nationwide. Periodic analyses were conducted to inform enrollment goals based on observed precision for positive percent agreement.

Enrollment: Participants who expressed interest in the study received an email or flyer with instructions for downloading the study app onto their phone or device and performing eligibility screening. The study app was a custom interface designed through the MyDataHelps platform. To enroll in the study, participants answered a series of questions to determine their eligibility and whether they meet the study's criteria for enrollment at a given time. All eligible participants underwent automated prioritization based on a-priori criteria that were adjusted on a daily or weekly basis (Supplemental Figure 1, Supplemental Figure 2). We adjusted prioritization criteria to preferentially enroll populations based on their region's SARS-CoV-2 transmission rates, community-level vaccination, and participants sociodemographic characteristics. This step occurred prior to consenting, and data collected was discarded if the participant did not enroll in the study before the data collection period ended; these data were never accessible to the study

team unless the participant consented to take part in the study. Periodic analyses were conducted to inform enrollment goals based on observed precision for positive percent agreement. Eligible participants who did not match the preferred criteria for enrollment during that time received a note saying that they cannot enroll in the study at this time but would receive communication from the study team when/if they became eligible (Supplemental Figure 3). These participants were placed on a “waiting list”, which was reviewed with regard to changing priority criteria throughout the study.

If eligible for enrollment, participants 18 years of age or older were asked to read the e-consent form and provide their signature through the study app (Supplemental Figure 4). Participants younger than 18 years were asked to review the e-assent and e-consent form with the support of their parent(s) or guardian(s) and provide assent for participation, while the parent/guardian provided consent for participation in the study. To acknowledge participants for their time participating in study, participants were eligible to receive up to \$250 for timely completion of all sample collections (Supplemental Figure 5). This study was approved by the Institutional Review Board of the WIRB-Copernicus Group.

Test Distribution: On enrollment, participants were asked to provide their shipping information through the study app. Participants were assigned to one of three OTC tests (Quidel QuickVue At-Home OTC COVID-19 Test, Abbott BinaxNOW COVID-19 Antigen Self Test, or BD Veritor™ At-Home COVID-19 Test) using an automated algorithm that was informed by the investigators’ discretion based on enrollment numbers and geographic location of the participants to allow for pragmatic distribution of the tests and fulfillment of the sample size (Supplemental Figure 6). The three readily available rapid antigen tests used in the study are all authorized for emergency use by the FDA and were selected for evaluation to provide multiple assessments and facilitate generalizing results to non-participating companies. These tests will be hereby referred to as test brands A, B, and C (randomly assigned), as the purpose of the study was not to directly compare different test performances. Participants residing together and enrolling with

the same address or enrolled in a group, such as a cohort of classmates, were assigned the same OTC test, to avoid mixing of test types between participants. Groups were given unique codes to input on enrollment (i.e. group codes), to ensure members were easily identifiable. As a result of this strategy, the assignment of tests to participants were non-random, and we intentionally did not pursue a strategy to compare performance between different tests because that was outside the scope of this project.

Because molecular assays require a prescription by a licensed physician, the study team contracted with PWNHealth, a national clinician network, and developed an automated order filing system working with Quest Diagnostics to place the requisite orders (Supplemental Figure 7). A total of 10 OTC tests and 7 home-collection kits supporting testing with the molecular comparators were provided to each participant. Participants received OTC tests and PCR home-collection kits by mail through separate shipments, unless participants enrolled at an approved in-person location, in which case participants were handed both types of testing kits directly. Once the test kits were received, participants were asked to confirm receipt through the study app prior to starting the testing period. If a participant needed additional OTC tests or PCR home-collection kits during the study period (e.g. tests were damaged upon delivery or lost in the mail), the study team provided additional tests.

Testing Schedule: Participants were asked to perform the OTC test and collect the specimen for molecular comparator testing on the same day roughly every 48 hours during the 15-day testing period (Table 2, Supplemental Figure 8). Immediately prior to testing, participants were asked to record any symptoms on the day of testing. The study app provided a push-notification testing reminder at the 44-hour mark and additional notifications every two hours until the 52-hour mark if the participant had not completed their tests. Participants were instructed via the app to have at least a 15-minute break between the OTC test and the sample collection for molecular tests. Participants were asked to only test with the OTC test on day 15 (Table 2). Participants were asked to record the results of the OTC tests when available by selecting one of the following

options: “Negative”, “Positive”, “Invalid”, or “Don’t Know”, and upload a photo of the test strip through the app (Supplemental Figure 9). Participants assigned to Test C were instead asked to download the Test C company-specific app which contained a test reader. The company test reader asked participants to upload a photo of the test strip and provided rapid antigen test results in real time. Test C users were asked to report the test result given by the test reader, rather than self-interpretation, to the study app. Test C participants were also asked to upload a picture in the study app for validation. If the participant tested positive by OTC test, they were referred to the OTC device’s instructions and told to contact their healthcare provider for any medical questions. In the event of an invalid OTC test result, the participant was asked to perform a second OTC test. If the second OTC result was also invalid, the participant was not asked to perform additional OTC tests on that day and was instructed to continue to the PCR sample collection (at least 15 minutes after OTC sample collection). Participants were responsible for shipping the PCR collection kit containing the sample using the pre-paid FedEx envelope based on the instructions provided by Quest Diagnostics, which were authorized by the FDA for emergency use.

Molecular Testing Procedures: Due to concerns of potential false-positive molecular tests, the FDA recommended the use of two types of molecular assays, and potentially, a third assay, if the prior two assays were discordant (oral communication, August 3, 2021).^{23,4} Once participant samples were received at Quest Diagnostics, the sample was divided into aliquots to perform Roche Cobas 6800 SARS-CoV-2 PCR and Quest Laboratory Developed RT-PCR (LDT) for all participants (Supplemental Figure 10). An additional aliquot was preserved by the Quest laboratory to allow for testing of the sample on Hologic Aptima Transcription Mediated Amplification (TMA) assay, and the UMass Chan team submitted an additional order for the Hologic Aptima TMA assay when the results of the Roche Cobas and Quest LDT were discordant or if at least one of the PCR assays resulted as inconclusive. All three molecular assays used are highly sensitive, authorized by EUA, and run per instructions per use. Remnant

samples of sample fluid from home-collection kits were frozen at -80 Celsius and shipped to UMass Chan Medical School for future testing. Results of the molecular PCR assays were available to participants through the Quest results portal and through communication from the study team. All participants that tested positive by molecular test received a phone call from a study team member. For determining performance statistics, a molecular test was considered as positive only if a majority of the three molecular assays were positive. In the event of discordant results, the participant was contacted by the study staff to explain the results and the results were made available through the study app.

Communication with the study team: Participants were provided a study hotline staffed with research coordinators to provide support during extended hours during the weekdays from 8 AM to 9 PM EST. Participants were also able to contact the study team via email outside of the hotline hours. All calls between coordinators and participants were documented in a call log containing the length of the call, the reason for the call, narrative about the call, and whether the issue was resolved.

Data Management: There were three primary sources of data (Supplemental Figure 11). All participant-reported data were collected through the MyDataHelps smartphone app and downloaded incrementally through secure file transfer protocol to UMass Chan servers. The molecular testing data were shared by Quest Diagnostics through an established datafeed between Quest and UMass Chan servers, which populated output of molecular assays. However, the PCR cycle threshold (CT) values and results of the tiebreaker assays were not part of routinely abstracted data and required manual abstraction. Finally, tracking reports were used to document all communications between participants and the study staff. All three sources of data were combined using a participant's unique identifier assigned by the MyDataHelps app. Paired OTC and PCR data were merged by matching on participant identifier and date of testing. When test dates were not aligned or there were unmatched results, a member of the study team reviewed all testing data from the participants to adjudicate paired

findings. The most common reason for mismatched dates was due to missing dates of collection or transcription errors, as date of collection for PCR sample is handwritten by the participant on the Quest requisition form. Study staff reconciled possible mismatched test results by reviewing data for requisition numbers entered in the app, shipment tracking data, and contact reports from the study. If mismatched data could not be adequately aligned, those data were considered as ineligible due to failed quality-checks. Additional adjudication undertaken by the study team prior to data lock included a manual review of all rapid antigen test images for participants who either had a self-reported positive, don't know, or invalid OTC result or if they tested positive on molecular test at any point during the study. All data were de-identified and shared with the FDA throughout the study. At the end of the study, the final and cleaned dataset was shared with the FDA and company-specific data were shared with the companies. All data will be made available on the NIH Data Hub once the main findings from this study are published.

RESULTS

Enrollment: In total, 7,361 participants enrolled in the study between October 18, 2021 and February 15, 2022 (Figure 1a). Due to complications with the Test C company app, Test C enrollment began on November 4, 2021, three weeks after Test A and B enrollment started (Figure 1b). Following initial surges in recruitment, the enrollment waitlist was implemented on November 24, 2021 to allow for refinement of enrollment from geographic hotspots (Figure 2). In total, 369 participants tested positive on molecular assays during the study and 167 of those were eligible for analyses (Figure 3).

Geographic and Sociodemographic Participant Characteristics: The geographic and sociodemographic characteristics of study participants changed over the course of the study, reflecting the shifts in the enrollment approach. In total, participants from 44 of the 48 mainland U.S. states enrolled in the study (Figure 2). In October, the majority of initial study recruitment occurred on a college campus in Wisconsin, resulting in a young, predominantly white, student

population (Figure 2; Table 3). Additionally, only 3.1% of enrolled participants in October were unvaccinated for SARS-CoV-2. In November, study recruitment was expanded throughout the United States, with enrollment from more than 20 different states. Participants from rural populations comprised 14% of the participants recruited in November. In December and January, study enrollment shifted to prioritize unvaccinated participants, preferentially pulling these participants from the study waitlist, resulting in 24.6% and 21.4% of monthly enrollment being unvaccinated (Table 3). In January, recruitment efforts also focused on increasing racial and ethnic diversity in the study population, and the minority of participants enrolled during January identified as White (42.6%). Additionally, 10.5% of participants enrolled in January were of Hispanic origin.

Operational Support and Logistics: Throughout the study, there were 11,646 contacts between coordinators and 4,389 distinct participants (Table 4). This accounted for 33,356 minutes of coordination time. Coordinators and participants communicated by phone call (39.7%), email (30.6%), voicemail (29.7%), and push notification within the study app (4.2%). The most common reason for contact was to share COVID-19 test results from molecular tests, which accounted for 24.3% of all calls. Other common reasons for contact included testing reminders (19.1%), shipping and receipt of test kits (16.7%), compensation (16.3%), and clarifying the testing schedule (10.4%). The company-generated app for Test C was initially only compatible with certain smartphones, and many participants were unable to use the company app effectively. Due to these challenges, we delayed enrollment for Test C by two weeks and troubleshooted the app with the company, to ensure participants were able to use the app appropriately (Figure 1a). With participants assigned to Tests A and B, there was not a company app, so similar issues did not arise. Further, the decentralized design necessitated an increased reliance on shipping vendors, allowing us to reach participants across the mainland US states. However, this also required a variety of tracking methods to keep track of the 58,888 kits shipped. This study spanned the holiday season, which resulted in additional shipping

delays; 1.2% (n =728) of packages were affected by logistical or participant errors that led to an unviable or missing RT-PCR test result.

DISCUSSION

Using an innovative digital, site-less study approach, we developed recruitment and enrollment strategies to capture new onset COVID-19 infections most effectively throughout the country, resulting in the detection of nearly three times as many new-onset COVID-19 cases as originally specified. The digital site-less approach allowed us to seamlessly and dynamically change enrollment patterns and sample different populations based on the evolving nature of the pandemic, to assess rapid diagnostics with agility, efficiency, and rigor among asymptomatic individuals.

The digital site-less approach offered great agility to alter the recruitment strategy to fit our needs throughout the evolution of the study. Initially, to iron out the study logistics, including shipping rapid antigen tests and molecular comparator collection kits directly to participants, we opened study recruitment in October 2021 to members of a Midwestern university marching band who were traveling together, where we anticipated a high amount of unmasked, close contact interactions. This group enrollment is reflected in the demographics of participants in October, who were younger and had a higher proportion of white participants than participants recruited in November through January. Later, we were able to geographically alter the sampling to enroll participants from throughout the United States and use a waitlist to selectively sample certain populations by zip-code, determined by community prevalence of COVID-19, and demographics to ensure we were curating a representative sample. For example, after seeing low enrollment of participants over age 65 years in October through December, in January, we selectively pulled those over 65 years of age off the waitlist, increasing the proportion of participants in this group from 1.5% in October to 13.4% in January. Additionally,

the recruitment strategy resulted in 16.6% of all enrolled participants being unvaccinated, matching nationwide estimates for those over 5 years of age (CDC data tracker). The waitlist approach also allows us to ensure that enrollment was not solely clustered among those who have greatest access to recruitment strategies, including news articles and social media, but rather gave the time for information to disseminate to and within various communities.

The remote design also resulted in gains in study efficiency, in terms of both coordinator time and cost. Instead of requiring coordinators to schedule and supervise serial sample collections for each participant, the virtual, site-less design allowed participants to test in their homes and on their time, increasing accessibility to individuals with various employment and school schedules, as well as living situations. According to the study coordination logs, direct participant-coordinator phone calls averaged to just over 1-hour per day (82.3 minutes over 161 days of the study) and consisted primarily of testing reminders and returning molecular test results to participants, as well as notifying positive participants of their infection. As clinical research coordinators on average support upwards of seven studies concurrently and approximately two-thirds of coordinators report being expected to work more time on studies than allotted, it is important to understand methods to optimize study coordination time, while maintaining quality and consistency.¹⁴ Here, we showed the ability for site-less digital trials to run with minimal participant-coordinator interaction, while maintaining high enrollment and adherence to the study protocol (data not shown).

Site-less cohort studies have become increasingly common, as advances in technology have made virtual recruitment and engagement more feasible.¹⁵ However, digital products and solutions in clinical research remain underutilized.¹⁶ Site-less studies can not only facilitate recruitment and participant engagement, but they can also enhance scientific rigor and design. Through coordination with shipping services, home-test distributors, and clinical labs, we organized shipments of study supplies to participants on enrollment, to allow participants to start testing at home within 24 to 48 hours of enrollment with minimal participant burden. Further, we

coordinated the pick-up and shipment of repeated biological samples to the central laboratory throughout the study period for comparator testing, which has seldom been done within the framework of a digital study. The success of these workflows opens the door for the adaptation of site-less studies to answer many complex research questions, by integrating survey responses, biological samples, and home health monitoring.

There are a few potential limitations to the study. We did not require participants to be under video observation while performing the rapid antigen tests or collecting samples for the molecular tests, which could have resulted in improper technique and sample collection. However, our study facilitated use of these tests ‘as intended’ and ‘as authorized’ by the FDA and therefore may represent better approximation of real-world performance of the tests than traditional clinical studies, which simulate a home environment in controlled research sites. Additionally, the smartphone app required participants to take images of the rapid antigen tests; therefore, we could verify the rapid antigen test results for all participants. Due to the scale and nature of this study, we relied on commercial vendors for assembly and distribution of the testing kits, which precluded our ability to provide study-specific written instructions enclosed within the kit; we instead relied on providing all study-specific instructions through the smartphone app. This resulted in confusion for some participants who did not start testing right away after receiving the test kits or did not start with performing both rapid antigen and PCR tests concurrently. However, our study team were able to detect these instances and intervene by contacting the participants who did not begin testing for an extended period of time after receiving the tests or were not performing paired testing. Because of the logistical constraints, we could not facilitate a cold-chain transportation of the collected specimen, which preclude our ability to perform viral culture studies, which would have provided additional insights into the performance of rapid antigen tests. The workflow for sample collection for molecular tests required participants to handwrite the date and time of sample collection on a form, and the data generated from the laboratory had to be linked with rapid antigen testing data based on

Participant ID and day of testing, which resulted in mismatches and required substantial manual effort for reconciling differences by reviewing tracking history of sample collection. Finally, the requirement of smartphone ownership to participate in the study disadvantages people without smartphones. However, smartphone use is ubiquitous and has been accelerated by the SARS-CoV-2 pandemic.¹⁷ Additionally, we submit that the barrier to participation created by requiring a smartphone is lesser than that created by requiring participants to take time off from their day during business hours to travel to a clinical study site, multiple times during the study.

These limitations notwithstanding, this study represents the most robust attempt at understanding the performance of serial use of rapid antigen tests for SARS-CoV-2 detection among asymptomatic participants. We summarize the advantages of our digital siteless approach in Table 5 and outline recommendations for future digital studies. The collaboration behind this study between the NIH, FDA, and the RADx Clinical Studies Core allowed for development of a protocol that was innovative and a clinical study that will provide tremendous value to federal agencies, academic researchers and companies for elucidating key questions related to rapid antigen tests' performance. Future collaborative efforts to develop best practice guidelines and infrastructure for performing digital clinical studies are needed to advance scientific community's ability to perform rigorous clinical research and answer vexing questions.

422

423 **Acknowledgment:** We are grateful to our study participants and to our collaborators from the
 424 National Institute of Health (NIBIB and NHLBI) who provided scientific input into the design of
 425 this study and interpretation of our results, but could not formally join as co-authors due to
 426 institutional policies and to the Food and Drug Administration (Office of In Vitro Diagnostics and
 427 Radiological Health) for their involvement in the primary TUAH study. We received meaningful
 428 contributions from Drs. Bruce Tromberg, Jill Heemskerk, Dennis Buxton, Erin Iturriaga, Jue
 429 Chen, Andrew Weitz, and Krishna Juluru. We are thankful to the developers of the rapid antigen
 430 tests, who donated their tests for research use in this study. We are thankful to county health
 431 departments across the country who helped with recruitment for this siteless study by spreading
 432 the word in their networks.

433

434 **Competing Interest Statement**

435 VK is principal, and TS, SS, CN, ES, and EH are employees of the health care technology
 436 company CareEvolution, which was contracted to configure the smartphone study app, provide
 437 operational and logistical support, and collaborate on overall research approach. LS and LR are
 438 employees of Quest Diagnostics LLC, which was contracted to provide direct-to-consumer kits,
 439 logistical support for nationwide RT-PCR testing, and operational support for producing
 440 molecular testing results. DDM reports consulting and research grants from Bristol-Myers
 441 Squibb and Pfizer, consulting and research support from Fitbit, consulting, and research support
 442 from Flexcon, research grant from Boehringer Ingelheim, consulting from Avania, non-financial
 443 research support from Apple Computer, consulting/other support from Heart Rhythm Society.
 444 YCM has received tests from Quanterix, Becton-Dickinson, Ceres, and Hologic for research-
 445 related purposes, consults for Abbott on subjects unrelated to SARS-CoV-2, and receives
 446 funding support to Johns Hopkins University from miDiagnostics. AS receives non-financial
 447 support from CareEvolution for collaborative research activities.

448

449 **Funding Statement**

450 This study was funded by the NIH RADx Tech program under 3U54HL143541-02S2 and NIH
 451 CTSA grant UL1TR001453. The views expressed in this manuscript are those of the authors
 452 and do not necessarily represent the views of the National Institute of Biomedical Imaging and
 453 Bioengineering; the National Heart, Lung, and Blood Institute; the National Institutes of Health,
 454 or the U.S. Department of Health and Human Services. Salary support from the National
 455 Institutes of Health U54HL143541, R01HL141434, R01HL137794, R61HL158541,
 456 R01HL137734, U01HL146382 (AS, DDM), U54EB007958-13 (YCM, MLR), A1272201400007C,
 457 UM1AI068613 (YCM).

458

References:

1. Policy for Coronavirus Disease-2019 Tests During the Public Health Emergency (Revised) | FDA. Accessed July 4, 2022. <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/policy-coronavirus-disease-2019-tests-during-public-health-emergency-revised>
2. Kanji JN, Zelyas N, MacDonald C, et al. False negative rate of COVID-19 PCR testing: a discordant testing analysis. *Virology Journal*. 2021;18(1):13. doi:10.1186/s12985-021-01489-0
3. Wei Z, Ryhana M, Elizabeth S, J BG, J MA. Comparison of Four Molecular In Vitro Diagnostic Assays for the Detection of SARS-CoV-2 in Nasopharyngeal Specimens. *Journal of Clinical Microbiology*. 2022;58(8):e00743-20. doi:10.1128/JCM.00743-20
4. Lowe CF, Matic N, Ritchie G, et al. Detection of low levels of SARS-CoV-2 RNA from nasopharyngeal swabs using three commercial molecular assays. *Journal of Clinical Virology*. 2020;128:104387. doi:<https://doi.org/10.1016/j.jcv.2020.104387>
5. Smith RL, Gibson LL, Martinez PP, et al. Longitudinal Assessment of Diagnostic Test Performance Over the Course of Acute SARS-CoV-2 Infection. *The Journal of Infectious Diseases*. 2021;224(6):976-982. doi:10.1093/infdis/jiab337
6. CDC COVID Data Tracker. Accessed May 12, 2021. <https://covid.cdc.gov/covid-data-tracker/#variant-proportions>
7. Herbert C, Kheterpal V, Suvarna T, et al. Design and Preliminary Findings of Adherence to the Self-Testing for Our Protection From COVID-19 (STOP COVID-19) Risk-Based Testing Protocol: Prospective Digital Study. *JMIR Form Res*. 2022;6(6):e38113. doi:10.2196/38113
8. Lamberti MJ, Getz K. Profiles of New Approaches to Improving the Efficiency and Performance of Pharmaceutical Drug Development. *Tufts White Paper*. Published online 2015.
9. Hirsch IB, Martinez J, Dorsey ER, et al. Incorporating Site-less Clinical Trials Into Drug Development: A Framework for Action. *Clinical Therapeutics*. 2017;39(5). doi:10.1016/j.clinthera.2017.03.018
10. Gray DM, Nolan TS, Gregory J, Joseph JJ. Diversity in clinical trials: an opportunity and imperative for community engagement. *The Lancet Gastroenterology and Hepatology*. 2021;6(8). doi:10.1016/S2468-1253(21)00228-4
11. Clark LT, Watkins L, Piña IL, et al. Increasing Diversity in Clinical Trials: Overcoming Critical Barriers. *Current Problems in Cardiology*. 2019;44(5). doi:10.1016/j.cpcardiol.2018.11.002
12. National Academies of Sciences Engineering and Medicine. *Improving Representation in Clinical Trials and Research: Building Research Equity for Women and Underrepresented Groups*. The National Academies Press; 2022. doi:<https://doi.org/10.17226/26479>.
13. Inan OT, Tenaerts P, Prindiville SA, et al. Digitizing clinical trials. *npj Digital Medicine*. 2020;3(1):101. doi:10.1038/s41746-020-0302-y
14. Speicher LA, Fromell G, Avery S, et al. The critical need for academic health centers to assess the training, support, and career development requirements of clinical research coordinators:

496 recommendations from the Clinical and Translational Science Award Research Coordinator
 497 Taskforce. *Clin Transl Sci*. 2012;5(6):470-475. doi:10.1111/j.1752-8062.2012.00423.x

498 15. Banks MA. Core Concept: In the wake of COVID-19, decentralized clinical trials move to center
 499 stage. *Proc Natl Acad Sci U S A*. 2021;118(47):e2119097118. doi:10.1073/pnas.2119097118

500 16. Marra C, Gordon WJ, Stern AD. Use of connected digital products in clinical research following
 501 the COVID-19 pandemic: a comprehensive analysis of clinical trials. *BMJ Open*.
 502 2021;11(6):e047341. doi:10.1136/bmjopen-2020-047341

503 17. Demographics of Mobile Device Ownership and Adoption in the United States | Pew Research
 504 Center. Accessed July 31, 2022. <https://www.pewresearch.org/internet/fact-sheet/mobile/>

505

506

Tables and Figures:

Table 1: Inclusion and Exclusion Criteria

Inclusion Criteria	
➤	At least 2 years old
➤	Access or parental access to a smartphone or device to download and use study app
➤	Reside in mainland United States
➤	Able to receive mail at home
➤	Willing to drop off comparator samples at FedEx drop-off site or allow for pick-up service to retrieve samples
➤	Report no COVID-19 symptoms within 14 days prior to study enrollment
Exclusion Criteria	
➤	Self-reported positive test for SARS-CoV-2 infection in the previous 3 months
➤	Facial trauma interfering with nasal swabs
➤	Lack capacity to consent
➤	Do not understand English or Spanish
➤	Currently in the correctional justice system
➤	No internet access on smartphone while at home

Table 2: Test Us At Home Testing Schedule

Study Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
OTC Device	X		X		X		X		X		X		X		X
Molecular Comparator	X		X		X		X		X		X		X		

X=Test completed on this day

512

513 **Table 3: Test Us at Home Participant Demographics by Month of Enrollment**

	October N=415	November N=3701	December N=1592	January N=1653	Total N=7361
Test Assigned: n (%)					
Test A	80 (19.3)	1,474 (39.8)	375 (23.6)	530 (32.1)	2,459 (33.4)
Test B	335 (80.7)	1,231 (33.3)	375 (23.6)	565 (34.2)	2,506 (34.0)
Test C	0 (0.0)	996 (26.9)	842 (52.9)	558 (33.8)	2,396 (32.6)
Vaccination Doses: n (%)					
1	51 (12.3)	1,176 (31.8)	418 (26.3)	32 (1.9)	1,677 (22.8)
2	350 (84.3)	2,062 (55.7)	513 (32.2)	352 (21.3)	3,277 (44.5)
3+	0 (0.0)	0 (0.0)	269 (16.9)	915 (55.4)	1,184 (16.1)
Unvaccinated	13 (3.1)	462 (12.5)	392 (24.6)	354 (21.4)	1,221 (16.6)
Don't know	1 (0.2)	1 (0.1)	0 (0.0)	0 (0.0)	2 (0.1)
Age Category: n (%)					
Less than 18 years	0 (0.0)	417 (11.3)	199 (12.5)	164 (9.9)	780 (10.6)
18-45 years	384 (92.5)	2,156 (58.3)	851 (53.5)	878 (53.1)	4,269 (58.0)
45-65 years	25 (6.0)	871 (23.5)	392 (24.6)	390 (23.6)	1,678 (22.8)
Over 65 years	6 (1.5)	257 (6.9)	150 (9.4)	221 (13.4)	634 (8.6)
Gender: n (%)					
Female	232 (55.9)	2,340 (63.2)	1,045 (65.6)	1,025 (62.1)	4,642 (63.1)
Male	131 (31.6)	1,208 (32.6)	458 (28.8)	523 (31.7)	2,320 (31.5)
Nonbinary	28 (6.8)	49 (1.3)	20 (1.3)	19 (1.2)	116 (1.6)
Transgender	14 (3.4)	20 (0.5)	11 (0.7)	2 (0.1)	47 (0.6)
No answer	10 (2.4)	84 (2.3)	58 (3.6)	84 (5.1)	236 (3.2)
Race/Ethnicity: n (%)					
White	358 (86.3)	3,167 (85.6)	1,249 (78.4)	703 (42.6)	5,477 (74.4)
African American/Black	15 (3.6)	95 (2.6)	94 (5.9)	159 (9.6)	363 (4.9)
Asian	5 (1.2)	98 (2.7)	50 (3.1)	302 (18.3)	455 (6.2)
Pacific Islander	0 (0.0)	1 (0.1)	1 (0.1)	25 (1.5)	27 (0.4)
Native American/Alaskan Native	3 (0.7)	19 (0.5)	9 (0.6)	10 (0.6)	41 (0.6)
Other	9 (2.2)	83 (2.2)	49 (3.1)	192 (11.6)	333 (4.5)
Multiracial	13 (3.1)	137 (3.7)	66 (4.1)	129 (7.8)	345 (4.7)

No answer	12 (2.9)	101 (2.7)	74 (4.7)	133 (8.1)	320 (4.4)
Hispanic: n (%)	24 (5.8)	243 (6.6)	88 (5.5)	174 (10.5)	529 (7.2)
Geographic Setting					
Rural	25 (6.0)	521 (14.1)	188 (11.8)	137 (8.3)	871 (11.8)
Suburban	238 (57.4)	1,997 (54.0)	900 (56.5)	797 (48.2)	3,932 (53.4)
Urban	144 (34.7)	1,126 (30.4)	456 (28.6)	643 (38.9)	2,369 (32.2)
No answer	8 (1.9)	57 (1.5)	48 (3.0)	76 (4.6)	189 (2.6)
Highest Level of Education: n (%)					
Did not finish high school	0 (0.0)	331 (8.9)	160 (10.1)	131 (7.9)	622 (8.5)
High school graduate	173 (41.7)	328 (8.9)	145 (9.1)	179 (10.8)	825 (11.2)
Some college	155 (37.4)	830 (22.4)	336 (21.1)	378 (22.9)	1,699 (23.1)
Bachelor's Degree or higher	78 (18.8)	2,027 (54.8)	833 (52.3)	830 (50.2)	3,768 (51.2)
Don't know	1 (0.2)	45 (1.2)	18 (1.1)	19 (1.2)	83 (1.1)
No answer	8 (1.9)	140 (3.8)	100 (6.3)	116 (7.0)	364 (4.9)
Employment Status: n (%)					
Unemployed	22 (5.3)	414 (11.2)	134 (8.4)	169 (10.2)	739 (10.0)
Retired	8 (1.9)	253 (6.8)	121 (7.6)	170 (10.3)	552 (7.5)
Student	265 (63.9)	656 (17.7)	273 (17.2)	239 (14.5)	1,433 (19.5)
Working now	106 (25.5)	2,230 (60.3)	967 (60.7)	934 (56.5)	4,237 (57.6)
Other	3 (0.7)	55 (1.5)	27 (1.7)	30 (1.8)	115 (1.6)
No answer	11 (2.7)	93 (2.5)	70 (4.4)	111 (6.7)	285 (3.9)

514

515

Table 4. Type of Contact and Reasons for Participant-Coordinator Contact

	Number of Participant-Coordinator Contacts
Total	11,646
Type of Contact, n (%)	
Phone Call	4,627 (39.7)
Voice mail	3,455 (29.7)
Email	3,564 (30.6)
Unscheduled Push notification	483 (4.2)
Who Initiated Contact?, n (%)	
Participant	5,371 (46.1)
Coordinator	5,261 (45.2)
Both	825 (7.1)
Missing	189 (1.6)
Reason, n^a	
App issues	1,123 (9.6)
Interpretation of Test Results	16 (0.1)
Symptom reporting	63 (0.5)
Issues with Rapid Antigen Tests	251 (2.2)
Testing Reminder	2,228 (19.1)
Compensation	1,903 (16.3)
Shipping of Tests	1,950 (16.7)
Study testing schedule	1,215 (10.4)
Giving Test Results	2,833 (24.3)
Other	1,794 (15.4)

^aReasons for contact were not mutually exclusive

Table 5. Summary of features of this study and opportunity for improvement

Challenges	Innovation	Opportunity for improvement
Widespread outreach to asymptomatic individuals	Democratized participation such that anyone, anywhere, anytime can choose to participate through a smartphone without needing to travel to a clinical study site or recruitment centers	Requirement of smartphone to enroll can prevent some people from participating. Approaches that loan study smartphones or leverage community-health workers as digital studies' liaisons can help overcome this limitation
Engagement of diverse communities	A dedicated team met with county health workers and community stakeholders from across the country to promote awareness of the study and participation among people of color, indigenous heritage, and rural locations	A network of community partners and stakeholders for digital clinical studies that inform, plan, and conduct outreach efforts can help further enhance participation of historically underrepresented populations
Recruitment from SARS-CoV-2 "hotspot"	Use of CDC daily case reports and projects to identify geographical regions where community outreach efforts were enhanced and which were identified as 'preferred regions for enrollment'	Promoting peer-to-peer recruitment from hotspots through the use of the smartphone apps
Refinement of enrollment	Adding a layer of prioritization to allow cohort composition such that participants from "hotspots" and those that improve diverse representation of the enrolled cohort	Automatically adaptive priority criteria that does not require manual input or decision-making but can allow manual override
Consenting participants including children > 2 years old	Self-contained description of consent, including brief description of key parts of the study displayed through the smartphone app	Use of participatory videos that explain the study objectives and activities embedded within the smartphone app
Assigning rapid antigen tests	Pre-specified criteria that assigned tests to participants automatically	Adaptive algorithm that refines criteria to create balanced cohorts based on multiple variables
Distributing testing kits	Asynchronous assembly and shipment of kits from commercial vendors	Synchronous assemble and shipment of kits such that all materials arrive in box instead of different boxes
Collecting samples for molecular tests	Direct-to-consumer kits for nasal swabs that were returned using FedEx overnight prepaid envelopes	Courier pickup options that can be invoked from within the study app

Performing up to three separate molecular assays in CLIA laboratories	Automated ordering of two tests and manual ordering of additional tests when discordance was observed	Automated ordering of discordant testing with streamlined data collection
Serial testing within a specific window (44-52 hours apart)	Automated app notification and text messages every two hours during the testing window	Providing the option of defining notification frequency during the testing window to the participants
Verification of rapid antigen testing results	In-app widget to allow participants to take an image of rapid antigen tests that were later verified by the participants	Automated readers that confirms that an image of rapid antigen test was collected and interprets the test result ^a
Data collection and quality	Data fidelity and timestamp for each element collected within the study app and collation of disparate datafeed from the molecular testing laboratory	Minimizing user entry of free-text or handwritten data
Participant reimbursement	Batch processing of prepaid Bank of America cards that are mailed to participants' home	Automated delivery of reimbursement through smartphone app once the requirements are met
a: future automated readers that help with interpretation of test result will need FDA authorization		

Figure 1: Test Us At Home Study Enrollment

Figure 1a:

Test Us At Home Enrollment: October 18, 2021-January 22, 2022





Figure 1b:


Test Us At Home Enrollment by Test



a: Test A and B enrollment opens to individuals with specific group codes on October 18, 2021; b: Enrollment opens nationwide on October 26, 2021; c: Test C enrollment starts on November 4, 2021; d: Enrollment waitlist approach implemented on November 24, 2021

Figure 2: Test Us at Home Enrollment by State and Month, October 2021-January 2022


medRxiv preprint doi: <https://doi.org/10.1101/2022.08.04.22278274>; this version posted August 5, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.



Note: COVID-19 monthly incidence calculated from CDC COVID Data Tracker⁶

Figure 3: CONSORT diagram of the Test Us At Home Study

CONSORT diagram for data included in the final dataset shared with the FDA (all data)



a= participants replaced their assigned rapid antigen tests with commercially obtained rapid antigen tests; b= dates of RT-PCR testing could not be verified based on triangulation of self-reported, shipping, and resulting data; A, B, and C refer to rapid antigen test assignment. c = at-least two positive molecular assays from a single sample per participant (Roche Cobas 6800, Quest LDT, Hologic Aptima)