



## Supplementary Material


### Supplementary Figures and Tables



- Supplementary Figure 1: Kinetics of antibody levels (A) Anti-spike IgG BAU/ml and (B) Anti-spike RBD Igs (AEU), in a subgroup of 51 COVID-19 patients. Subgroup was used in the neutralizing assay. Calculations were done with the Friedman's test for repeated measures with Pairwise Wilcoxon signed-rank test for multiple comparison correction.  $P < 0.05$ ,  $**P < 0.01$ ,  $***P < 0.001$ ,  $****P < 0.0001$ .



- Supplementary Figure 2: Cross-reactive neutralization. Comparison of neutralizing ID<sub>50</sub> of samples collected early in the pandemic (pre-April 2021) during Wuhan strain prevalence and those collected later in the pandemic (post-April 2021) during the prevalence of other non-Wuhan strains such as alpha, beta, delta and omicron. Comparison's done using a Mann Whitney U test.  $P < 0.05$ ,  $**P < 0.01$ ,  $***P < 0.001$ ,  $****P < 0.0001$ .



3. Supplementary Figure 3: Viral load comparisons among clinical groups. (A) RT PCR tests done in AKUH, Nairobi and (B) KCH, Kilifi. Calculated using Kruskal-Walli's test and Man Whitney U test respectively.  $P < 0.05$ ,  $**P < 0.01$ ,  $***P < 0.001$ ,  $****P < 0.0001$ .

### Supplementary Tables

1. Supplementary Table 1: Linear mixed model outputs of association of anti-spike IgG antibodies levels with other variables. Captures patients with at least two plasma samples. Estimates indicates effect of change of the variables on measure of anti-spike IgG. The estimates are natural log transformed. Each estimate has a corresponding P value to indicate significance level.

	<i>Estimate</i>	<i>Std. Error</i>	<i>Pr(&gt; t )</i>	<i>Significance level</i>
<i>(Intercept)</i>	3.901206	0.336231	$<2e-16$	***
<i>Time</i>	0.355905	0.026349	$< 2e-16$	***
<i>Severity</i>	0.184741	0.058897	0.00213	**
<i>Age</i>	0.013027	0.005742	0.02491	*
<i>Gender</i>	-0.146325	0.152025	0.33760	ns

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

2. Supplementary Table 2: Modelling of neutralizing antibody kinetics. Estimates indicates effect of change of the variables on measure of neutralizing antibodies (Day 0 used as reference point). Fitted effects are obtained from a linear mixed model output. Each estimate has a corresponding P value to indicate significance level.

	<b>Estimate (S.E)</b>	<b>Fitted Values (95% C.I)</b>	<b>p-value</b>
Day 0	ref	862.99 (104.9-1621.1)	
Day 28	1066.9 (403.4)	1929.93 (1171.8-2688.1)	<i>0.00949</i>
Day 180	-232.1 (403.4)	630.94 (-127.1-1389.1)	<i>0.5664</i>