

Supplementary Table S1. Association analysis of each *HLA-DRB1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39)	Relapse (-: n=160)	P_{uncorr}^1	OR (95%CI)	log-rank P_{uncorr}^2
	n (%)	n (%)			
<i>DRB1*01:01</i>	5 (12.8)	28 (17.5)	0.63	0.69 (0.25-1.93)	0.53
<i>DRB1*04:01</i>	1 (2.6)	1 (0.6)	0.35	4.18 (0.26-68.4)	0.37
<i>DRB1*04:05</i>	7 (17.9)	35 (21.9)	0.67	0.78 (0.32-1.92)	0.58
<i>DRB1*04:06</i>	4 (10.3)	8 (5.0)	0.26	2.17 (0.62-7.62)	0.25
<i>DRB1*04:10</i>	1 (2.6)	3 (1.9)	0.59	1.38 (0.14-13.6)	0.65
<i>DRB1*08:02</i> ³	7 (17.9)	11 (6.9)	0.055	2.96 (1.07-8.23)	0.040
<i>DRB1*08:03</i>	4 (10.3)	15 (9.4)	0.77	1.10 (0.35-3.53)	0.83
<i>DRB1*09:01</i> ³	22 (56.4)	62 (38.8)	0.049	2.05 (1.01-4.15)	0.049
<i>DRB1*11:01</i>	1 (2.6)	8 (5.0)	1.00	0.50 (0.06-4.12)	0.47
<i>DRB1*12:01</i>	3 (7.7)	11 (6.9)	0.74	1.13 (0.30-4.26)	0.87
<i>DRB1*12:02</i>	1 (2.6)	3 (1.9)	0.59	1.38 (0.14-13.6)	0.58
<i>DRB1*13:02</i>	1 (2.6)	14 (8.8)	0.31	0.27 (0.03-2.15)	0.22
<i>DRB1*14:05</i>	1 (2.6)	5 (3.1)	1.00	0.82 (0.09-7.19)	0.79
<i>DRB1*14*54</i>	3 (7.7)	15 (9.4)	1.00	0.81 (0.22-2.93)	0.79
<i>DRB1*15:01</i>	5 (12.8)	16 (10.0)	0.57	1.32 (0.45-3.86)	0.60
<i>DRB1*15:02</i>	9 (23.1)	34 (21.3)	0.83	1.11 (0.48-2.56)	0.80
<i>DRB1*16:02</i>	1 (2.6)	3 (1.9)	0.59	1.38 (0.14-13.6)	0.74

OR: odds ratio, CI: confidence interval.

*¹Carrier frequency (%) of each *HLA-DRB1* allele was compared between the patients with and without relapse by Fisher's exact test. Uncorrected P values (P_{uncorr}) are shown.

*²Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DRB1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.

*³For *DRB1*08:02* and *DRB1*09:01*, FDR Q values are shown in the main document (Table 4).

Supplementary Table S2. Association analysis of each *HLA-DQA1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39)	Relapse (-: n=159)	P _{uncorr} ^{*1}	OR (95%CI)	log-rank P _{uncorr} ^{*2}
	n (%)	n (%)			
<i>DQA1*01:01</i>	5 (12.8)	28 (17.6)	0.63	0.69 (0.25-1.91)	0.52
<i>DQA1*01:02</i>	7 (17.9)	30 (18.9)	1.0	0.94 (0.38-2.33)	0.96
<i>DQA1*01:03</i>	13 (33.3)	47 (29.6)	0.70	1.19 (0.56-2.52)	0.65
<i>DQA1*01:04</i>	3 (7.7)	21 (13.2)	0.42	0.55 (0.15-1.94)	0.38
<i>DQA1*03:01</i>	7 (17.9)	27 (17)	1.0	1.07 (0.43-2.67)	0.94
<i>DQA1*03:02</i> ^{*3}	24 (61.5)	64 (40.3)	0.020	2.38 (1.16-4.87)	0.020
<i>DQA1*03:03</i>	9 (23.1)	37 (23.3)	1.0	0.99 (0.43-2.27)	0.97
<i>DQA1*04:01</i>	4 (10.3)	5 (3.1)	0.077	3.52 (0.90-13.8)	0.075
<i>DQA1*05:05</i>	1 (2.6)	14 (8.8)	0.31	0.27 (0.03-2.14)	0.18
<i>DQA1*05:08</i>	2 (5.1)	3 (1.9)	0.26	2.81 (0.45-17.4)	0.18
<i>DQA1*06:01</i>	1 (2.6)	4 (2.5)	1.0	1.02 (0.11-9.39)	0.70

OR: odds ratio, CI: confidence interval.

^{*1}Carrier frequency (%) of each *HLA-DQA1* allele was compared between the patients with and without relapse by Fisher's exact test.

Uncorrected P values (P_{uncorr}) are shown.

^{*2}Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DQA1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.

*³ For *DQA1*03:02*, FDR Q value is shown in the main document (Table 4).

Supplementary Table S3. Association analysis of each *HLA-DQB1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39)	Relapse (-: n=160)	P_{uncorr}^{*1}	OR (95%CI)	log-rank P_{uncorr}^{*2}
	n (%)	n (%)			
<i>DQB1*03:01</i>	5 (12.8)	36 (22.5)	0.27	0.51 (0.18-1.39)	0.21
<i>DQB1*03:02</i>	7 (17.9)	27 (16.9)	0.82	1.08 (0.43-2.69)	0.93
<i>DQB1*03:03</i> ^{*3}	23 (59.0)	65 (40.6)	0.048	2.10 (1.03-4.28)	0.043
<i>DQB1*04:01</i>	7 (17.9)	34 (21.3)	0.83	0.81 (0.33-2.00)	0.63
<i>DQB1*04:02</i>	5 (12.8)	10 (6.3)	0.18	2.21 (0.71-6.87)	0.18
<i>DQB1*05:01</i>	5 (12.8)	28 (17.5)	0.63	0.69 (0.25-1.93)	0.53
<i>DQB1*05:02</i>	2 (5.1)	12 (7.5)	1.0	0.67 (0.14-3.11)	0.65
<i>DQB1*05:03</i>	2 (5.1)	12 (7.5)	1.0	0.67 (0.14-3.11)	0.63
<i>DQB1*06:01</i>	13 (33.3)	47 (29.4)	0.70	1.20 (0.57-2.54)	0.63
<i>DQB1*06:02</i>	5 (12.8)	15 (9.4)	0.55	1.42 (0.48-4.18)	0.50
<i>DQB1*06:04</i>	1 (2.6)	13 (8.1)	0.31	0.30 (0.04-2.35)	0.26

OR: odds ratio, CI: confidence interval.

^{*1}Carrier frequency (%) of each *HLA-DQB1* allele was compared between the patients with and without relapse by Fisher's exact test.

Uncorrected P values (P_{uncorr}) are shown.

^{*2}Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DQB1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.

*³For *DQB1*03:03*, FDR Q value is shown in the main document (Table 4).

Supplementary Table S4. Association analysis of each *HLA-DPB1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39)	Relapse (-: n=160)	P _{uncorr} ^{*1}	OR (95%CI)	log-rank P _{uncorr} ^{*2}
	n (%)	n (%)			
<i>DPB1*02:01</i>	22 (56.4)	72 (45.0)	0.22	1.58 (0.78-3.20)	0.27
<i>DPB1*02:02</i>	2 (5.1)	13 (8.1)	0.74	0.61 (0.13-2.83)	0.58
<i>DPB1*03:01</i>	5 (12.8)	14 (8.8)	0.54	1.53 (0.52-4.55)	0.38
<i>DPB1*04:01</i>	1 (2.6)	10 (6.3)	0.70	0.39 (0.05-3.18)	0.41
<i>DPB1*04:02</i>	8 (20.5)	42 (26.3)	0.54	0.73 (0.31-1.70)	0.54
<i>DPB1*05:01</i>	22 (56.4)	87(54.4)	0.86	1.09 (0.54-2.20)	0.81
<i>DPB1*09:01</i>	8 (20.5)	31 (19.4)	0.83	1.07 (0.45-2.56)	0.89
<i>DPB1*13:01</i>	2 (5.1)	4 (2.5)	0.33	2.11 (0.37-11.9)	0.45

OR: odds ratio, CI: confidence interval.

*¹Carrier frequency (%) of each *HLA-DPB1* allele was compared between the patients with and without relapse by Fisher's exact test.

Uncorrected P values (P_{uncorr}) are shown.

*²Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DPB1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.