1 Genomic surveillance reveals early detection and transition of Delta to Omicron

2 Lineages of SARS-CoV-2 Variants in wastewater treatment plants of Pune, India

- ³ Vinay Rajput ^{a,b,1}, Rinka Pramanik^{a,1}, Vinita Malik^a, Rakeshkumar Yadav^{a,b}, Pradnya
- 4 Kadam^c, Unnati Bhalerao^c, Manisha Tupekar^c, Dipti Deshpande^c, Priyanki Shah^d, LS
- 5 Shashidhara^{c,d}, Radhika Boargaonkar^e, Dhawal Patil^e, Saurabh Kale^e, Asim Bhalerao^f,
- Nidhi Jain^f, Sanjay Kamble^g, Syed Dastager^{a,b}, Krishanpal Karmodiya^c, Mahesh
 Dharne^{a,b*}
- ⁸ ^a National Collection of Industrial Microorganisms (NCIM), Biochemical Sciences
- 9 Division, CSIR-National Chemical Laboratory (NCL), Pune 411008, Maharashtra,
- 10 India
- ^b Academy of Scientific and Innovative Research (AcSIR), Ghaziabad 201002, Uttar
 Pradesh, India
- ^c Department of Biology, Indian Institute of Science Education and Research (IISER),
- 14 Pune 41108, Maharashtra, India
- ^d The Pune Knowledge Cluster (PKC), Savitribai Phule Pune University (SPPU),
- 16 Pune, Maharashtra, India
- ^e Ecosan Services Foundation (ESF), Pune 411030, Maharashtra, India
- ^f Fluid Robotics Private Limited (FRPL), Pune 411052, Maharashtra, India
- ^g Chemical Engineering and Process Development Division, CSIR-National Chemical
- 20 Laboratory, Pune 411008, Maharashtra, India
- 21
- ²² ¹These authors contributed equally
- 23
- 24
- 25 * Corresponding Author Details: National Collection of Industrial Microorganisms
- 26 (NCIM), Biochemical Sciences Division, CSIR-National Chemical Laboratory, Pune
- 27 411008, India. E-mail address: ms.dharne@ncl.res.in (Mahesh S. Dharne).

28 Highlights

- Omicron fragments were detected in the sewershed samples prior to clinical
 samples.
- Omicron sub-lineages BA.2.12, BA.2.38, BA.2.75, BA.3, BA.4, and XBB were
 prevalent.
- Lineage composition analysis indicated transition from Delta to Omicron variant
 indicated cause of third wave in India.
- Overall, 426 lineages of 17 highly transmissible variants of SARS-CoV-2 were
 detected in the study.

37

38 Abstract

The COVID-19 pandemic has emphasized the urgency for rapid public health 39 40 surveillance methods in early detection and monitoring of the transmission of infectious diseases. The wastewater-based epidemiology (WBE) has emerged as a 41 promising tool to analyze and enumerate the prevalence of infectious pathogens in a 42 population ahead of time. In the present study, real time quantitative polymerase chain 43 reaction (RT-qPCR) and Illumina sequencing was performed to determine the SARS-44 CoV-2 load trend and dynamics of variants over a longitudinal scale in 442 45 wastewater (WW) samples collected from 10 sewage treatment plants (STPs) of Pune 46 47 city, India, during November 2021 to April-2022. In total 426 distinct lineages 48 representing 17 highly transmissible variants of SARS-CoV-2 were identified. The SARS-CoV-2 Omicron variant fragments were detected in WW samples prior to its 49 detection in clinical cases. Moreover, highly contagious sub-lineages of Omicron, 50 such as BA.2.12 (0.8-0.25%), BA.2.38 (0.8-1.0%), BA.2.75 (0.01-0.02%), BA.3 51 (0.09-6.3%), BA.4 (0.24-0.29%), and XBB (0.01-13.7%) fragments were significantly 52 53 detected. The longitudinal analysis also suggested the presence of the BA.5 lineage in 54 November 2021, which was not reported in the clinical settings throughout the 55 duration of this study, indicative of silent variant persistence. Overall, the present study demonstrated the practicality of WBE in early detection of SARS CoV-2 56 variants, which could be useful in tracking future outbreaks of SARS-CoV-2. Such 57 58 approaches could be implicated in the monitoring of the infectious agents before they 59 appear in clinical cases.

Keywords: SARS-CoV-2; wastewater-based epidemiology; Early warning; Next generation sequencing; COVID-19; wastewater

62 Abbreviations

Wastewater, WW; COVID-19, coronavirus disease 19; SARS-CoV-2, severe acute
respiratory syndrome coronavirus 2; WHO, The World Health Organization; VOI,
variants of interest; VOC, variants of concern; WBE, wastewater-based epidemiology;
E, envelope E; N, nucleocapsid; WWTP, wastewater treatment plants; PEG-8000,
polyethylene glycol-8000; IISER, Indian Institute of Science Education and Research,
NCL, National Chemical Laboratory

69 **1.** Introduction

The SARS-CoV-2 pandemic, initially detected in Wuhan, China in 2019, 70 71 spreads rapidly all over the world (Global research on coronavirus disease (COVID-19)," W.H.O). As the virus spreads, the emergence of new variants has become a 72 concern for public health officials. The clinical testings have limitations in detecting 73 74 and tracking the spread of variants (Faria et al., 2021; Li et al., 2020). Also, the SARS-CoV-2 virus epidemic has sparked global efforts to understand its transmission 75 dynamics and the emergence of new variations (Kucirka et al., 2020; Rambaut et al., 76 77 2020). To address this challenge, researchers have begun exploring alternative methods for monitoring the spread of SARS-CoV-2, such as wastewater-based 78 79 epidemiology (WBE) (de Roda Husman et al., 2020; van der Meer et al., 2020).

80 WBE involves analyzing viral particles present in sewage to infer the spread of infection within a population. WBE has been used to study other viral diseases, 81 82 such as influenza and hepatitis E (Heijnen and Medema, 2011; Rosa et al., 2014), and has recently been applied to SARS-CoV-2. Studies by Peccia et al., 2020b and Ahmed 83 84 et al., 2022 have provided essential insights into the feasibility and applicability of WBE for SARS-CoV-2, highlighting its potential as an early detection system for 85 population infection dynamics. Several studies have been conducted globally, with 86 notable contributions from researchers in Germany, Italy, Netherlands, Finland, USA, 87 88 Japan, France, Canada and India (Maida et al., 2022; Tiwari et al., 2022b; Prado et al., 89 2021; Stephens et al., 2022; Ho et al., 2022; Amman et al., 2022; Kumar et al., 2020). 90 These studies provides essential information on the detection, quantification and 91 genetic characterization of SARS-CoV-2 in sewage samples (Srivastava et al., 2021; Dharmadhikari et al., 2022; Lamba et al., 2023). The WBE has the potential to detect 92 viral signals earlier than clinical surveillance methods and can offer a less biased 93 assessment of viral heterogeneity and public health (Joshi et al., 2022; Jahn et al., 94 95 2022; Morvan et al., 2022). The ongoing COVID-19 outbreak has brought attention to 96 the need for rapid and accurate surveillance of the spread of SARS-CoV-2.

The impact of SARS-CoV-2 has been felt globally since its emergence in 2019
(Chakraborty and Maity, 2020). The virus, which causes COVID-19, has resulted in
substantial interruptions to everyday life and economic activities, as well as
widespread disease and death. (Shang et al., 2021, Li et al., 2020). One of the critical

101 concerns in the COVID-19 epidemic is the advent of novel SARS-CoV-2 variants, 102 which may be more transmissible and lead to more severe disease (Aleem et al., 2022). 103 Omicron variant is known to be associated with a high number of cases across the globe (Nyberg et al., 2022). The virus has significantly impacted India, with several 104 105 clinical cases and deaths reported (Ranjan, 2022). Among the Indian cities affected, 106 Pune has recorded one of the highest numbers of SARS-CoV-2 Omicron cases. To 107 comprehend the propagation and evolution of SARS-CoV-2, researchers have used 108 next-generation sequencing (NGS) platforms to analyze genomic sequences of the 109 virus. These studies have shown that different SARS-CoV-2 variants have emerged in 110 vast geographical areas, with varying infectivity and transmission capability levels. In 111 light of this, NGS platforms have proven to be a valuable tool in understanding the evolution of the virus and the dangers it poses to society. Studies by Agrawal et al., 112 113 2022 in Germany, Landgraff et al., 2021 in Canada, Wilton, 2021 in London, Crits-Christoph, 2021 in California, and Dharmadhikari et al., 2022 in India have used NGS 114 platforms to identify SARS-CoV-2 mutations in wastewater. These studies have 115 116 proven that WBE can capture SARS-CoV-2 and its mutants in wastewater, providing 117 insight into the spread of the virus in communities.

118 The present study is the first in-depth longitudinal study employing 119 wastewater-based SARS-CoV-2 sequencing to delineate the overall dynamics of 120 SARS-CoV-2 lineages in Pune city, India. Here, we utilized WBE to monitor the 121 emergence of SARS-CoV-2 variants, specifically the Omicron lineage in Pune City, 122 analyzing its presence in wastewater samples and assessing its impact on the local 123 population and healthcare system. Understanding the dynamics of the Omicron 124 variant and its sub-lineages will provide insights that can inform public health response and help mitigate the spread of the virus. To achieve our objectives, we 125 126 systematically analyzed wastewater samples collected from different locations in 127 Pune City, using RT-qPCR and sequencing techniques to detect and quantify the presence of the Omicron variant. This study will provide critical insights into the 128 129 dynamics of the Omicron variant in Pune City and inform the development of 130 effective public health measures to combat its spread.

131 **2.** Materials and Methods

132 2.1. Wastewater sample collection and SARS-CoV-2 sequencing

Sewage wastewater samples were taken from the inlets of 10 sewage treatment 133 plants (STPs) in Pune (Fig.1, Table.S1). The sample collection included a grab-134 135 sampling method, executed following the CDC, USA guidelines of COVID-19 wastewater sampling, and approved by the Institutional Biosafety Committee (IBSC). 136 137 Sampling was carried out two times in a week from 10 STPs, in the period of 138 November 2021 - April 2022 in 1L sterile polypropylene bottles and stored at 4°C until processing. The processing was carried out immediately as per CDC, USA 139 140 guidelines of COVID-19 sewage water sampling. The sampling bottle was wiped 141 with 70% ethanol and subjected to heat inactivation at 60°C. 80 ml of the sample was aliquoted in a 250 ml HDPE bottle (HiMedia, India) and subjected to viral RNA 142 143 concentration using the polyethylene glycol (PEG) precipitation method. The detailed protocol of sample processing and RT-qPCR is given in supplementary information 144 (Supplementary File.2) Sequencing was carried out at Next Generation Genomics 145 facility at IISER, Pune, India. The library preparation was carried out using Illumina 146 147 Covid seq RUO kits (Illumina, USA). Sequencing was carried out using Illumina 148 NextSeq 550 sequencer.

149 2.2. Bioinformatics Analysis

In this study, an in-house bioinformatics pipeline was designed to identify the 150 predominant lineages and their signature mutations of SARS-CoV-2 in the sample. 151 152 The pipeline consisted of several crucial steps to ensure the accuracy of the results. 153 The first step involved quality control and filtering of the raw reads using fastp (Chen 154 et al., 2018), which helps to ensure the quality of the data being analyzed. The filtered 155 reads were then aligned with the reference genome of SARS-CoV-2 (MN908947.3) 156 using BWA-Mem (Li and Durbin, 2010), providing basic alignment and coverage 157 statistics using SAMtools coverage/bedcov (Li et al., 2009). To determine single nucleotide variants (SNVs), iVar (Grubaugh et al., 2019) tool was used with 158 159 parameters of a minimum base quality score of 20, minimum frequency of 0.03, and minimum read depth of 10. To predict the lineage composition of the SARS-CoV-2 160 161 variants in a sample, we utilized LCS (Valieris et al., 2022b), a mixture model designed to determine the variant composition in environmental samples. We used 162 163 LCS lineage abundance estimates to compute Shannon diversity and Bray-Curtis dissimilarity metrics, as well as an Adonis statistical analysis using R-programming 164

packages such as vegan v2.5-7, ggplot, and phyloseq. Finally, data analysis was
performed using scripting in R/Python/Bash to manually check variant-associated
mutations and to plot the results.

168 **3. Results**

169 **3.1.** Quantification and correlation of SARS-CoV2

170 We analyzed data collected from 10 sewage treatment plants (STPs) located across the Pune city, covering an estimated 92% of the city's population between 171 172 November 2021 and April 2022 (Fig.1, Table.S1). An average of two samples were taken from each location per week to assess the level of community spread. 173 Wastewater samples were processed and analyzed within 24h of sample collection. 174 175 RT-qPCR was employed to test 442 samples for the presence of SARS-CoV-2 genetic material using three different genetic target regions: N, RdRp, and E gene. A sample 176 177 was considered positive for SARS-CoV-2 when any two genes showed Ct values 178 lower than 35. In total, 142 out of 442 samples were positive, while the remaining 300 179 were negative, representing a total positive detection rate of 32.12%. (Table.S2)

180 To gain insights to the prevalence of SARS-CoV-2 infection in the community, we calculated the amount of virus present in each sample using a viral load 181 calculation tool. The results of this analysis, depicting the levels of SARS-CoV-2 in 182 183 wastewater and clinically reported in Pune cases (http://cms.unipune.ac.in/~bspujari/Covid19/Pune2/) are illustrated in (Fig.2, Fig.3). 184 185 A strong correlation for the entire city viral load exhibits a significant association with daily clinically reported COVID-19 positive cases ($R^2=0.81$, p<22e-16; Fig.S1.A) and 186 7-day moving average COVID-19 cases (R²=0.83, p<22e-16; Fig.S1.B). The RT-187 qPCR analysis revealed an increase in viral load during the SARS-CoV-2 surge in 188 January 2022, with the virus detected in the samples throughout the period from 189 November 2021 to April 2022. The viral load was observed lowered in November and 190 191 December 2021, with an average viral copy of 3.2 gc/ml and 2.2 gc/ml, respectively, 192 while a sharp increase in viral load was noted in January 2023, with an average 193 detection of 83.9 gc/ml. Overall wastewater viral trend forms an early mirror image over clinically reported COVID-19 cases, demonstrating the usefulness of WBE in 194 195 detecting the early transmission of SARS-CoV-2 infection in the community.

196 The SARS-CoV-2 signals were detected across all STPs throughout the study. When COVID-19 entered Pune in its third wave, all STPs began to give early 197 198 warnings 08 to 17 days before the city's clinical COVID-19 cases substantially increased. STP sites such as Old Naidu, Mundhwa, and New Naidu and Kharadi gave 199 early warning signals from December 8th to December 16th, 2021, sequentially, 200 whereas STP sites like Tanajiwadi, BhairobaNala, Bopodi, and Erandwane began 201 providing early warning signals after December 20th, 2021. However, STP sites of 202 Vitthalwadi and Baner gave signals at the same time the clinical cases substantially 203 204 started increasing (Fig.2, Fig.3), which probably indicates the effective clinical testing in the region. This phenomenon of detecting 08 to 17 days early viral signals from 205 206 wastewater proves the usefulness of WBE as a system for early detection.

3.2. Prediction and diversity of SARS-CoV-2 variants in wastewater

208 To understand the dynamics of SARS-CoV-2 variants in the city, we conducted a large-scale SARS-CoV-2 sequencing effort on 442 samples collected 209 210 over six months from November 2021 to April 2022 (Table.S3). A significant 211 proportion of the samples which is 52%, failed to yield sufficient sequencing data due 212 to low viral load as evidenced by high cycle threshold (Ct) values. However, the remaining 48% of samples met our stringent quality criteria, including detection 213 214 confidence of SARS-CoV-2 and minimum genome coverage of 55%, and were 215 subsequently used for lineage prediction analysis. The study's findings offer insightful 216 information on the evolution and propagation of SARS-CoV-2 variants within our 217 study population.

The relative frequencies and diversity of SARS-CoV-2 lineages in wastewater 218 219 were analyzed using the LCS tool in a study conducted from November 2021 to April 220 2022. A total of 426 distinct lineages representing 17 highly transmissible variants of 221 SARS-CoV-2 were identified in WW during this period (Fig.2, Fig.3). The Delta 222 lineage was found to be the most dominant during November and December 2021, with an overall percentage of 80% and 55% respectively (Table.S5), followed by the 223 224 Kappa lineage, which accounted for 12% and 30% of the total lineages during the same period, while BA.3 and BA.2 lineages accounts for 2.4-6.3% and 0-3% 225 respectively. The BA.1 lineage accounted for only 0.3% and 0.2% in January and 226 227 February'2022, respectively. The remaining 8% of lineages were classified as "others".

228 However, small signals of the Omicron lineage started to be detected during 229 November 2021 (2.6%) and December 2021 (7.9%), with a 10-fold increase in 230 January 2022 (70%) to become the most prevalent lineage from January 2022 onwards (Fig.4). Interestingly, the study also detected strong signals of the XBB 231 232 lineage in March samples accounts for 13.7% to total lineages. This emphasizes the 233 significance of continuing monitoring of SARS-CoV-2 variants in WW to better 234 understand their spread and transmission dynamics. Identifying highly transmissible 235 variants in WW may also affect public health interventions and measures.

3.3. Diversity and correlation between Wastewater and Clinical SARS-CoV-2 data

238 We aimed to correlate the lineage composition between clinical and 239 wastewater samples of SARS-CoV-2 in Pune city, India. We retrieved 3223 SARS-240 CoV-2 genomes from GISAID from November 2021 to April 2022, submitted from 241 Pune city (Table.S6). A total of 98 distinct lineages and sub lineages were identified 242 during this period. Based on the predominance of each lineage in India, the lineages 243 were categorized according to the INSACOG categorization for further study 244 (INSACOG Dashboard, Table.S7). The results showed that the overall clinical 245 lineages followed a similar pattern as the wastewater samples. The Omicron lineage and its sub lineages were the dominant lineages from January to April 2022. Omicron 246 247 BA.2 accounted for 51.6% of all lineages, followed by Omicron BA.2.10 (14.8%), 248 Omicron BA.2.X (13.8%), Omicron BA.1.X (6.3%), Omicron BA.1 (5.6%), and 249 Omicron BA.2.38 (4.9%). (Fig.S3). However, the Delta lineage was the most 250 dominant lineage during November and December 2021. Specifically, the 251 Delta_B.1.617.2 lineage accounted for 88.1% of the total lineages, followed by early 252 signals of Omicron_BA.1.X (5.2%), Omicron_BA.2 (2.8%), and Omicron_BA.1 253 (2.1%). Similarly, the wastewater samples had dominant lineages of Delta_B.1.617.2 254 and Kappa during November and December 2021. Later, the Omicron variant takeover the Delta variant and became a dominant variant by January' 2022 (Fig.4). 255 256 This transition from Delta to the omicron variant highlights the significance of ongoing monitoring and observation of SARS-CoV-2 lineages in both clinical and 257 258 wastewater samples to gain a better understanding of their spread and transmission 259 dynamics.

260 To understand the diversity of SARS-CoV-2 lineages circulating in the 261 community, we calculated Simpson indices for all the samples collected from various 262 sites. These indices measure the total diversity of a community and are commonly used in ecology and epidemiology. The results of our analysis revealed a significant 263 264 difference in diversity between months (Kruskal-Wallis, p=2.6e-12) but no significant 265 difference by location (Fig.S2.A, Fig.S2.B). To further investigate the variation in 266 diversity across time, we used the Bray-Curtis dissimilarity matrix with the Adonis test to compute beta diversity. This approach allows for comparing community 267 composition between different samples or groups. The analysis found a significant 268 difference in beta diversity across the months (R2 = 0.55, P > 0.001) but no significant 269 270 variation by location (Fig.S2.C, Fig.S2.D). These results suggest that temporal rather than geographic factors primarily influence the diversity of SARS-CoV-2 lineages in 271 Pune. Overall, these findings discern the dynamics of SARS-CoV-2 lineage diversity 272 in the population and highlight the importance of continued monitoring to track 273 emerging variants of concern. 274

275 **3.4.** Early detection of Omicron and its sub-lineages

To determine the population's exposure to SARS-CoV-2 variants, we utilized 276 the iVar tool to analyze each sample for co-occurring signature mutations of the 277 278 Omicron variant (BA.1, BA.2, BA.3, BA.4, BA.5, BA.1.x, BA.2.x, BA.3.x, BA.4.x, 279 BA.5.x). Despite some limitations in the sequencing data due to low RNA 280 concentrations in certain samples, we could still detect characteristic signals of the 281 Omicron variant. Our analysis revealed that wastewater samples taken from the STP at Bhairobanala was the first one to have the Omicron signals : S: D614G, S: H655Y, 282 S: N679K, S: P681H, N: P13L) and Tanajiwadi (S: N440K, S: S477N, S: T478K, S: 283 284 E484A, S: Q493R, S: Q498R, S: N501Y, S: Y505H, S: T547K, S: D614G, E: T9I, on November 1st, 2021, before it was classified as a VOC by the World Health 285 286 Organization ("Tracking SARS-CoV-2 variants," WHO) on November 26th, 2021 (Table.S8). The Omicron was also detected in samples collected from the STP at 287 288 Bopodi ((S: L452R, S: T478K, E: T9I) on November 2nd, 2021, before it was confirmed in Pune's first clinical case on December 5th, 2021. Furthermore, our 289 290 analysis revealed that Omicron and its sub lineages were detected in wastewater 291 samples collected from nine out of ten STPs in Pune, indicating widespread

292 community exposure to this variant before its detection in clinical cases (Table.S9). 293 These findings emphasize the use of wastewater-based surveillance in tracking the 294 emergence and spread of SARS-CoV-2 variants in the community. It is important to 295 note that our investigation was carried out in India, and the Omicron variant was 296 discovered in Botswana on November 2011. As such, our data suggest that Omicron 297 may have been circulating in wastewater prior to its detection in clinical cases. The 298 findings of this investigation offer crucial insights into the early detection and spread 299 of SARS-CoV-2 in the community and support the implementation of WBE as a 300 complementary approach to traditional clinical surveillance.

301 Moreover, to evaluate the rise of variation in the lineage of SARS-CoV-2 in 302 the population, we employed a linear mixed model analysis to examine the correlation 303 between the initial detection of variant of concern in wastewater and the date when 304 clinical samples were collected from Pune city and submitted to GISAID. Our examination revealed a trend of increasing presence of the Omicron variant 305 (B.1.1.529) and its sub lineages (BA.1, BA.2, BA.2.12, BA.2.X, BA.2.38, BA.3, 306 BA.4, BA.5) starting from January 2022. As time progressed, however, we noted a 307 308 reduction in the frequency of the Kappa (B.1.617.1) and Delta (B.1.617.2) lineages, with the Omicron variant completely dominating by February 2022 (Fig.5). 309 310 Additionally, we observed an upward trend in the prevalence of the XBB lineage 311 starting from January 2022. These findings emphasize the crucial importance of ongoing monitoring of the variations in SARS-CoV-2 in order to inform public health 312 313 strategies.

314 4. **Discussion**

The emergence of new SARS-CoV-2 variants has been a key cause of concern 315 316 in the ongoing COVID-19 epidemic. One such variant, Omicron, has been identified in India (Gao et al., 2021). This variant has raised concerns due to its high 317 318 transmissibility and potential to evade immunity from previous infections or vaccines (Shrestha et al., 2021; Zhang, 2022; Yadav and Kumar, 2022). Our study utilizes 319 WBE to identify, track, and assess the SARS-CoV-2 genetic variants in Pune city, 320 India. We collected 442 wastewater samples from 10 locations, representing an 321 322 estimated 7.4 million individuals. WBE has been used in several studies globally to 323 identify and quantify the SARS-CoV-2 fragments in wastewater (Srivastava et al.,

2021; Dharmadhikari et al., 2022; Joshi et al., 2022; Jahn et al., 2022; Morvan et al., 2022). By applying this model at the city level, we aim to prevent the spread of COVID-19 and better understand the dynamics of SARS-CoV-2 transmission in Pune city.

328 In this study, we used RT-qPCR and Illumina sequencing to quantify and 329 monitor the genetic characteristics of SARS-CoV-2 variants. Our findings show a 330 strong correlation between the viral load and clinical cases of COVID-19. 331 Additionally, our data suggest that a significant amount of viral shedding likely 332 occurs early in infection, particularly from asymptomatic individuals, before they seek medical attention and are tested for Covid-19 (Lamba et al., 2023). The SARS-CoV-2 333 334 persists in faeces samples longer than in respiratory and serum samples (Prakash et al., 2021). In the present study, the SARS-CoV-2 was detected before an increase in 335 clinical cases highlighting the potential utility of WBE as a tool for early detection 336 and prevention of SARS-CoV-2 transmission. By monitoring STPs, public health 337 officials can identify outbreaks and implement strict management strategies at the 338 339 earliest stages of severe disease.

340 The SARS-CoV-2 genetic material was quantified using RT-qPCR, followed 341 by Illumina sequencing to determine the trends of SARS-CoV-2 lineages disseminated in the community. Detecting SARS-CoV-2 titers in wastewater is 342 challenging amid the fragmentation and PCR inhibitors, making it difficult to quantify 343 344 and sequence for estimating lineage/variant composition (Baaijens et al., 2021). 345 However, in this study, LCS (Lineage decomposition for SARS-CoV-2), a model for 346 predicting the lineage composition in a sample, was utilized, which captures and 347 classifies variant-specific mutations (Valieris et al., 2022). Our results showed that 348 relative to the clinical testing, SARS-CoV-2 RNA content in wastewater provides a 349 more accurate indicator for local infection dynamics (Karthikeyan, 2021); Randazzo 350 et al., 2020; Polo et al., 2020). We found a wide range of SARS-CoV-2 lineages in 351 wastewater, with some lineages that infect a substantial proportion of people 352 exhibiting similar patterns. Additionally, we discovered low prevalent mixed lineages in wastewater that were undetectable from clinical data, suggesting that such lineages 353 354 infected only a small number of individuals.

Furthermore, weak signals from novel variants imply the arrival of a new lineage into the population that could not be emphasized in clinical data- indicating the limited testing or testing for symptomatic individuals exclusively. We discovered

358 early XBB signals and other Omicron sub-lineages that were not previously reported 359 in clinical data. This discrepancy highlights the lack of required sequencing, i.e. at 360 least 0.5% of the positive cases (Grimaldi et al., 2022), which impacts the ability to detect SARS-CoV-2 variants quickly and undermines the global pandemic 361 362 preparedness. Our research emphasizes the effects of socioeconomic disparities, 363 population density, and wastewater treatment as powerful strategies to stop the spread 364 of SARS-CoV-2 (Sharifi and Khavarian-Garmsir, 2020). It is imperative to continue 365 monitoring and testing wastewater for SARS-CoV-2 to prevent the spread of new 366 variants and to ensure global pandemic preparedness. We aimed to quantify the 367 SARS-CoV-2 genetic material in wastewater samples and determine the genomic 368 trends of SARS-CoV-2 lineages circulating in the community. Our results showed that wastewater samples exhibit greater alpha diversity than clinical samples. 369 Additionally, the viral composition of the city did not show regional variability, as the 370 beta diversity of SARS-CoV-2 did not differ by STP locations. However, there was a 371 significant variation (35%) across clinical and wastewater samples, and distinct and 372 373 common clusters of SARS-CoV-2 lineages were formed over time, suggesting 374 introduction and elimination of various lineages in the community over time 375 (Karthikeyan et al., 2022; Gu et al., 2022). These findings emphasize the significance 376 of wastewater surveillance as a complementary tool to clinical testing in understanding the dynamics of SARS-CoV-2 transmission in the community. 377

378 We investigated the presence of different lineages and sub-lineages of SARS-379 CoV-2 in wastewater samples collected from Pune, India. Our findings showed the presence of several variants of the Omicron lineage, which have been linked to 380 381 increased transmissibility, adverse health impacts, and hospitalization (Khandia et al., 382 2022; Araf et al., 2022.). Importantly, our results indicated that three out of the ten 383 catchment areas had detectable levels of Omicron in the wastewater samples before 384 the first clinical case of Omicron-positive in Botswana on November 11th, 2021: 10 385 days earlier in Bhairobanala, 9 days earlier in Bopodi, and 10 days earlier in 386 Tanajiwadi. Additionally, nine out of the ten catchment areas had Omicron signals in the wastewater samples before the first reported clinical case of Omicron in Pune city 387 388 on December 5th, 2021: 19 days earlier in Baner, 34 days earlier in Bhairobanala, 33 389 days earlier in Bopodi, 23 days earlier in Erandwane, 24 days earlier in Kharadi, 18 390 days earlier in New Naidu, 25 days earlier in old Naidu, 34 days earlier in Tanajiwadi, 391 and 24 days earlier in Mundhwa. This implies that a substantial portion of the

population was infected and shedding the virus before being diagnosed throughclinical testing (He et al., 2020).

394 Furthermore, we discovered six highly transmissible lineages of Omicron in 395 the wastewater samples before their detection in clinical samples from the city: 396 BA.2.12, BA.2.38, BA.2.75, BA.3, BA.4, and XBB detected 20, 8, 153, 242, 127, and 397 279 days earlier, respectively. Additionally, we found weak indications of lineage 398 BA.5 in November 2022, which has yet to be reported in any clinical cases from Pune 399 city. Our findings demonstrate the utility of WBE for detecting and monitoring the 400 presence of SARS-CoV-2 variants in the community. Testing of wastewater captures 401 the combined contributions of every individual in a catchment region, providing a less 402 biased representation of viral diversity and public health (Karthikeyan et al., 2022). In 403 contrast, delays in detection from clinical samples may be due to limited testing and 404 sequencing or asymptomatic or home-testing cases (Mercer and Salit, 2021). 405 Additionally, our study showed that sequencing data from wastewater samples could be used to monitor the prevalence of variants and estimate their growth rate, using 406 407 significantly fewer samples than required for clinical samples. Overall, these results 408 substantiates the importance of WBE as a complementary approach to traditional 409 clinical surveillance for detecting and monitoring the spread of SARS-CoV-2 variants. 410 Furthermore, it also highlights the potential of WBE to detect the emergence of new variants before they are identified in clinical samples. 411

412 5. Conclusion

413 The present study highlighted the efficacy of wastewater-based epidemiology (WBE) 414 as a formidable tool in the early warning, detection, and monitoring of SARS-CoV-2 415 variants. By conducting a comprehensive analysis of wastewater samples collected between November 2021 and April 2022, we could detect the Omicron variant 416 417 fragments before clinical cases, providing valuable insights for public health officials 418 to implement preventative measures. Our findings highlight the potential of WBE as a practical approach for real-time surveillance of infectious outbreaks, not only for 419 420 SARS-CoV-2 but also for other pathogens. Integrating WBE into existing public health surveillance methods can significantly enhance our ability to understand the 421 422 transmission dynamics and evolution of infectious diseases and, ultimately, help in 423 preventing future outbreaks.

424 Data availability: All data produced in the present study are available upon
425 reasonable request to the authors. Clinical COVID-19 data was used from CMS
426 dashboard (http://cms.unipune.ac.in/~bspujari/Covid19/Pune2/)

427 Declaration of competing interest: The authors declare that they have no known
428 competing financial interests or personal relationships that could have appeared to
429 influence the work reported in this paper.

430 **Funding**

This work was financially supported by Rockefeller Foundation, USA (project codeGAP 332926).

433 Acknowledgments

434 Authors wish to thank Directors of CSIR-NCL, IISER-Pune, and PKC for encouragement and support. We gratefully acknowledge Pune Municipal Corporation 435 436 (PMC) for permission to carry out the sampling and use clinical COVID-19 data. 437 Authors are indebted to Prof. Aurnab Ghose, IISER-Pune (for grant administration support and discussions), Dr. Dayanand Panse-ESF (for support and pre-preparations 438 439 in sampling logistics) and Ms. Rachel Samson (for processing and RNA extractions of the subset samples). Authors also acknowledge https://biorender.com/ for creation of 440 images. The manuscript has been thoroughly checked for plagiarism content using 441 442 iThenticate software.

443 **Reference**

Agrawal, S., Orschler, L., Schubert, S., Zachmann, K., Heijnen, L., Tavazzi, 444 445 S., Gawlik, B.M., de Graaf, M., Medema, G., Lackner, S., 2022. Prevalence and 446 circulation patterns of SARS-CoV-2 variants in European sewage mirror clinical data of 54 447 European cities. Water Research 214, 118162. https://doi.org/10.1016/j.watres.2022.118162 448

Ahmed, W., Bivins, A., Metcalfe, S., Smith, W.J.M., Ziels, R., Korajkic, A.,
McMinn, B., Graber, T.E., Simpson, S.L., 2022. RT-qPCR and ATOPlex sequencing
for the sensitive detection of SARS-CoV-2 RNA for wastewater surveillance. Water
research 220, 118621. https://doi.org/10.1016/j.watres.2022.118621

Aleem, A., Samad, A.B.A., Slenker, A.K., 2022. Emerging Variants of
SARS-CoV-2 And Novel Therapeutics Against Coronavirus (COVID-19) - PubMed.

Amman, Markt, Endler, Hupfauf, Agerer, Schedl, Richter, Zechmeister,
Bicher, Heiler, Triska, Thornton, Penz, Senekowitsch, Laine, Keszei, Klimek, Nägele,
Mayr, Daleiden, Steinlechner, Niederstätter, Heidinger, Rauch, Scheffknecht, Vogl,
Weichlinger, Wagner, Slipko, Masseron, Radu, Allerberger, Popper, Bock, Schmid,
Oberacher, Kreuzinger, Insam, Bergthaler, 2022. Viral variant-resolved wastewater
surveillance of SARS-CoV-2 at national scale. Nature Biotechnology 40, 1814–1822.
https://doi.org/10.1038/s41587-022-01387-y

Araf, Y., Akter, F., Tang, Y., Fatemi, R., Parvez, Md.S.A., Zheng, C., Hossain,
Md.G., 2022. Omicron variant of SARS CoV 2: Genomics, transmissibility, and
responses to current COVID 19 vaccines. Journal of Medical Virology 94, 1825–
1832. https://doi.org/10.1002/jmv.27588

Baaijens, J.A., Zulli, A., Ott, I.M., Petrone, M.E., Alpert, T., Fauver, J.R.,
Kalinich, C.C., Vogels, C.B.F., Breban, M.I., Duvallet, C., McElroy, K., Ghaeli, N.,
Imakaev, M., Mckenzie-Bennett, M., Robison, K., Plocik, A., Schilling, R., Pierson,
M., Littlefield, R., Spencer, M., Simen, B.B., Hanage, W.P., Grubaugh, N.D., Peccia,
J., Baym, M., 2021. Variant abundance estimation for SARS-CoV-2 in wastewater
using RNA-Seq quantification. Cold Spring Harbor Laboratory.

Chakraborty, I., Maity, P., 2020. COVID-19 outbreak: Migration, effects on
society, global environment and prevention. The Science of the Total Environment
728. https://doi.org/10.1016/j.scitotenv.2020.138882

- Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one
 FASTQ preprocessor. Bioinformatics (Oxford, England) 34, i884–i890.
 https://doi.org/10.1093/bioinformatics/bty560
- 478 Crits-Christoph, A., 2021. Genome Sequencing of Sewage Detects
 479 Regionally Prevalent SARS-CoV-2 Variants. mBio.

de Roda Husman, A.M., et al., 2020. Wastewater-based epidemiology of
SARS-CoV-2. The Lancet Microbe, 1(1), e1-e3.

482	Dharmadhikari, T., Rajput, V., Yadav, R., Boargaonkar, R., Patil, D., Kale,
483	S., Kamble, S.P., Dastager, S.G., Dharne, M.S., 2022. High throughput sequencing
484	based direct detection of SARS-CoV-2 fragments in wastewater of Pune, West India.
485	Science of the Total Environment 807, 151038.
486	https://doi.org/10.1016/j.scitotenv.2021.151038
487 488	Faria, N.R., et al., 2021. Tracking changes in SARS-CoV-2 genomes to understand the evolution of the pandemic. Science, 372(6537), eabd3060.
489	Gao, S., Guo, H., Luo, G., 2021. Omicron variant (B.1.1.529) of SARS CoV-
490	2, a global urgent public health alert! Journal of Medical Virology 94, 1255-1256.
491	https://doi.org/10.1002/jmv.27491
492	GenePath Diagnostics - Covid-19 Viral Load Calculation Tool (RUO). (n.d.).
493	Retrieved January 30, 2023, from https://coviquant.genepathdx.com/
494	Global research on coronavirus disease (COVID-19) [WWW Document], n.d.
495	URL https://www.who.int/emergencies/diseases/novel-coronavirus-2019/global-
496	research-on-novel-coronavirus-2019-ncov (accessed 2.2.23).
497	Grimaldi, Panariello, Annunziata, Giuliano, Daniele, Pierri, Colantuono, Salvi,
498	Bouché, Manfredi, Cuomo, Concilio, D., Tiberio, Fiorenza, Portella, Cimmino,
499	Sorrentino, Fusco, Granata, Cerino, Limone, Atripaldi, Ballabio, Cacchiarelli, 2022.
500	Improved SARS-CoV-2 sequencing surveillance allows the identification of new
501	variants and signatures in infected patients. Genome Medicine 14, 1-15.
502	https://doi.org/10.1186/s13073-022-01098-8

Grubaugh, Gangavarapu, Quick, Matteson, Jesus, D., Main, Tan, Paul,
Brackney, Grewal, Gurfield, Rompay, V., Isern, Michael, Coffey, Loman, Andersen,
2019. An amplicon-based sequencing framework for accurately measuring intrahost
virus diversity using PrimalSeq and iVar. Genome Biology 20, 1–19.
https://doi.org/10.1186/s13059-018-1618-7

Gu, Xie, Adam, Tsui, Chu, Chang, Cheuk, Gurung, Krishnan, Ng, Liu, Wan,
Cheng, Edwards, Leung, Wu, Tsang, Leung, Cowling, Peiris, Lam, Dhanasekaran,
Poon, 2022. Genomic epidemiology of SARS-CoV-2 under an elimination strategy in
Hong Kong. Nature Communications 13, 1–10. https://doi.org/10.1038/s41467-02228420-7

He, Lau, Wu, Deng, Wang, Hao, Lau, Wong, Guan, Tan, Mo, Chen, Liao,
Chen, Hu, Zhang, Zhong, Wu, Zhao, Zhang, Cowling, Li, Leung, 2020. Temporal
dynamics in viral shedding and transmissibility of COVID-19. Nature Medicine 26,
672–675. https://doi.org/10.1038/s41591-020-0869-5

- Heijnen, L., Medema, G., 2011. Surveillance of influenza A and the
 pandemic influenza A (H1N1) 2009 in sewage and surface water in the Netherlands PubMed. Journal of water and health 9. https://doi.org/10.2166/wh.2011.019
- Ho, J., Stange, C., Suhrborg, R., Wurzbacher, C., Drewes, J.E., Tiehm, A.,
 2022. SARS-CoV-2 wastewater surveillance in Germany: Long-term RT-digital
 droplet PCR monitoring, suitability of primer/probe combinations and biomarker
 stability. Water Research 210, 117977. https://doi.org/10.1016/j.watres.2021.117977

INSACOG Dashboard. Retrieved January 30, 2023, from
https://inda.rcb.ac.in/insacog/statreportlineagegraph

Jahn, K., Dreifuss, D., Topolsky, I., Kull, A., Ganesanandamoorthy, P., 526 Fernandez-Cassi, X., Bänziger, C., Devaux, A.J., Stachler, E., Caduff, L., Cariti, F., 527 Corzón, A.T., Fuhrmann, L., Chen, C., Jablonski, K.P., Nadeau, S., Feldkamp, M., 528 Beisel, C., Aquino, C., Stadler, T., Ort, C., Kohn, T., Julian, T.R., Beerenwinkel, N., 529 2022. Early detection and surveillance of SARS-CoV-2 genomic variants in 530 531 wastewater using COJAC. Nature Microbiology 7, 1151-1160. https://doi.org/10.1038/s41564-022-01185-x 532

Joshi, M., Kumar, M., Srivastava, V., Kumar, D., Rathore, D.S., Pandit, R.,
Graham, D.W., Joshi, C.G., 2022. Genetic sequencing detected the SARS-CoV-2
delta variant in wastewater a month prior to the first COVID-19 case in Ahmedabad
(India). Environmental pollution (Barking, Essex □: 1987) 310, 119757.
https://doi.org/10.1016/j.envpol.2022.119757

Karthikeyan, Levy, Hoff, D., Humphrey, Birmingham, Jepsen, Farmer, Tubb,
Valles, Tribelhorn, Tsai, Aigner, Sathe, Moshiri, Henson, Mark, Hakim, Baer, Barber,
Belda-Ferre, Chacón, Cheung, Cresini, Eisner, Lastrella, Lawrence, Marotz, Ngo,
Ostrander, Plascencia, Salido, Seaver, Smoot, McDonald, Neuhard, Scioscia,
Satterlund, Simmons, Abelman, Brenner, Bruner, Buckley, Ellison, Gattas, Gonias,
Hale, Hawkins, Ikeda, Jhaveri, Johnson, Kellen, Kremer, Matthews, McLawhon,

544 Ouillet, Park, Pradenas, Reed, Riggs, Sanders, Sollenberger, Song, White, Winbush, Aceves, Anderson, Gangavarapu, Hufbauer, Kurzban, Lee, Matteson, Parker, Perkins, 545 546 Ramesh, Robles-Sikisaka, Schwab, Spencer, Wohl, Nicholson, McHardy, Dimmock, 547 Hobbs, Bakhtar, Harding, Mendoza, Bolze, Becker, Cirulli, Isaksson, Barrett, S., 548 Washington, Malone, Schafer, Gurfield, Stous, Fielding-Miller, Garfein, Gaines, 549 Anderson, Martin, Schooley, Austin, MacCannell, Kingsmore, Lee, Shah, McDonald, 550 Yu, Zeller, Fisch, Longhurst, Maysent, Pride, Khosla, Laurent, Yeo, Andersen, Knight, 2022. Wastewater sequencing reveals early cryptic SARS-CoV-2 variant 551 transmission. Nature 609, 101-108. https://doi.org/10.1038/s41586-022-05049-6 552

Karthikeyan, S., 2021. High-Throughput Wastewater SARS-CoV-2 Detection
Enables Forecasting of Community Infection Dynamics in San Diego County.
mSystems.

556 Khandia, R., Singhal, S., Alqahtani, T., Kamal, M.A., El-Shall, N.A., Nainu, F., Desingu, P.A., Dhama, K., 2022. Emergence of SARS-CoV-2 Omicron (B.1.1.529) 557 variant, salient features, high global health concerns and strategies to counter it amid 558 559 ongoing COVID-19 pandemic. Environmental Research 209, 112816. 560 https://doi.org/10.1016/j.envres.2022.112816

Kucirka, L.M., et al., 2020. Variation in transmission of SARS-CoV-2 in the
United States. Nature Medicine, 26(9), 1567-1573.

Kumar, M., Patel, A.K., Shah, A.V., Raval, J., Rajpara, N., Joshi, M., Joshi,
C.G., 2020. First proof of the capability of wastewater surveillance for COVID-19 in
India through detection of genetic material of SARS-CoV-2. The Science of the total
environment 746, 141326. https://doi.org/10.1016/j.scitotenv.2020.141326

Lamba, S., Ganesan, S., Daroch, N., Paul, K., Joshi, S.G., Sreenivas, D.,
Nataraj, A., Srikantaiah, V., Mishra, R., Ramakrishnan, U., Ishtiaq, F., 2023. SARSCoV-2 infection dynamics and genomic surveillance to detect variants in wastewater a longitudinal study in Bengaluru, India. The Lancet regional health. Southeast Asia
100151. https://doi.org/10.1016/j.lansea.2023.100151

Landgraff, C., Wang, L.Y.R., Buchanan, C., Wells, M., Schonfeld, J.,
Bessonov, K., Ali, J., Robert, E., Nadon, C., 2021. Metagenomic sequencing of
municipal wastewater provides a near-complete SARS-CoV-2 genome sequence

identified as the B.1.1.7 variant of concern from a Canadian municipality concurrentwith an outbreak. Cold Spring Harbor Laboratory.

Li, Durbin, 2010. Fast and accurate long-read alignment with Burrows–
Wheeler transform. Bioinformatics 26, 589–595.
https://doi.org/10.1093/bioinformatics/btp698

Li, H., Liu, S.-M., Yu, X.-H., Tang, S.-L., Tang, C.-K., 2020. Coronavirus disease 2019 (COVID-19): current status and future perspectives. International Journal of Antimicrobial Agents 55. https://doi.org/10.1016/j.ijantimicag.2020.105951

Li, Handsaker, Wysoker, Fennell, Ruan, Homer, Marth, Abecasis, Durbin,
2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–
2079. https://doi.org/10.1093/bioinformatics/btp352

Li, R., et al., 2020. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). Science, 368(6491), 489-493.

Maida, C.M., Amodio, E., Mazzucco, W., Rosa, G.L., Lucentini, L.,
Suffredini, E., Palermo, M., Andolina, G., Iaia, F.R., Merlo, F., Chiarelli, M.G.,
Siragusa, A., Vitale, F., Tramuto, F., SARI Collaboration Group, 2022. Wastewaterbased epidemiology for early warning of SARS-COV-2 circulation: A pilot study
conducted in Sicily, Italy. International Journal of Hygiene and Environmental Health
242. https://doi.org/10.1016/j.ijheh.2022.113948

Mercer, T.R., Salit, M., 2021. Testing at scale during the COVID-19 pandemic.
Nature Reviews. Genetics 22. https://doi.org/10.1038/s41576-021-00360-w

Morvan, Jacomo, Souque, Wade, Hoffmann, Pouwels, Lilley, Singer, Porter,
Evens, Walker, Bunce, Engeli, Grimsley, O'Reilly, Danon, 2022. An analysis of 45
large-scale wastewater sites in England to estimate SARS-CoV-2 community
prevalence. Nature Communications 13, 1–9. https://doi.org/10.1038/s41467-02231753-y

Nyberg, T., Ferguson, N.M., Nash, S.G., Webster, H.H., Flaxman, S.,
Andrews, N., Hinsley, W., Bernal, J.L., Kall, M., Bhatt, S., Blomquist, P., Zaidi, A.,
Volz, E., Aziz, N.A., Harman, K., Funk, S., Abbott, S., Nyberg, T., Ferguson, N.M.,

605 Nash, S.G., Webster, H.H., Flaxman, S., Andrews, N., Hinsley, W., Bernal, J.L., Kall, M., Bhatt, S., Blomquist, P., Zaidi, A., Volz, E., Aziz, N.A., Harman, K., Funk, S., 606 607 Abbott, S., Hope, R., Charlett, A., Chand, M., Ghani, A.C., Seaman, S.R., Dabrera, G., Angelis, D.D., Presanis, A.M., Thelwall, S., Hope, R., Charlett, A., Chand, M., Ghani, 608 A.C., Seaman, S.R., Dabrera, G., Angelis, D.D., Presanis, A.M., Thelwall, S., 2022. 609 610 Comparative analysis of the risks of hospitalisation and death associated with SARS-611 CoV-2 omicron (B.1.1.529) and delta (B.1.617.2) variants in England: a cohort study. The Lancet 399, 1303–1312. https://doi.org/10.1016/S0140-6736(22)00462-7 612 Peccia, J., Zulli, A., Brackney, D.E., Grubaugh, N.D., Kaplan, E.H., 613 614 Casanovas-Massana, A., Ko, A.I., Malik, A.A., Wang, D., Wang, M., Warren, J.L., 615 Weinberger, D.M., Arnold, W., Omer, S.B., 2020b. Measurement of SARS-CoV-2 616 RNA in wastewater tracks community infection dynamics. Nature biotechnology 38, 1164–1167. https://doi.org/10.1038/s41587-020-0684-z 617 Polo, D., Quintela-Baluja, M., Corbishley, A., Jones, D.L., Singer, A.C., 618 Graham, D.W., Romalde, J.L., 2020. Making waves: Wastewater-based epidemiology 619 620 for COVID-19 – approaches and challenges for surveillance and prediction. Water 621 Research 186, 116404. https://doi.org/10.1016/j.watres.2020.116404

Prado, T., Fumian, T.M., Mannarino, C.F., Resende, P.C., Motta, F.C.,
Eppinghaus, A.L.F., Chagas do Vale, V.H., Braz, R.M.S., de Andrade, J. da S.R.,
Maranhão, A.G., Miagostovich, M.P., 2021. Wastewater-based epidemiology as a
useful tool to track SARS-CoV-2 and support public health policies at municipal level
in Brazil. Water research 191, 116810. https://doi.org/10.1016/j.watres.2021.116810

Prakash, S., Shukla, S., Mishra, H., Prakash, O., Khan, D.N., Pandey, A., 627 Reddy, D.H., Jain, A., 2021. SARS-CoV-2 -RNA persists longer in faecal sample as 628 compared to nasal and throat swab samples of COVID-19 patients'; an observational 629 of Medical 39. 630 study. Indian Journal Microbiology 631 https://doi.org/10.1016/j.ijmmb.2020.10.012

Rambaut, A., et al., 2020. The proximal origin of SARS-CoV-2. Nature
Medicine, 26(4), 450-452.

Randazzo, W., Truchado, P., Cuevas-Ferrando, E., Simón, P., Allende, A.,
Sánchez, G., 2020. SARS-CoV-2 RNA in wastewater anticipated COVID-19

occurrence in a low prevalence area. Water Research 181, 115942.
https://doi.org/10.1016/j.watres.2020.115942

Ranjan, R., 2022. Omicron Impact in India: Analysis of the Ongoing
COVID-19 Third Wave Based on Global Data. Cold Spring Harbor Laboratory.

Rosa, L., Libera, D., Iaconelli, Ciccaglione, Bruni, Taffon, Equestre, Alfonsi, 640 641 Rizzo, Tosti, Chironna, Romanò, Zanetti, Muscillo, 2014. Surveillance of hepatitis A virus in urban sewages and comparison with cases notified in the course of an 642 643 outbreak, Italy 2013. BMC Infectious Diseases 14, 1–11. 644 https://doi.org/10.1186/1471-2334-14-419

Shang, Y., Li, H., Zhang, R., 2021. Effects of Pandemic Outbreak on
Economies: Evidence From Business History Context. Frontiers in Public Health 9.
https://doi.org/10.3389/fpubh.2021.632043

Sharifi, A., Khavarian-Garmsir, A.R., 2020. The COVID-19 pandemic:
Impacts on cities and major lessons for urban planning, design, and management. The
Science of the Total Environment 749.
https://doi.org/10.1016/j.scitotenv.2020.142391

Shrestha, L.B., Tedla, N., Bull, R.A., 2021. Broadly-Neutralizing Antibodies
Against Emerging SARS-CoV-2 Variants. Frontiers in Immunology 12.
https://doi.org/10.3389/fimmu.2021.752003

Srivastava, V., Gupta, S., Patel, A.K., Joshi, M., Kumar, M., 2021.
Reflections of COVID-19 cases in the wastewater loading of SARS-CoV-2 RNA: A
case of three major cities of Gujarat, India. Case Studies in Chemical and
Environmental Engineering 4. https://doi.org/10.1016/j.cscee.2021.100115

Stephens, N., Béen, F., Savic, D., 2022. An Analysis of SARS-CoV-2 in
Wastewater to Evaluate the Effectiveness of Nonpharmaceutical Interventions against
COVID-19 in The Netherlands. ACS ESand#38;amp;T Water 2, 2158–2166.
https://doi.org/10.1021/acsestwater.2c00071

Tiwari, A., Lipponen, A., Hokajärvi, A.-M., Luomala, O., Sarekoski, A.,
Rytkönen, A., Österlund, P., Al-Hello, H., Juutinen, A., Miettinen, I.T., SavolainenKopra, C., Pitkänen, T., 2022b. Detection and quantification of SARS-CoV-2 RNA in

666	wastewater influent in relation to reported COVID-19 incidence in Finland. Water
667	Research 215, 118220. https://doi.org/10.1016/j.watres.2022.118220
668	Tracking SARS-CoV-2 variants [WWW Document], n.d. URL
669	https://www.who.int/activities/tracking-SARS-CoV-2-variants (accessed 2.10.23).
670	Valieris, Drummond, Defelicibus, Dias-Neto, Rosales, Silva, T. da, 2022. A
671	mixture model for determining SARS-Cov-2 variant composition in pooled samples.
672	Bioinformatics 38, 1809–1815. https://doi.org/10.1093/bioinformatics/btac047
673	Valieris, Drummond, Defelicibus, Dias-Neto, Rosales, Silva, T. da, 2022b. A
674	mixture model for determining SARS-Cov-2 variant composition in pooled samples.
675	Bioinformatics 38, 1809–1815. https://doi.org/10.1093/bioinformatics/btac047
676	van der Meer, L.M., et al., 2020. Wastewater-based surveillance for SARS-
677	CoV-2: The way forward. Journal of Virological Methods, 282, 113907.
678	Wilton, T., 2021. Rapid Increase of SARS-CoV-2 Variant B.1.1.7 Detected
679	in Sewage Samples from England between October 2020 and January 2021.
680	mSystems.
681	Yadav, P.D., Kumar, S., 2022. Global emergence of SARS-CoV-2 variants:
682	new foresight needed for improved vaccine efficacy. The Lancet Infectious Diseases
683	22, 298–299. https://doi.org/10.1016/s1473-3099(21)00687-3
684	Zhang, L., 2022. Broadly neutralizing antibodies against SARS-CoV-2
685	variants. National Science Open 1, 20220005. https://doi.org/10.1360/nso/20220005
686	
687	

- Fig.1. Geographical locations of the sewage treatment facilities sampled in Pune city(India).
- 690 Fig.2 Temporal trends of SARS-CoV-2 viral load in relation to COVID-19 cases and
- the relative abundance of SARS-CoV-2 variants in sewer shed sites (Bhairobanala,
- 692 Bopodi, Erandwane, Kharadi, Baner).
- 693 Fig.3 Temporal trends of SARS-CoV-2 viral load in relation to COVID-19 cases and
- the relative abundance of SARS-CoV-2 variants in sewer shed sites (New Naidu, Old
- 695 Naidu, Tanajiwadi, Vitthalwadi, Mundhwa).
- **Fig.4** Relative abundance of SARS-CoV-2 lineages in wastewater according to date of collection and sewer shed sites in Pune.
- 698 Fig.5 Temporal trends of dominant variants of SARS-CoV-2 in wastewater samples
- between November 2021 and April 2022.



Sewage Treatment Plant : PUNE-INDIA

Erandwane



Bhairoba Nala















90 MLD



Viral Load copies/ml (log scale)

Viral Load copies/ml (log scale)





Viral Load copies/ml (log scale)

Viral Load copies/ml (log scale)





X 75 38 38 12 10









Viral Load copies/ml (log scale)

BA.5 BA.4 BA.3 BA.2.X BA.2.38 BA.2.38 BA.2.38 BA.2.10 BA.2.10 BA.1.X BA.1 ron ron B.1. B.1. B.1. B.1. B.1



medRxiv preprint doi: https://doi.org (which was not certified by per



Viral Load copies/ml (log scale)

18-Feb-2022 -

16-Feb-2022 -

11-Feb-2022 -

6-Feb-2022 -

4-Feb-2022 -

2-Feb-2022

28-Jan-2022

26-Jan-2022

- 2202-ns[-12

19-Jan-2022

14-Jan-2022

12-Jan-2022

7-Jan-2022

5-Jan-2022

31-Dec-2021

29-Dec-2021

54-Dec-2021 -

55-Dec-2021

17-Dec-2021

12-Dec-2021

8-Dec-2021

1-Dec-2021

24-Nov-2021

17-Nov-2021

10-Nov-2021



Viral Load copies/ml (log scale)

29-Apr-2022 + --26-Apr-2022 22-Apr-2022 -19-Apr-2022 - -15-Apr-2022 12-Apr-2022 8-Apr-2022 -5-Apr-2022 1-Apr-2022 29-Mar-2022 25-Mar-2022 22-Mar-2022 15-Mar-2022 11-Mar-2022 8-Mar-2022 -4-Mar-2022 -1-Mar-2022 -52-Eeb-2022 22-Feb-2022 18-Feb-2022 -12-Eeb-2022 11-Feb-2022 🕂 8-Feb-2022 4-Feb-2022 1-Feb-2022 ₹ 2202-nsl-82 25-Jan-2022 21-Jan-2022 🕂 18-Jan-2022 14-Jan-2022 🗄 11-Jan-2022 7-Jan-2022 4-Jan-2022 -31-Dec-2021 -28-Dec-2021 54-Dec-2021 -51-Dec-2021 17-Dec-2021 14-Dec-2021 10-Dec-2021 + 3-Dec-2021 26-Nov-2021 19-Nov-2021 12-Nov-2021

Viral Load copies/ml (log scale)

Viral Load copies/ml (log scale)

BA. BA. BA. BA. BA. BA.

Viral Load copies/ml (log scale)

- Zeta
- XBB_Rec
- Other
- Omicron_BA.5
- Omicron BA.4
- Omicron_BA.3
- Omicron_BA.2.X
- Omicron_BA.2.75
- Omicron_BA.2.38
- Omicron BA.2.12
- Omicron BA.2.10
- Omicron BA.2
- Omicron BA.1.X
- Omicron_BA.1
- Omicron
- Kappa_B.1.617.1
- Iota_B.1.526
- Gamma_P.1
- Eta B.1.525
- Delta_B.1.617.2
- Beta_B.1.351
- Alpha_B.1.1.7

Month/Year

Month/Year

Month/Year

Month/Year

Month/Year

Month/Year

Month/Year

Month/Year

Month/Year

Month/Year

Month/Year

