




N =																	
JANSSEN	21	56	58	56	55	57	56	58	49	40	32	26	23				
MODERNA	133	160	177	177	168	144	105	103	69	48	42	34	22				
PFIZER		72	115	108	105	108	90	79	54	37	22	17	18	13			

**Supplemental Figure 1. Anti-spike IgG titers vs months after date of full immunization, comparing by vaccine type, among patients with prior COVID-19 in the initial series cohort**

Of note, the boxplots shown here are bounded by the upper and lower limits of the assay used. The dots represent the outliers, defined as greater than 1.5\*IQR above the third quartile or less than 1.5\*IQR below the first quartile.

The tables of N show the number of titers for each month, by vaccine type.



N =							
MODERNA	115	103	104	105	90	71	68
PFIZER	95	91	93	80	67	38	47

**Supplemental Figure 2. Anti-spike IgG titers vs months after date of full immunization by third vaccine dose, comparing by vaccine type, among patients with prior COVID-19 in the third dose cohort**

Of note, the boxplots shown here are bounded by the upper and lower limits of the assay used. The dots represent the outliers, defined as greater than  $1.5 \times \text{IQR}$  above the third quartile or less than  $1.5 \times \text{IQR}$  below the first quartile.

The tables of N show the number of titers for each month, by vaccine type.