Teledentistry for improving access to, and quality of oral health care: A protocol for an overview of systematic reviews and meta-analyses 4

5	Pascaline Kengne Talla ^{1*} , Paul Allison ¹ , André Bussières ^{2,3} , Nicolas Giraudeau ⁴ , Svetlana,
6	Komarova ¹ , Quentin Basiren ¹ , Frédéric Bergeron ⁵ , Elham Emami ¹
7	
8	
9	1. Faculty of Dental Medicine and Oral Health Sciences, McGill University, Montreal, Quebec,
10	Canada
11	2. Université de Québec à Trois rivières, Departement de Chiropratique,
12	3. School of Physical and Occupational Therapy, McGill University, Montreal, Quebec, Canada
13	4. Faculté d'Odontologie, Université de Montpellier, Montpellier, France
14	5. Laval University, Department of preventive and social medicine, Quebec City, Quebec, Canada
15	*Corresponding author
16	Email: <u>pascaline.kengnetalla@mcgill.ca (PKT)</u>
17	
18	
19	
20	
21	

22 Abstract

23 Digital technologies are becoming essential to address and optimize the suboptimal performance of healthcare systems. Teledentistry involves the use of information and communication technology to 24 improve access to oral health care and the quality of oral health care delivery. Several systematic 25 26 reviews (SRs) have been conducted to synthesize evidence on the effectiveness of teledentistry but 27 with conflicting results. The aim of this review is to comprehensively summarize available SRs and 28 provide evidence on the impact of teledentistry on access to oral care, patients' and oral healthcare 29 providers' outcomes, quality of oral health care and costs. This protocol has been registered with the International Prospective Register of Systematic Reviews (PROSPERO CRD42022373964). Six 30 31 electronic databases including MEDLINE (Ovid), Embase (Embase.com), CINAHL (EBSCO), 32 Web of Science, Cochrane Library and Epistemonikos will be searched for SRs of quantitative, qualitative, and mixed reviews evaluating teledentistry modalities involving both patients and/or 33 34 oral health care providers (OHCPs). We will include only studies published in English or French. 35 The primary outcomes will be considered from the patients' perspective (e.g., access to oral health care, patient-reported outcomes, and experiences). The secondary outcomes will include outcomes 36 37 from patients and OHCPs (e.g., clinical outcomes, safety, behaviors, and costs). Two independent 38 reviewers will perform data screening, data extraction and will assess the quality of included studies 39 using the AMSTAR 2 and ROBIS tools. Data will be synthesized narratively and presented by 40 tables and graphs. We will report any overlap of primary studies in the SRs. A statement on the 41 strength of evidence for each outcome will be provided if possible. This review will inform decision-makers, patients, OHCPs, and researchers on the potential effectiveness, benefits, and 42 43 challenges of teledentistry and support them in making recommendations for its use. Results will be disseminated through peer-reviewed publications, presentations at conferences, and on social 44 45 media.

48 Introduction

49 Oral diseases globally affect more than 3.5 billion people, highlighting the need for interventions 50 that could improve accessibility to and affordability of oral health care [1]. Information and 51 communication technologies (ICTs) are promising approaches to address some of the inadequacies 52 of healthcare systems, to improve patients' access to and experiences of care, to reduce the costs of 53 care delivery and to promote high value care [2-8]. In addition, digital technologies can improve the 54 quality of health care (e.g equity, safety, effectiveness, patient-centered and timely care) [9, 10]. 55 The COVID-19 pandemic has accelerated the use of these technologies in all health-related 56 disciplines including dentistry [11], paving the way for the development of virtual dental care. They 57 have reshaped the delivery of oral health care including patient-clinician interaction, screening, 58 diagnosis of oral diseases, monitoring of patients, treatment planning and management of care [12].

59

60 Emerging evidence suggests that teledentistry, a branch of telehealth, is cost-effective at the micro, meso and macro levels involving patients, oral health care providers (OHCPs) and allied health care 61 62 workers, and decision makers [13-16]. Teledentistry includes synchronous and asynchronous 63 modes, and uses different vehicles such as telephones, smartphones, tablets and computers, as well 64 as various approaches such as calls, text and voice messaging, videos, and applications [17, 18]. 65 Teledentistry has been used in the context of dental education, training, and transfer of information 66 or for the delivery of dental care/services. It involves the interaction between OHCPs and their peers, or with other health care providers or with their patients and/or caregivers to improve 67 68 patients' outcomes and the quality of care.

69

Multiple reviews have synthesized the evidence on the benefits and implementation challenges of teledentistry [19-22], the process of teledentistry, and the outcomes and experiences [13, 14, 23-25] from the perspective of patients, healthcare organizations and OHCPs [26]. However, the results are often conflicting [22, 27, 28], inconsistent or inconclusive [21, 29, 30]. Although systematic

74 reviews are a compelling means of synthesizing research, a systematic review of existing systematic reviews (SRs) can provide a broader assessment of the quality and credibility of available evidence 75 76 [31, 32], and offer valuable information for patients, families, health professionals, researchers, and 77 policy-makers. The information generated by such overview can be used to enhance both clinical 78 practice and population health. In this study, we will use the term 'overview' due to the lack of 79 consistency in the literature on the terminology of the compilation data from multiple SRs to 80 provide a single summary of relevant evidence [33, 34]. A published overview on teledentistry 81 evaluated its accuracy and effectiveness for the delivery of oral health care [35]. A major limitation 82 of this overview is the lack of risk of bias assessments of the included SRs. Moreover, it has only 83 focused on accuracy of screening, diagnosis, and therapeutic management of dental care outcomes. 84 Other health outcomes related to access to oral care (e.g., utilization of services) and patients' and OHCPs' behaviors to improve the quality of oral healthcare would be important to consider to 85 inform practice, policy decision-making, and future research [36]. Another available overview is 86 87 limited to tele-orthodontics to improve compliance in orthodontic patients [37]. Therefore, there is a 88 need to address the gaps in literature by conducting a comprehensive overview of existing SRs 89 using a rigorous methodology with valid quality assessment of the evidence [38]. This proposed 90 overview of existing SRs aims to compile and contrast the existing evidence from systematic 91 reviews published on teledentistry [31]. Accurate information resulting from this overview will 92 assist decision-makers on the effectiveness of teledentistry and inform the development of 93 guidelines to support OHCPs in its implementation.

94

95 **Research question**

96 We will answer to the following research question: "From the perspective of a range of 97 stakeholders, to what extent is teledentistry effective in improving access to, and quality of oral 98 healthcare, while reducing related costs?"

99 Methods

100	The review is guided by the Joanna Briggs Institute (JBI) Manual for Evidence synthesis for
101	umbrella reviews [39], and the Cochrane handbook for overview of reviews [40], and is written in
102	accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols
103	(PRISMA-P) guidance [41]. The PRISMA checklist is available as S1 checklist.
104	

105 Study registration

106 This overview has been registered in the International Prospective Register of Systematic Reviews107 (PROSPERO CRD42022373964).

108

109 Eligibility criteria

We will use the "PICOSS" format: Participants, Intervention, Comparator, Outcome, Study designand Setting.

112

113 **Participants/Population**

We will include SRs involving patients receiving oral healthcare services performed by any licensed OHCPs (general dental practitioners, dental specialists, dental hygienists, dental nurses/assistants, and dental therapists) and the SRs involving OHCPs with or without the patients.

117

118 Intervention

Teledentistry refers to use of information and communication technologies including the transmission of clinical information and images between an oral health professional and patient or between two health professionals, including at least one oral health professional, who are separated by distance for dental consultations, diagnosis and treatment planning [18]. Teledentistry is a modality used to provide remote access to healthcare services to patients. It includes the use of a group of technologies and modalities, which can be categorized as follows [17]: i) Store and

125 forward, which is used to keep patients' oral health records such as radiographs and photographs; ii) 126 Remote patient monitoring, used for patient data collection from a remote site and then transferring 127 them to a dental practitioner in another location; iii) Live video, involving the use of a real-time 128 interaction between a patient and an OHCP, using audio-visual communication for screening, diagnosis, treatment planning or follow-up; and iv) Mobile health, making use of mobile 129 130 communication devices such as phones and tablets to provide virtual oral healthcare services. Other 131 terms referring to teledentistry encompass e-health, virtual care, telemedicine oral health and 132 mobile oral health. Irrespective of the term used, we will include all interventions that involve oral 133 health care delivery through telecommunication systems in the presence of at least one member of 134 dental staff.

135

136 **Comparator**

We will include any SRs where the comparator could be one of the following types of interventions or cases: in-person or face-to face interventions or usual care; no intervention; synchronous versus asynchronous; synchronous versus mobile health; synchronous versus remote monitoring; and other digital technologies (e.g. electronic dental records, virtual reality).

141

142 **Outcomes**

143 The primary outcomes are reported from the patients' perspective. They will include access to oral 144 health care (e.g. use of oral healthcare services, number of consultations, use in emergency cases, 145 delay of treatment, waiting time); patient-reported outcomes (e.g. oral health related- and overall quality of life; self-reported clinical outcomes; pain management, oral functions; psychosocial 146 147 impact) and experiences with oral health care (e.g. satisfaction with care; communication with 148 OHCPs, patient-centered care and empowerment; acceptance and understanding of information and confidence in the treatment, and experience with the technology such as ability to use the 149 150 application).

151

152 Secondary outcomes will be reported as they relate to:

 Patient indicators: Clinical outcomes reported by OHCPs (e.g. plaque index, gingival index and white spot lesions); Adherence/compliance to treatment (e.g. medication, oral health prevention and promotion practices); Knowledge, attitude, and behavior; Barriers and enablers towards the use of teledentistry; Safety; Adverse outcomes; and Costs (e.g. travel time, transportation, missing work/school, loss of the productivity and consultation time).

OHCP indicators: Accuracy of diagnosis; Awareness, knowledge, attitude, and behavior;
 Barriers and enablers towards the adoption of teledentistry; Prescribing (e.g. testing,
 medication); Monitoring of patients; Coordination and management of oral health care;
 Communication with patients, dental specialists, dental staff and other health professionals;
 Costs (e.g. equipment, number of patients per day, number of consultations per day, waiting
 time, training in the use of the equipment for teledentistry); and Equity.

164

165 Study designs

166 We will include systematic reviews with or without meta-analysis of quantitative (randomized 167 control trials, quasi-experimental studies such as non-randomized controlled trials, before and after 168 controlled studies and interrupted series studies, or observational studies), qualitative and mixed 169 methods studies using any teledentistry modalities (asynchronous, remote monitoring, real-time, 170 and mhealth). The term SR refers to a reproducible, standardized and transparent approach aiming 171 to identify, evaluate and summarize the evidence from primary studies on a particular topic, thereby making it more accessible to decision-makers [42]. We will include only SRs that have conducted 172 173 searches in at least two databases, have clear inclusion/exclusion criteria, performed quality 174 assessment, and synthesized included studies [32, 42, 43].

175

176 Setting

We will capture the evidence from SRs on teledentistry conducted in any dental setting (e.g., dental
offices, school, community, hospital, home care), geographical region (e.g., rural, urban), and
country (low-, middle- or high income).

180

181 Exclusion criteria

We will exclude any types of knowledge synthesis systematic reviews lacking a formal methodological quality or risk of bias assessment or where a search was conducted in a single database (26). We will exclude studies on teledentistry focusing only on education and training in dentistry and research without a care delivery component.

186

187 Search strategy

188 We will search electronic databases in MEDLINE (Ovid), Embase, CINAHL (EBSCO), Web of 189 Science, The Cochrane Library and Epistemonikos (https://www.epistemonikos.org/) using a search 190 strategy that has been developed using an interactive process by members of the research team with 191 the support of an expert librarian. The search strategy will be conducted from database inception 192 using the following keywords: ("teledentistry" OR "remote care*" OR "mobile health") AND 193 ("reviews" OR "meta-analysis" OR "systematic review). We will also perform a search in 194 Sociological Abstract (Proquest), Academic Search Premier (EBSCO), and Proquest Dissertations 195 & Theses. We will contact experts in the field by email if additional data are required. In addition, 196 we will check the reference lists of included SRs and any identified overviews for any eligible articles. See S1 file for the search strategy conducted in Medline. We will update the search prior 197 198 to the publication of the review to identify any new relevant systematic reviews. There will be no 199 restrictions for countries, age of participants, publication date and settings. However, we will 200 consider only studies published in English or French because of the limited resources for 201 translation.

202

203 Study selection

204 SRs identified by the search will be imported into Covidence software [44]. The research team will discuss the selection criteria to ensure a shared understanding before pairs of assessors will 205 206 independently pilot the initial screening phase (i.e., titles and abstracts) on 10% of total number of 207 citations retrieved. Disagreements at each stage will be resolved through discussion or consultation 208 with a third reviewer whenever necessary. We will adjust the grid following this calibration 209 exercise. Reviewers will consider "include", "exclude", or "Maybe" as modalities of the studies 210 selection. All studies rated "include" or "unclear" will be considered for the second phase (i.e. full 211 text review). The decisions and reasons for exclusion will be recorded in Covidence software.

212

213 Data extraction

Two research team members will independently perform data extraction using a form in Excel that is based on the JBI data extraction form for review of systematic reviews[39]. The data extraction form is provided as S2 file. We will conduct a pilot data extraction on 10% of eligible SRs until a consensus on extracted data is reached between reviewers. Any discrepancies will be resolved through discussion, or by consultation with a third reviewer. We will not extract data from the primary studies included in the reviews, nor re-synthetize their findings. Extracted information will include:

- Review characteristics: first author, year of publication, country, type of review with or
 without meta-analysis; aim of the study; publication language; published protocol; number
 of databases searched, range date of search strategy; any restrictions (e.g. language,
 geographic or date), inclusion/exclusion criteria;
- 225 2. Participants: number and profile of participants (e.g patients, OHCPs);

- 3. Intervention and comparators: intervention(s) of interest and comparators; domains in
 dentistry; reported definition of intervention from authors;
- 4. Outcomes: primary outcomes (access to oral health care, patient -reported outcomes, patient reported-experiences); secondary outcomes relevant to patients and OHCPs; effect size metric(s) reported (e.g. risk ratio) for categorical outcomes and (e.g., standard mean differences) for continuous outcomes;
- 5. Setting: type of settings likely community, dental office, university, hospital;
- 233 6. Methods: type of study designs included (e.g. randomized controlled trials, observational 234 studies or both); number of included studies, number of studies reporting data for meta-235 analyses, use of theoretical framework in the intervention or for data analysis; main primary 236 outcomes of interest; risk of bias tools; statistical methods used to combine studies; 237 synthesis and summary of data, estimates including heterogeneity measures, and any 238 additional analyses (e.g. subgroup analysis or sensitivity analysis); the level of evidence, for instance the Grading of Recommendations, Assessment, Development and Evaluation 239 240 (GRADE) from included studies;
- 7. Other details: main conclusions, limitations, next steps, funding source and conflicts ofinterest.
- If we have missing data, we will search for additional information such as protocols or contactstudy authors to find the available information.
- 245

246 Quality assessment of reviews

The quality assessment of SRs is an essential component when we conduct an overview of systematic reviews [32]. It includes both the methodological quality and the risk of bias which are distinct concepts [45, 46]. The quality assessment examines compliance with the highest possible standards in conducting and reporting their research process [47, 48], while risk of bias assessment

(the appraisal of internal validity) examines any concerns with the design, conduct, analysis,interpretation, or reporting of a study which could affect the study's results [49].

253

254 To conduct the quality assessment, we will use two valid and reliable tools covering complementary 255 criteria: i) the AMSTAR-2 checklist (the Assessment of Multiple Systematic Reviews 2) to evaluate 256 the methodological quality [50]; and ii) the ROBIS tool to assess comprehensively the risk of bias 257 [51]. AMSTAR-2 is applicable to systematic reviews of both randomized controlled trials (RCTs) 258 and non-randomized controlled trials (non-RCTs). This tool consists of 16 items, with 7 critical 259 domains (items 2, 4, 7, 9, 11, 13, and 15). The assessment comprises three options, "yes" 260 (item/question fully addressed), "no" (item/question not addressed), or "partial" (item/question not 261 fully addressed). The overall confidence on the results of the review is classified into high, 262 moderate, low, and critically low [50].

263

264 The ROBIS tool aims to assess the risk of bias in reviews related to interventions, etiology, 265 diagnosis and prognosis [51]. It consists of 24 questions across three phases: Phase 1 assesses the 266 relevance (optional); phase 2 identifies concerns with the review process; and phase 3 judges the 267 risk of bias in the review. Phase 2 and 3 questions are answered with the following options: yes: 268 probably ves; probably no; no; or no information. The concerns regarding phase 2 and phase 3 269 domains are classified as high, low, or unclear. We will focus only on phases 2 and 3 to give an 270 overall score on the risk of bias in each review. Phase 2 identifies the concerns in the review across 271 four domains: study eligibility criteria; identification and selection of studies; data collection and 272 study appraisal; and synthesis and findings. There are signaling questions and a judgment of 273 concerns about risk of bias for each domain (low, high or unclear). Phase 3 summarizes the 274 concerns identified during the Phase 2 assessment [51]. Finally, we will provide a judgment 275 regarding the overall risk of bias.

Despite the relevance of these tools and the growing number of reporting guidelines, they are mostly for SRs of quantitative reviews and there is a lack of critical appraisal tools [52] to assess the quality of SRs of qualitative studies [53]. In fact, some criteria from AMSTAR-2 and ROBIS tools are not adapted for quality assessment of factors such as risk of bias, publication bias, heterogeneity, meta-analysis in SRs of qualitative research [53]. However, we will use both these tools to conduct the quality assessment within the limit of their utilization.

283

Two reviewers will independently assess each review using both tools. Any discrepancies will be resolved by discussion or a consultation with a third reviewer. We will not perform any quality assessment of primary studies in included reviews.

287

288 Data synthesis

289 We will perform a narrative synthesis of the data. We will present results in tabular form in tables 290 describing characteristics of included studies (e.g. first author's name, language of publication, 291 country, settings, year of publication, profile of participants, study purpose), information on 292 teledentistry (e.g. definition, teledentistry modalities), methods (e.g. SR with or without meta-293 analysis, type of analysis), additional results (e.g. assessment of quality, appraisal tool used, 294 heterogeneity of the results, and level of evidence), and outcomes (e.g. primary outcomes and 295 secondary outcomes from included studies). We will categorize the SRs into sub-groups, according 296 to the type of intervention such as teledentistry modalities. If possible, we will perform a narrative 297 synthesis of the subgroups.

298

An important concern in conducting an overview is the likelihood of overlap in primary studies across included reviews, which may result in overestimates in results (36) and confuse clinicians making decisions amongst competing interventions in their clinical practices. Thus, we will

302 summarize key details from the studies included and will not perform a meta-synthesis of included

303 meta-analyses. We will report any overlap between SRs in the tables as a matrix.

304

We will report the overall score of the two measurement tools (AMSTAR-2 and ROBIS), the level of evidence from the credibility assessment, and the percentage overlap between primary studies within included SRs and SR-meta- analysis (SR-MAs).

308

309 We will assess the certainty of evidence defined as any of evaluation of the totality or strength of 310 the evidence on the impact of teledentistry using the following criteria for credibility assessment as 311 proposed in four categories [38, 54]: Class I (Convincing evidence): associations with a statistical 312 significance of P-value $< 10^{-6}$, include more than 1000 cases (or more than 20 000 participants for 313 continuous outcomes), have the largest component study reporting a significant result (P < 0.05), 314 have a 95% prediction interval that excludes the null, does not have large heterogeneity ($I^2 < 50\%$). 315 and shows no evidence of small study effects (P > 0.10) and excess significance bias (P > 0.10); 316 Class II (Highly suggestive evidence): associations with a significance of P < 0.001, include more 317 than 1000 cases (or more than 20 000 participants for continuous outcomes), and have the largest 318 component study reporting a statistically significant result (P < 0.05); Class III (Suggestive 319 evidence): associations that report a significance of P < 0.01 with more than 1000 cases (or more 320 than 20000 participants for continuous outcomes); Class IV (Weak evidence): remaining significant 321 associations with P < 0.05. If we do not have sufficient data, we will analyse the certainty of 322 evidence using data reported in the included SRs.

323

324 **Discussion**

This review will contribute to a comprehensive body of knowledge on the potential effectiveness ofteledentistry. To our knowledge, this overview is the first concerning teledentistry that will include

327 a broad definition of teledentistry incorporating various modalities, participants, study designs, 328 types of SRs and a combination of evidence, thereby increasing the understanding of the potential 329 of teledentistry. This overview has implications for dental practice, policy, education and research. 330 Beyond using a broad definition of teledentistry, this review will provide valuable knowledge on 331 the use of digital technologies, including teledentistry, to improve oral health care systems, 332 Teledentistry can be used to enhance the quality of oral health care as well as enhance oral health 333 equity. The lack of high-quality information on the effectiveness of teledentistry has often been 334 reported as a strong barrier to its implementation. This review has the potential to close that gap and 335 will contribute to inform decision-makers, researchers, clinicians, and patients on the effectiveness 336 of teledentistry for the delivery of care and to improve patients' outcomes and experiences, as well 337 as inform the needs for the future research. For instance, results will highlight teledentistry's 338 benefits and challenges, thereby identifying where is could potentially be used. These findings may 339 also support the development and implementation of guidelines on teledentistry in clinical practice.

340

341 We anticipate some limitations. A first limitation relates to the overlap between the primary studies 342 included in more than one SR in our overview, which can cause an overestimation of the effects for 343 a given outcome. We will report the percentage of overlap between the included studies and will 344 discuss of the impact of the overlap on our results. The second limitation is the restriction of our 345 overview to SRs only, which could result in missing some relevant primary studies published 346 during the completion of our review. To mitigate this issue, we will repeat our search just before we 347 finish our manuscript and will identify any additional relevant published SRs, but we will only 348 include them if they fulfill our inclusion criteria. We will document and report any deviations to the 349 protocol during the review. In addition, we will mention any limitation in the lack of available 350 critical appraisal tool for any designs (quantitative, qualitative and mixed reviews) [53].

352 Dissemination plan

Results of this overview will be disseminated through presentations at national and international
scientific and professional conferences, publication in peer-review journals and social media, such
as Twitter, plus the website of the Faculty of Dental Medicine and Oral health Sciences, at McGill
University.

357

358 Acknowledgement

359 The authors thank the summer students for their involvement in the first steps of data screening.

360

361 Supporting information

362 S1 checklist: PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis

363 Protocols) 2015 checklist: Recommended items to address in a systematic review protocol.

- 364 S1 file: Search strategy
- 365 S2 file: Data extraction sheet

366

367 Funding

368 None

369 **Competing interests**

370 None

371 Availability of data and materials

372 Not applicable

374 **References**

Peres MA, Macpherson LMD, Weyant RJ, Daly B, Venturelli R, Mathur MR, et al. Oral
 diseases: a global public health challenge. The Lancet. 2019;394(10194):249-60.

Gandhi S, Chen S, Hong L, Sun K, Gong E, Li C, et al. Effect of Mobile Health
 Interventions on the Secondary Prevention of Cardiovascular Disease: Systematic Review and
 Meta-analysis. The Canadian journal of cardiology. 2017;33(2):219-31.

380 3. Fjeldsoe BS, Marshall AL, Miller YD. Behavior change interventions delivered by mobile
381 telephone short-message service. American journal of preventive medicine. 2009;36(2):165-73.

382 4. Abaza H, Marschollek M. mHealth Application Areas and Technology Combinations*. A
383 Comparison of Literature from High and Low/Middle Income Countries. Methods of information in
384 medicine. 2017;56(7):e105-e22.

5. Toniazzo MP, Nodari D, Muniz F, Weidlich P. Effect of mHealth in improving oral
hygiene: A systematic review with meta-analysis. Journal of clinical periodontology.
2019;46(3):297-309.

388 6. Irving M, Stewart R, Spallek H, Blinkhorn A. Using teledentistry in clinical practice as an
and telecare. 2018;24(3):129-46.

391 7. Aquilanti L, Santarelli A, Mascitti M, Procaccini M, Rappelli G. Dental Care Access and the
392 Elderly: What Is the Role of Teledentistry? A Systematic Review. International journal of
393 environmental research and public health. 2020;17(23).

Bashshur RL, Shannon GW, Krupinski EA, Grigsby J, Kvedar JC, Weinstein RS, et al.
 National telemedicine initiatives: essential to healthcare reform. Telemedicine journal and e-health :
 the official journal of the American Telemedicine Association. 2009;15(6):600-10.

397 9. Ibrahim MS, Mohamed Yusoff H, Abu Bakar YI, Thwe Aung MM, Abas MI, Ramli RA.
398 Digital health for quality healthcare: A systematic mapping of review studies. Digital health.
399 2022;8:20552076221085810.

400 10. World Health organisation (WHO). Global strategy on digital health 2020-2025. Geneva:

- 401 World Health Organization; 2021. Licence: CC BY-NC-SA 3.0 IGO. Geneva
 402 <u>https://www.who.int/docs/default-</u>
- 403 source/documents/gs4dhdaa2a9f352b0445bafbc79ca799dce4d.pdf; 2021.

404 11. Achmad H, Tanumihardja M, Ramadhany YF. Teledentistry as a solution in dentistry during
405 the covid-19 pandemic period: A systematic review. International Journal of Pharmaceutical
406 Research. 2020;12:272-8.

- 407 12. Daniel SJ, Kumar S. Teledentistry: a key component in access to care. J. 2014;14
 408 Suppl:201-8.
- 409 13. Estai M, Bun S, Kanagasingam Y, Tennant M. Cost savings from a teledentistry model for

school dental screening: an Australian health system perspective. Australian Health Review.
2018;42(5):482-90.

- 412 14. Estai M, Kanagasingam Y, Tennant M, Bunt S. A systematic review of the research
 413 evidence for the benefits of teledentistry. Journal of telemedicine and telecare. 2018;24(3):147-56.
- 414 15. Marino R, Ghanim A. Teledentistry: a systematic review of the literature. Journal of
 415 Telemedicine & Telecare. 2013;19:179-83.
- 416 16. Khan SA, Omar H. Teledentistry in practice: literature review. Telemed J E Health.
 417 2013;19:565-7.

418 17. American Dental Association (ADA). ADA Policy on Teledentistry 2015 [Available from:

- 419 https://www.ada.org/en/about-the-ada/ada-positions-policies-and-statements/statement-on-
- 420 <u>teledentistry</u>.
- 421 18. World Health organisation (WHO). Mobile technologies for oral health: an implementation
- 422 guide. Geneva: World Health Organization and International Telecommunication Union, 2021.
- 423 Licence: CC BY-NC-SA 3.0 IGO https://www.who.int/publications/i/item/9789240035225. 2021.
- 424 19. Tan SHX, Lee CKJ, Yong CW, Ding YY. Scoping review: Facilitators and barriers in the
- 425 adoption of teledentistry among older adults. Gerodontology. 2021;38:351-65.

426 20. Böhm da Costa C, Peralta FDS, Ferreira de Mello ALS. How Has Teledentistry Been
427 Applied in Public Dental Health Services? An Integrative Review. Telemedicine journal and e428 health : the official journal of the American Telemedicine Association. 2019.

429 21. Estai M, Kanagasingam Y, Tennant M, Bunt S. A systematic review of the research 430 benefits teledentistry. Journal evidence for the of of telemedicine and telecare. 431 2018;24:1357633X16689433.

432 22. Aquilanti L, Santarelli A, Mascitti M, Procaccini M, Rappelli G. Dental Care Access and the
433 Elderly: What Is the Role of Teledentistry? A Systematic Review. International journal of
434 environmental research and public health. 2020;17:1-13.

435 23. Alabdullah JH, Daniel SJ. A Systematic Review on the Validity of Teledentistry.
436 Telemedicine journal and e-health : the official journal of the American Telemedicine Association.
437 2018;24:639-48.

438 24. Daniel SJ, Wu L, Kumar S. Teledentistry: A Systematic Review of Clinical Outcomes,
439 Utilization and Costs. Journal of dental hygiene : JDH / American Dental Hygienists' Association.
440 2013;87:345-52.

441 25. Madrid Troconis C, Rodriguez J, Puello P. Impact of Teledentistry Programs on Dental
442 Service in Rural Areas: A Systematic Review. International Journal of Applied Engineering
443 Research. 2018:14417-23.

444 26. Islam MRR, Islam R, Ferdous S, Watanabe C, Yamauti M, Alam MK, et al. Teledentistry as
445 an Effective Tool for the Communication Improvement between Dentists and Patients: An
446 Overview. Healthcare. 2022;10(8):1586.

Lima IFP, Vieira WD, Bernardino ID, Costa PA, Lima APB, Pithon MM, et al. Influence of
reminder therapy for controlling bacterial plaque in patients undergoing orthodontic treatment: A
systematic review and meta-analysis. Angle Orthod. 2018;88:483-93.

450 28. Mohammed H, Rizk MZ, Wafaie K, Ulhaq A, Almuzian M. Reminders improve oral 451 hygiene and adherence to appointments in orthodontic patients: a systematic review and meta-452 analysis. European journal of orthodontics. 2019;41(2):204-13.

Patil S, Hedad IA, Jafer AA, Abutaleb GK, Arishi TM, Arishi SA, et al. Effectiveness of
mobile phone applications in improving oral hygiene care and outcomes in orthodontic patients. J
Oral Biol Craniofac Res. 2021;11:26-32.

Al-Moghrabi D, Alkadhimi A, Tsichlaki A, Pandis N, Fleming PS. The influence of mobile
applications and social media-based interventions in producing behavior change among orthodontic
patients: A systematic review and meta-analysis. Am J Orthod Dentofacial Orthop. 2021;01:01.

459 31. Becker LA, Pieper D, Hartling L. Chapter V: Overviews of Reviews. In: Higgins JPT TJ,

Chandler J, Cumpston M, Li T, Page MJ, Welch VA editor. Cochrane Handbook for Systematic
Reviews of Interventions version 63 (updated February 2022) Cochrane, 2022 Available from
wwwtrainingcochraneorg/handbook2022.

463 32. Krnic Martinic M, Pieper D, Glatt A, Puljak L. Definition of a systematic review used in
464 overviews of systematic reviews, meta-epidemiological studies and textbooks. BMC Medical
465 Research Methodology. 2019;19(1):203.

Kim JSM, Pollock M, Kaunelis D, Weeks L. Guidance on review type selection for health
technology assessments: key factors and considerations for deciding when to conduct a de novo
systematic review, an update of a systematic review, or an overview of systematic reviews.
Systematic Reviews. 2022;11(1):206.

34. Rouleau G, Hong QN, Kaur N, Gagnon MP, Côté J, Bouix-Picasso J, et al. Systematic
Reviews of Systematic Quantitative, Qualitative, and Mixed Studies Reviews in Healthcare
Research: How to Assess the Methodological Quality of Included Reviews? Journal of Mixed
Methods Research. 2021:15586898211054243.

474 35. Gurgel-Juarez N, Torres-Pereira C, Haddad AE, Sheehy L, Finestone H, Mallet K, et al.

475 Accuracy and effectiveness of teledentistry: a systematic review of systematic reviews. Evid. 2022.

- 476 36. Hennessy EA, Johnson BT. Examining overlap of included studies in meta-reviews:
 477 Guidance for using the corrected covered area index. Research synthesis methods. 2020;11(1):134478 45.
- 479 37. Wafaie K, Rizk MZ, Basyouni ME, Daniel B, Mohammed H. Tele-orthodontics and sensor-
- 480 based technologies: a systematic review of interventions that monitor and improve compliance of
- 481 orthodontic patients [published online ahead of print, 2023 May 3]. *Eur J Orthod*. 2023;cjad004.
- 482 doi:10.1093/ejo/cjad004
- 483 38. Fusar-Poli P, Radua J. Ten simple rules for conducting umbrella reviews. Evidence Based
 484 Mental Health. 2018;21(3):95-100.
- 485 39. Aromataris E, F, R., , Godfrey C, Holly C, Khalil H, Tungpunkom P. Chapter 10: Umbrella
- 486 Reviews. In: Aromataris E, Munn Z (Editors). JBI Manual for Evidence Synthesis. JBI, 2020.
- 487 Available from <u>https://synthesismanual.jbi.global</u>. <u>https://doi.org/10.46658/JBIMES-20-11</u>. 2020.
- 488 40. Pollock M, Fernandes RM, Becker LA, Pieper D, Hartlin L. Chapter V: Overviews of
- 489 Reviews. In: Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA editor. Cochrane
- 490 Handbook for Systematic Reviews of Interventions version 63 (updated February 2022) Cochrane,
- 491 2022 Available from www.trainingcochraneorg/handbook2022.
- 492 41. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred
 493 reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement.
 494 Systematic Reviews. 2015;4(1):1.
- 495 42. Centre for Reviews and Dissemination. Systematic Reviews: CRD's guidance for
 496 undertaking reviews in health care. Ont, Canada: University of York, ISBN 978-1-900640-47-3.
 497 <u>https://www.york.ac.uk/media/crd/Systematic Reviews.pdf</u>; 2009.
- 498 43. Puljak L. If there is only one author or only one database was searched, a study should not
 499 be called a systematic review. Journal of clinical epidemiology. 2017;91:4-5.
- 500 44. Covidence. Covidence systematic review software, Veritas Health Innovation, Melbourne,
- 501 Australia. Available at <u>www.covidence.org</u> 2021 [

502

507

503 45. Gates A, Gates M, Duarte G, Cary M, Becker M, Prediger B, et al. Evaluation of the 504 reliability, usability, and applicability of AMSTAR, AMSTAR 2, and ROBIS: protocol for a 505 descriptive analytic study. Systematic Reviews. 2018;7(1):85.

506 46. Banzi R, Cinquini M, Gonzalez-Lorenzo M, Pecoraro V, Capobussi M, Minozzi S. Quality

assessment versus risk of bias in systematic reviews: AMSTAR and ROBIS had similar reliability

508 but differed in their construct and applicability. Journal of clinical epidemiology. 2018;99:24-32.

509 47. Rouleau G, Gagnon M-P, Côté J, Payne-Gagnon J, Hudson E, Dubois C-A, et al. Effects of

510 E-Learning in a Continuing Education Context on Nursing Care: Systematic Review of Systematic

511 Qualitative, Quantitative, and Mixed-Studies Reviews. J Med Internet Res. 2019;21(10):e15118-e.

512 48. Perry R, Whitmarsh A, Leach V, Davies P. A comparison of two assessment tools used in
513 overviews of systematic reviews: ROBIS versus AMSTAR-2. Systematic Reviews. 2021;10(1):273.

514 49. Storman M, Storman D, Jasinska KW, Swierz MJ, Bala MM. The quality of systematic
515 reviews/meta-analyses published in the field of bariatrics: A cross-sectional systematic survey using
516 AMSTAR 2 and ROBIS. Obesity reviews : an official journal of the International Association for
517 the Study of Obesity. 2020;21(5):e12994.

518 50. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical 519 appraisal tool for systematic reviews that include randomised or non-randomised studies of 520 healthcare interventions, or both. BMJ. 2017;358:j4008.

51. Whiting P, Savović J, Higgins JP, Caldwell DM, Reeves BC, Shea B, et al. ROBIS: A new
tool to assess risk of bias in systematic reviews was developed. Journal of clinical epidemiology.
2016;69:225-34.

524 52. Munthe-Kaas HM, Glenton C, Booth A, Noyes J, Lewin S. Systematic mapping of existing 525 tools to appraise methodological strengths and limitations of qualitative research: first stage in the 526 development of the CAMELOT tool. BMC Medical Research Methodology. 2019;19(1):113.

- 527 53. Rouleau G, Hong QN, Kaur N, Gagnon MP, Côté J, Bouix-Picasso J, et al. Systematic
- 528 Reviews of Systematic Quantitative, Qualitative, and Mixed Studies Reviews in Healthcare
- 529 Research: How to Assess the Methodological Quality of Included Reviews? Journal of Mixed
- 530 Methods Research.0(0):15586898211054243.
- 531 54. Papatheodorou S. Umbrella reviews: what they are and why we need them. European
- 532 Journal of Epidemiology. 2019;34.
- 533
- 534