



Munich Berlin Symptom Questionnaire (MBSQ) – Questionnaire for Adults

| Su | ırname: | | Name (physician): | | |
|----------|--|-----------------------|--|---|-------------------|
| Na | ame: | | Date (physician): | | |
| | ate of birth: | | Institution: | | |
| | | | | | |
| 10 | day's date: Completion time: | min | Date of disease onset: | | |
| The | questionnaire has to be used in medical interview. On medical evaluation of this questionnaire has to be bacterists. The diagnosis cannot be established without a | sed on the s ı | ipplementary scoring she fferential diagnostics. | et for adults. ME/CFS is | |
| | | | During the last 6 mo | | 4 |
| | | Is not present | Frequency 1 = sometimes 2 = about ½ of the time 3 = most of the time 4 = always | Severity 1 = mild 2 = moderate 3 = severe 4 = very severe | Physician's notes |
| I | Fatigue/ Daily Function | | | • | |
| 1 | Fatigue (exhaustion, tiredness) | 0 | 1 2 3 4 | 1 2 3 4 | |
| 2 | Limitations in daily life | 0 | 1 2 3 4 | 1 2 3 4 | |
| 3 | If fatigue is present, did it start new or at a definable tim | | | Yes No | |
| 4 | If fatigue is present, is it the result of ongoing, excessive | e exertion? | | Yes No | |
| 5 | If fatigue is present, is it alleviated by rest? | | | Yes No | |
| <u> </u> | | | | | |
| II | Post-Exertional Symptoms | | | | |
| 6 | Loss of physical or mental stamina | 0 | 1 2 3 4 | 1 2 3 4 | |
| 7 | Increased symptoms after mild daily activities | 0 | 1 2 3 4 | 1 2 3 4 | |
| 8 | If post-exertional symptoms are present, how long does it take after mild daily activities until the increased symptoms have subsided? | 0 ≤ 1 h 0 11-13 h | O 2-3 h O 14-23 h | O 4-10 h O ≥ 24 h | |
| 9 | If post-exertional symptoms are present, which three mild daily activities lead to a worsening of the symptoms? | 2 | | | |
| 10 | If post-exertional symptoms are present, which three symptoms worsen after mild daily activities? | 1 2 | | | |
| | | | D 1 11 1 10 | 4 | |
| | | Not present | During the last 6 more Frequency 1 = sometimes 2 = about ½ of the time 3 = most of the time 4 = always | Severity 1 = mild 2 = moderate 3 = severe 4 = very severe | Physician's notes |
| III | Sleep | | · · · · · · · · · · · · · · · · · · · | | |
| 11 | Unrefreshing sleep Unusually much sleep during the day | 0 | 1 2 3 4 | 1 2 3 4 | |
| 13 | Problems falling asleep | 0 | 1 2 3 4 | 1 2 3 4 | |
| 14 | Problems sleeping through | 0 | 1 2 3 4 | 1 2 3 4 | |
| 15 | Day / night reversal | 0 | 1 2 3 4 | 1 2 3 4 | |
| | | | | | |
| IV | Pain | | 4 0 0 4 | | |
| 16 17 | Myofascial pain (can include achy and sore muscles) Joint pain without swelling or redness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 18 | Headaches | 0 | 1 2 3 4 | 1 2 3 4 | |
| | | | | | |
| ٧ | Neurocognitive Manifestations | | | | |
| 19 | Confusion | 0 | 1 2 3 4 | 1 2 3 4 | |
| | | 0 | 1 2 3 4 | 1 2 3 4 | |
| 21 | Concentration problems Memory problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| ~~ | METHOLY PLONETIE | U | 1 2 3 4 | 1 4 3 4 | |

| | | | During the last 6 m | onths | |
|---------------------|---|--------------|---|----------|-------------------|
| | | Not present | Frequency 1 = sometimes 2 = about ½ of the time 3 = most of the time 4 = always | Severity | Physician's notes |
| 23 | Orientation problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 24 | Comprehension problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 25 | Word-finding problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 26 | If cognitive symptoms (V.20-25) are present, do they we effort / stress or time pressure? | orsen due to | | Yes No | |
| 27 | Perceptual and sensory disturbances, e.g., blurred vision | 0 | 1 2 3 4 | 1 2 3 4 | |
| 28 | Coordination problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 29 | Muscle twitches | 0 | 1 2 3 4 | 1 2 3 4 | |
| 30 | Muscle weakness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 31 | Overload phenomena - hypersensitivity to (bright) light | 0 | 1 2 3 4 | 1 2 3 4 | |
| 32 | Overload phenomena - hypersensitivity to noise | 0 | 1 2 3 4 | 1 2 3 4 | |
| 33 | Overload phenomena - hypersensitivity to touch | 0 | 1 2 3 4 | 1 2 3 4 | |
| 34 | Overload phenomena – emotional overload | 0 | 1 2 3 4 | 1 2 3 4 | |
| VI | Autonomic Manifestations | | | | |
| 35 | Dizziness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 36 | Palpitations with or without cardiac arrhythmias | 0 | 1 2 3 4 | 1 2 3 4 | |
| 37 | Circulatory problems when getting up | 0 | 1 2 3 4 | 1 2 3 4 | |
| 38 | Circulatory problems in an upright posture | 0 | 1 2 3 4 | 1 2 3 4 | |
| 39 | Unusual paleness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 40 | Bladder problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| _ | Gastrointestinal problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| | Exertional shortness of breath | 0 | 1 2 3 4 | 1 2 3 4 | |
| ,,,, | | | | | • |
| VII | Neuroendocrine Manifestations | T I | | | |
| 43 | Loss of thermostatic stability (e.g., subnormal body | | | | |
| | temperature, and marked diurnal fluctuation, sweating episodes, feelings of feverishness, cold | 0 | 1 2 3 4 | 1 2 3 4 | |
| | limbs) | | | | |
| 14 | Intolerance of extremes hot and cold | 0 | 1 2 3 4 | 1 2 3 4 | |
| 14 45 | Marked weight change – loss of appetite or abnormal | | 1 2 3 4 | 1 2 3 4 | |
| 70 | appetite | 0 | 1 2 3 4 | 1 2 3 4 | |
| 46 | Worsening of symptoms with stress | 0 | 1 2 3 4 | 1 2 3 4 | |
| VIII | Immunologic Manifestations | | | | |
| 47 | Flu like symptoms / general malaise | 0 | 1 2 3 4 | 1 2 3 4 | |
| 47 48 | Painful lymph nodes | 0 | 1 2 3 4 | 1 2 3 4 | |
| 40 49 | | 0 | 1 2 3 4 | 1 2 3 4 | |
| 49 50 | New sensitivities to food, medication or chemicals | 0 | 1 2 3 4 | 1 2 3 4 | |
| F | ther Questions About the Symptom Course | | | | |
| | ere are any symptoms, | | | | |
| | at are the three main symptoms of this questionnaire? | 1 | | | |
| | • • • | 2 | | | |

For evaluation of this questionnaire please use the supplementary scoring sheet for adults.





Munich Berlin Symptom Questionnaire (MBSQ) – Questionnaire for Pediatric and Adolescent Patients

| 1110 | mon Bornir Oymptom Quootionnano (mBOQ) | Quoonom | nano ioi i odiatilo ana 7 | idologodii i dilolito | |
|----------|---|------------------------------------|---|---|-------------------|
| Su | ırname: | | Name (physician): | | |
| Na | ame: | | Date (physician): | | |
| Da | ate of birth: | | Institution: | | |
| - | | min | Date of disease onset | | |
| | day's date: Completion time: | | | | |
| For med | <u>patients</u> : Please complete this questionnaire on your on <u>doctors</u> : This questionnaire has to be used in medical dical visit . The medical evaluation of this questionnal ical diagnosis . The diagnosis cannot be established we | I interview. (ire has to be | Open questions and comprobased on the supplementa | ehensive problems shou ry scoring sheet for adu | |
| | | | During the last 3 mo | onths | |
| | | Is not present | Frequency 1 = sometimes 2 = about ½ of the time 3 = most of the time 4 = always | Severity 1 = mild 2 = moderate 3 = severe 4 = very severe | Physician's notes |
| I | Fatigue / Daily Function | _ | | | |
| 1 | Fatigue (exhaustion, tiredness) | 0 | 1 2 3 4 | 1 2 3 4 | |
| 3 | Limitations in daily life - school/ education Limitations in daily life - social | 0 | 1 2 3 4 | 1 2 3 4 | |
| 4 | Limitations in daily life - social Limitations in daily life - personal | 0 | 1 2 3 4 | 1 2 3 4 | |
| | | · · | 1 2 3 4 | 1 2 3 4 | |
| 5 | If fatigue is present, did it start new or at a definable tim | | | Yes No | |
| 6 | If fatigue is present, is it the result of ongoing, excessive | e exertion? | | Yes No | |
| 7 | If fatigue is present, is it alleviated by rest? | | | Yes No | |
| ш | Post-Exertional Symptoms | | | | |
| 8 | | 0 | 1 2 3 4 | 1 2 3 4 | |
| 9 | Loss of physical or mental stamina Increased symptoms after light everyday activities | 0 | 1 2 3 4 | 1 2 3 4 | |
| 10 | If post-exertional symptoms are present, | - | | | |
| | how long does it take after mild daily activities until the increased symptoms have subsided? | O ≤ 1 h O 11-13 | | O 4-10 h O ≥ 24 h | |
| 11 | If post-exertional symptoms are present, which three mild daily activities lead to a worsening of the symptoms? | | | | |
| 12 | If post-exertional symptoms are present, which three symptoms worsen after mild daily activities? | 1 2 3 | | | |
| | | 1 | During the last 3 mg | onthe | |
| | | Not present | Frequency 1 = sometimes 2 = about ½ of the time 3 = most of the time 4 = always | Severity 1 = mild 2 = moderate 3 = severe 4 = very severe | Physician's notes |
| | Sleep | 1 - | | | |
| 13 | Unrefreshing sleep | 0 | 1 2 3 4 | 1 2 3 4 | |
| 14 15 | Unusually much sleep during the day Problems falling asleep | 0 | 1 2 3 4 | 1 2 3 4 | |
| 16 | Problems sleeping through | 0 | 1 2 3 4 | 1 2 3 4 | |
| 17 | Day/ night reversal | 0 | 1 2 3 4 | 1 2 3 4 | |
| | | | <u> </u> | <u> </u> | |
| IV | Pain | | | | |
| 18 | Myofascial pain (can include achy and sore muscles) | 0 | 1 2 3 4 | 1 2 3 4 | |
| 19 | Joint pain without swelling or redness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 20 | Headaches Abdominal pain | 0 | 1 2 3 4 | 1 2 3 4 | |
| - 4 | ravaviiii(d) VdIII | . U | 1 4 3 4 | 1 4 3 4 | |

| | | | During the last 3 m | | |
|------|---|-------------|---|----------------------------|-------------------|
| | | | Frequency | Severity | |
| | | . | 1 = sometimes | 1 = mild | Physician's notes |
| | | Not present | 2 = about ½ of the time 3 = most of the time | 2 = moderate 3 = severe | |
| | | | 4 = always | 4 = very severe | |
| ٧ | Neurocognitive Manifestations | ı | 1 amayo | 1 1013 001010 | |
| 22 | Confusion | 0 | 1 2 3 4 | 1 2 3 4 | |
| 23 | Slowness of thought | 0 | 1 2 3 4 | 1 2 3 4 | |
| 24 | Concentration problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 25 | Memory problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 26 | Orientation problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| | Comprehension problems | 0 | | 1 2 3 4 | |
| 27 | | 0 | 1 2 3 4 | | |
| 28 | Word-finding problems | U | 1 2 3 4 | 1 2 3 4 | |
| 29 | If cognitive symptoms (V.23-28) are present, do they wo | rsen due to | | | |
| | Effort / stress or time pressure? | | | Yes No | |
| | ' | 1 | | | |
| 30 | Absent mindedness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 31 | Difficulty recalling information | 0 | 1 2 3 4 | 1 2 3 4 | |
| 32 | Need to focus on one thing at a time | 0 | 1 2 3 4 | 1 2 3 4 | |
| 33 | Trouble expressing thought | 0 | 1 2 3 4 | 1 2 3 4 | |
| 34 | Lose train of thought | 0 | 1 2 3 4 | 1 2 3 4 | |
| 35 | New troubles with math or other educational subjects | 0 | 1 2 3 4 | 1 2 3 4 | |
| 36 | Perceptual and sensory disturbances, e.g., blurred | 0 | 1 2 3 4 | 1 2 3 4 | |
| | vision | - | | | |
| 37 | Coordination problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 38 | Muscle twitches | 0 | 1 2 3 4 | 1 2 3 4 | |
| 39 | Muscle weakness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 40 | Overload phenomena - hypersensitivity to (bright) light | 0 | 1 2 3 4 | 1 2 3 4 | |
| 41 | Overload phenomena - hypersensitivity to noise | 0 | 1 2 3 4 | 1 2 3 4 | |
| 42 | Overload phenomena - hypersensitivity to touch | 0 | 1 2 3 4 | 1 2 3 4 | |
| 43 | Overload phenomena – emotional overload | 0 | 1 2 3 4 | 1 2 3 4 | |
| 43 | Overload prieriorneria – emotioriai overload | U | 1 2 3 4 | 1 2 3 4 | |
| | A | | | | |
| VI | Autonomic Manifestations | | | | |
| 44 | Dizziness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 45 | Palpitations with or without cardiac arrhythmias | 0 | 1 2 3 4 | 1 2 3 4 | |
| 46 | Circulatory problems when getting up | 0 | 1 2 3 4 | 1 2 3 4 | |
| 47 | Circulatory problems in upright posture | 0 | 1 2 3 4 | 1 2 3 4 | |
| 48 | Unusual paleness | 0 | 1 2 3 4 | 1 2 3 4 | |
| 49 | Bladder problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 50 | Gastrointestinal problems | 0 | 1 2 3 4 | 1 2 3 4 | |
| 51 | Exertional shortness of breath | 0 | 1 2 3 4 | 1 2 3 4 | |
| | | | | | |
| VII | Neuroendocrine Manifestations | | | | |
| 52 | Loss of thermostatic stability (e.g., subnormal body | | | | |
| | temperature, and marked diurnal fluctuation, | | | | |
| | sweating episodes, feelings of feverishness, cold | 0 | 1 2 3 4 | 1 2 3 4 | |
| | limbs) | | | | |
| 53 | Intolerance of extremes hot and cold | 0 | 1 2 3 4 | 1 2 3 4 | |
| 54 | Marked weight change – loss of appetite or abnormal | - | | | |
| 7 . | appetite | 0 | 1 2 3 4 | 1 2 3 4 | |
| 55 | Worsening of symptoms with stress | 0 | 1 2 3 4 | 1 2 3 4 | |
| | . O : | <u> </u> | <u> </u> | <u> </u> | |
| VIII | Immunologic Manifestations | | | | |
| 56 | Flu like symptoms / general malaise | 0 | 1 2 3 4 | 1 2 3 4 | |
| 57 | Painful lymph nodes | 0 | 1 2 3 4 | 1 2 3 4 | |
| 58 | Fever | 0 | 1 2 3 4 | 1 2 3 4 | |
| 59 | Recurrent sore throat | 0 | 1 2 3 4 | 1 2 3 4 | |
| 60 | New sensitivities to food, medication or chemicals | 0 | 1 2 3 4 | 1 2 3 4 | |
| UU | New sensitivities to 1000, medication of chemicals | U | 1 2 3 4 | 1 2 3 4 | |
| - | 41 O C Al (41 O () | | | | |
| | ther Questions About the Symptom Course | | | | |
| | ere are any symptoms, | 1 | | | |
| wha | at are the three main symptoms of this questionnaire? | | | | |
| | | | | | |
| | | 3 | | | |





Munich Berlin Symptom Questionnaire – Scoring Sheet for Adults (≥ 18 Years of Age)

| Only if frequency and severity are reported with ≥2, the respective item counts positively for evaluation |
|---|
|---|

| Surname: | Surname (physician): |
|----------------|----------------------|
| Name: | Name (physician): |
| Date of birth: | Date (physician): |
| Today's date: | Institution: |

| Cai | nadian Consensus Criteria¹ | 101 | M Criteria ² |
|-----|--|-----|--|
| 0 | Duration of Illness (Onset of the Symptoms:) The symptomatology is present for at least 6 months. | 0 | Duration of Illness (Onset of the Symptoms:,) The symptomatology is present for at least 6 months. |
| 0 | Medical History, Physical Examination, and Differential Diagnostics did not indicate any other cause for the symptoms (in particular, no indications for Addison's disease, Cushing's disease, hypo- /hyperthyroidism, anemia, hemochromatosis, diabetes mellitus, haemato-oncological, rheumatological and treatable sleep disorders) The symptoms must have begun or have been significantly altered after the onset of the illness. | 0 | Medical History, Physical Examination, and Differential Diagnostics did not indicate any other cause for the symptoms The symptoms must have begun or have been significantly altered after the onset of the illness. |
| 0 | Fatigue / Daily Function All of the following points must apply: Fatigue: 1.1 ≥2 Limitations: at least 1 of the 3 points 1.2 ≥2 Fatigue new start: 1.3 Yes | 0 | Fatigue / Daily Function All of the following points must apply: Fatigue: 1.1 ≥2 Limitations: at least 1 of the 3 points 1.2 ≥2 Fatigue new start: 1.3 Yes Fatigue due to exertion: 1.4 No Fatigue alleviated by rest: 1.5 No |
| 0 | Post-Exertional Symptoms All of the following points must apply: loss of stamina; increased symptoms after everyday activities: all 2 points II.6-7 ≥2 duration of worsening of symptoms: II.8 ≥14 hours* | 0 | Post-Exertional Symptoms All of the following points must apply: all 2 points II.6-7 ≥2 |
| 0 | Sleep At least 1 of the following 5 points must apply: III.11-15 ≥2 | 0 | Sleep The following point must apply: III.11 ≥2 |

| Canadian Consensus Criteria¹ | | IOI | IOM Criteria ² | |
|------------------------------|--|-----|---|--|
| 0 | Pain At least 1 of the following 5 points must apply: VI.16-18 ≥2 | | Pain Not included in IOM criteria | |
| 0 | Neurocognitive Manifestations At least 2 of the following symptom groups must apply: Confusion: V.19 ≥2 Concentration and/or Memory: 1 of the following 2 points: V.21-22 ≥2 Orientation: V.23 ≥2 Comprehension and/or word-finding: 1 of the following 2 points: V.24-25 ≥2 Perception and overload phenomena: 1 of the following 5 points: V.27, V.31-34 ≥2 Coordination and muscular system: 1 of the following 3 points: V.28-30 ≥2 | 0 | Neurocognitive and Autonomous Manifestations At least 1 of the following 2 categories must apply: The 2 following symptom groups must apply: At least 1 of the following 6 points must apply: V.20-22, V.24-25, V.28 ≥2 The following point must apply: V.26 Yes At least 1 of the following 2 points must apply: VI.37-38 ≥2 | |
| 0 | Autonomic, Neuroendocrine, Immunologic Manifestations At least 2 of the following 3 categories must apply: At least 1 of the following 8 points must apply: VI.35-42 ≥2 At least 1 of the following 4 points must apply: VII.43-46 ≥2 At least 1 of the following 4 points must apply: VIII.47-50, ≥2 | | Neuroendocrine, Immunologic Manifestations Not included in IOM criteria | |

| O Patient meets Canadian Consensus Criteria for ME/CFS | O Patient meets IOM criteria for ME/CFS |
|--|---|
| O Patient does not meet Canadian Consensus Criteria for ME/CFS | O Patient does not meet IOM criteria for ME/CFS |

In discussion with the patient & in the physician's letter, it must be pointed out that ME/CFS is a diagnosis of exclusion, which must be re-evaluated in case of new clinical aspects.

¹ Carruthers, Bruce & Jain, Anil & Meirleir, Kenny & Peterson, Daniel & Klimas, Nancy & Lerner, A. & Bested, Alison & Flor-Henry, Pierre & Joshi, Pradip & Powles, Peter & Sherkey, Jeffrey & Sande, Marjorie. (2003). Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. Journal Of Chronic Fatigue Syndrome. 11. 7-115. 10.1300/J092v11n01_02.

² Clayton, Ellen. (2015). Beyond Myalgic Encephalomyelitis/Chronic Fatigue Syndrome An IOM Report on Redefining an Illness. JAMA. 313. 10.1001/jama.2015.1346.

^{*} In the original publication of Carruthers et al.¹ PEM requires a duration of >24 hours. However, according to the current state of international research, we recommend diagnosing PEM with a duration of ≥14 hours (see Cotter J et al. (2018) A Brief Questionnaire to Assess Post-Exertional Malaise. Diagnostics (Basel). 2018;8(3):66. Doi: 10.3390/diagnostics8030066 und Kedor, C., Freitag, H., Meyer-Arndt, L. et al. A prospective observational study of post-COVID-19 chronic fatigue syndrome following the first pandemic wave in Germany and biomarkers associated with symptom severity. Nat Commun 13, 5104 (2022). https://doi.org/10.1038/s41467-022-32507-6).





<u>Munich Berlin Symptom Questionnaire – Scoring Sheet for Children and Adolescents (0-17 Years of Age)</u>

Only if frequency and severity are reported with ≥2, the respective item counts positively for evaluation.

| Surname: | Surname (physician): |
|----------------|----------------------|
| Name: | Name (physician): |
| Date of birth: | Date (physician): |
| Today's date: | Institution: |

| Pediatric Case Definition for ME/CFS by Jason et al.¹ | | CI | inical Diagnostic Worksheet by Rowe et al. ² | | |
|---|---|----|---|--|--|
| 0 | Duration of Illness (Onset of the Symptoms:) The symptomatology is present for at least 3 months. | 0 | Duration of Illness (Onset of the Symptoms:) The symptomatology is present for at least 3 months. | | |
| 0 | Medical History, Physical Examination and Differential Diagnostics did not indicate any other cause for the symptoms (in particular, no indications for untreated hypothyreoidism, sleep apnea, narcolepsy, malignancies, leukemia, unresolved hepatitis, multiple sclerosis, juvenile rheumatoid arthritis, lupus erythematosus, HIV/ AIDS, severe obesity (BMI greater than 40), celiac disease, lyme disease, childhood schizophrenia or psychiatric disorders, bipolar disorder, active alcohol or substance abuse, active anorexia nervosa or bulimia nervosa, depressive disorders) | 0 | Medical History, Physical Examination and Differential Diagnostics did not indicate any other cause for the symptoms (in particular, no evidence of adrenocortical insufficiency, overtraining syndrome, GI disorders: Celiac disease irritable bowel syndrome, eosinophilic gastroenteritis; Chiari malformation, cervical spinal stenosis, neuroborreliosis or other tick-borne disease, major depression, narcolepsy, obstructive or central sleep apnea, postcommotion syndrome, severe anemia, systemic lupus erythematosus and similar autoimmune disease, untreated hypor/hyperthyroidism). | | |
| 0 | Fatigue / Daily Function All of the following points must apply: Fatigue: I.1 ≥2 Limitations: all 3 points: I.2-4 ≥2 Fatigue due to exertion: I.6 No Fatigue alleviated by rest: I.7 No | 0 | Fatigue / Daily Function All of the following points must apply: Fatigue: I.1 ≥2 Limitations: at least 1 of the 3 points: I.2-4 ≥2 Fatigue due to exertion: I.6 No Fatigue alleviated by rest: I.7 No | | |
| 0 | Post-Exertional Symptoms All of the following points must apply: loss of stamina; increased symptoms after everyday activities: all 2 points: II.8-9 ≥2 duration of worsening of symptoms: II.10 ≥14 hours | 0 | Post-Exertional Symptoms All of the following points must apply: loss of stamina; increased symptoms after everyday activities: all 2 points: II.8-9 ≥2 duration of worsening of symptoms: II.10 >24 hours | | |





| Pediatric Case Definition for ME/CFS by Jason et al. ¹ | | CI | Clinical Diagnostic Worksheet by Rowe et al. ² | | |
|---|--|----|--|--|--|
| 0 | Sleep At least 1 of the following 5 points must apply: III.13-17 ≥2 | 0 | Sleep, Pain, and Neurocognitive Symptoms At least 2 of the following 3 categories must apply: At least 1 of the following 5 points must apply: III.13-17 ≥2 | | |
| 0 | Pain At least 1 of the following 4 points must apply: IV.18-21 ≥2 | | At least 1 of the following 6 points must apply: IV.18-21, IV.57, IV.59 ≥2 At least 1 of the following 8 points must apply: V.23-25, V.27-28, V.30, V.33 ≥2; V.29: Yes | | |
| 0 | Neurocognitive Manifestations At least 2 of the following 11 points must apply: V.23-25; V.27-28; V.30-35 ≥2 | | | | |
| 0 | Autonomic, Neuroendocrine and Immunologic Manifestations At least 2 of the following 3 categories must apply: At least 1 of the following 5 points must apply: VI.44-47; VI.51 ≥2 At least 1 of the following 4 points must apply: VII.52-55 ≥2 At least 1 of the following 4 points must apply: VIII.56-59 ≥2 | 0 | Autonomic, Neuroendocrine, and Immunologic Manifestations Not included in the Clinical Diagnostic Worksheet by Rowe | | |
| 0 | Patient meets the Pediatric Case Definition for ME/CFS by Jason | 0 | Patient meets the Clinical Diagnostic Worksheet criteria for ME/CFS by Rowe | | |
| 0 | Patient does not meet the Pediatric Case Definition for ME/CFS by Jason | 0 | Patient does not meet the Clinical Diagnostic Worksheet criteria for ME/CFS by Rowe | | |

In the discussion with patients and parents as well as in the physician's letter it must be pointed out that ME/CFS is clinical diagnosis which must be re-evaluated in the case of new clinical aspects.

¹Jason, Leonard & Jordan, Karen & Miike, Teruhisa & Bell, David & Lapp, Charles & Torres-Harding, Susan & Rowe, Kathy & Gurwitt, Alan & Meirleir, Kenny & Hoof, Elke & Psych, Clin & Chairperson,. (2006). A Pediatric Case Definition for Myalgic Encephalomyelitis and Chronic Fatigue Syndrome. Journal Of Chronic Fatigue Syndrome. 13. 1-44. 10.1300/J092v13n02_01.

² Rowe, Peter & Underhill, Rosemary & Friedman, Kenneth & Gurwitt, Alan & Medow, Marvin & Schwartz, Malcolm & Speight, Nigel & Stewart, Julian & Vallings, Rosamund & Rowe, Katherine. (2017). Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Diagnosis and Management in Young People: A Primer. Frontiers in Pediatrics. 5. 10.3389/fped.2017.00121.

* In the original publication by Rowe et al.² a duration of at least 6 months is required for the diagnosis. If the duration of the disease is <6 months, it is only recommended to express an urgent suspicion. We recommend a diagnosis in children and adolescents after a duration of 3 months in order to provide symptom-oriented care at an early stage.





<u>Munich Berlin Symptom Questionnaire – Additional Scoring Sheet</u> <u>for Adolescents (12-17 Years)</u>

Only if frequency and severity are reported with ≥2, the respective item enters positively into the evaluation.

| Surname: | Surname (physician): |
|----------------|----------------------|
| Name: | Name (physician): |
| Date of birth: | Date (physician): |
| Today's date: | Institution: |

| Canadian Consensus Criteria ³ | | IOM Criteria ⁴ | |
|--|--|---------------------------|--|
| 0 | Duration of Illness (Onset of the Symptoms:) | 0 | Duration of Illness (Onset of the Symptoms:) |
| | The symptomatology has been present for at least 3 months. | | The symptomatology has been present for at least 3 months.* |
| 0 | did not indicate any other cause for the symptoms | 0 | Medical History, Physical Examination and Differential Diagnostics did not indicate any other cause for the symptoms |
| | (in particular no indications for Addison's disease, Cushing's disease, hypo-/hyperthyroidism, anemia, hemochromatosis, diabetes mellitus, haemato-oncological, rheumatological and treatable sleep disorders) | | |
| | The symptoms must have begun or have been significantly altered after the onset of the illness. | | The symptoms must have begun or have been significantly altered after the onset of the illness. |
| | | | |
| 0 | Fatigue / Daily Function | 0 | Fatigue / Daily Function |
| | All of the following points must apply: Fatigue: I.1 ≥2 | | All of the following points must apply: Fatigue: I.1 ≥2 |
| | Limitations: at least 1 of the 3 points: I.2-4 \geq 2 | | Limitations: at least 1 of the 3 points: I.2-4 ≥2 |
| | Fatigue new start: I.5 Yes | | Fatigue new start: I.5 Yes Fatigue due to exertion: I.6 No |
| | | | Fatigue alleviated by rest: I.7 No |
| 0 | Post-Exertional Symptoms | 0 | Post-Exertional Symptoms |
| | All of the following points must apply: loss of stamina; increased symptoms after everyday activities: all 2 points II.8-9 ≥2 duration of worsening of symptoms: II.10 >24 hours | | All of the following points must apply: all 2 points: II.8-9 ≥2 |





| Ca | anadian Consensus Criteria³ | IC | M Criteria⁴ |
|----|--|----|--|
| 0 | Sleep At least 1 of the following 5 points must apply: III.13-17 ≥2 | 0 | Sleep The following point must apply: III.13 ≥2 |
| 0 | Pain At least 1 of the following 3 points must apply: IV.18-20 ≥2 (except abdominal pain) | 0 | Pain Not included in IOM criteria |
| 0 | Neurocognitive Manifestations At least 2 of the following symptom groups must apply: Confusion: V.22 ≥2 Concentration and/or Memory: 1 of the following 2 points: V.24-25 ≥2 Orientation: V.26 ≥2 Comprehension and/or word-finding: 1 of the following 2 points: V.27-28 ≥2 Perception and overload phenomena: 1 of the following 5 points: V.36, V.40-43 ≥2 Coordination und muscular system: 1 of the following 3 points: V.37-39 ≥2 | 0 | Neurocognitive and Autonomic Manifestations At least 1 of the following 2 categories must apply: The 2 following symptom groups must apply: At least 1 of the following 6 points must apply: V.23-25, V.27-28, V.37 ≥2 The following point must apply: V.29, Yes At least 1 of the following 2 points must apply: VI.46-47 ≥2 |
| 0 | Autonomic, Neuroendocrine, Immunologic Manifestations At least 2 of the following 3 categories must apply: At least 1 of the following 8 points must apply: VI.44-51 ≥2 At least 1 of the following 4 points must apply: VII.52-55 ≥2 At least 1 of the following 4 points must apply: VIII.56-57, VIII.59-60 ≥2 | 0 | Neuroendocrine, Immunologic Manifestations Not included in IOM criteria |
| | | | |

| O Patient meets Canadian Consensus Criteria for ME/CFS | O Patient meets IOM criteria for ME/CFS |
|--|---|
| O Patient does not meet Canadian Consensus Criteria for ME/CFS | O Patient does not meet IOM criteria for ME/CFS |

In the discussion with patients and parents as well as in the physician's letter it must be pointed out that ME/CFS is clinical diagnosis which must be re-evaluated in the case of new clinical aspects.

³ Carruthers, Bruce & Jain, Anil & Meirleir, Kenny & Peterson, Daniel & Klimas, Nancy & Lerner, A. & Bested, Alison & Flor-Henry, Pierre & Joshi, Pradip & Powles, Peter & Sherkey, Jeffrey & Sande, Marjorie. (2003). Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. 11. 7-115. 10.1300/J092v11n01_02.

⁴ Clayton, Ellen. (2015). Beyond Myalgic Encephalomyelitis/Chronic Fatigue Syndrome An IOM Report on Redefining an Illness. JAMA. 313. 10.1001/jama.2015.1346.

^{*} In the original publication by Clayton et al.4 a duration of at least 6 months is required for the diagnosis. We recommend diagnosing children and adolescents after a disease duration of 3 months to provide them with symptom-oriented care at an early stage.