

Table S1. Number of days for each activity-counting pattern

Activity patterns	No of days	% of days	Daily activity count (mean ± SD)
AD	18980	41.00	14786.06±5454.00
M	19156	41.38	14004.09±5405.45
E	5033	10.87	12586.48±5551.76
BP	1932	4.17	4640.34±5755.53
IM	1190	2.57	7016.28±7419.17
Total	46291	100	

Table S2. Number of participants for behavior

Activity-count Behavior	No of participants	% of participants	Total activity count (mean ± SD)
AD dominant	1792	27.09	14996.28±4636.84
M dominant	1964	29.69	13628.09±4594.38
AD+M dominant	1855	28.05	13307.74±5063.63
E dominant	1002	15.15	11589.80±5254.22
Total	6613	100	

Table S3. Proportion of activity-counting patterns in each behavior

Activity-count Behavior	Activity-counting pattern probability (Mean ± SD)				
	AD	M	E	BP	IM
AD dominant	0.7900±0.1281	0.1201±0.0192	0.0668±0.1016	0.0140±0.0499	0.0091±0.0395
M dominant	0.1269±0.1114	0.8341±0.1303	0.0135±0.0505	0.0175±0.0580	0.0081±0.0371
AD+M dominant	0.4342±0.1340	0.4396±0.1168	0.0433±0.0782	0.0594±0.1264	0.0236±0.0707
E dominant	0.2407±0.1772	0.0677±0.0958	0.4916±0.2396	0.1061±0.1793	0.0940±0.1604

Table S4: Sensitivity analysis (Odd ratios for depressive symptoms by activity behavior)

Model	Behavior pattern				P value
	M dominant	AD dominant	AD+M dominant	E dominant	p-value
Model 1	1.00 (ref)	1.21(0.90,1.64)	1.24(0.92,1.67)	2.48(1.86,3.30)	<0.0001
Model 2	1.00 (ref)	1.32(0.96,1.79)	1.32(0.98,1.78)	2.88(2.11,3.96)	<0.0001
Model 3	1.00 (ref)	1.23(0.91,1.66)	1.22(0.91,1.64)	2.48(1.87,3.31)	<0.0001
Model 4	1.00 (ref)	1.16(0.86,1.57)	1.22(0.90,1.64)	2.56(1.92,3.41)	<0.0001
Model 5	1.00 (ref)	1.18(0.87,1.60)	1.27(0.95,1.72)	2.52(1.89,3.38)	<0.0001
Model 6	1.00 (ref)	1.23(0.92,1.67)	1.19(0.89,1.61)	2.35(1.76,3.14)	<0.0001

Notes

Model 1: Activity behavior (Crude)

Model 2: Activity behavior, Age

Model 3: Activity behavior, BMI

Model 4: Activity behavior, Gender

Model 5: Activity behavior, work situation

Model 6: Activity behavior, total activity