Interrogating Changes in COVID-19 Vaccine Intent During the National Vaccine Rollout in the Philippines

Alexandria Caple¹, Arnie Dimaano², Marc Martin Sagolili², April Anne Uy², Panjee Mariel Aguirre³, Dean Alano², Kaila Andrine Baul², Giselle Sophia Camaya², Angelo Raine Carlos², Brent John Ciriaco³, Princess Jerah Mae Clavo², Dominic Cuyugan², Cleinne Florence Geeseler Fermo², Paul Jeremy Lanete², Ardwayne Jurel La Torre³, Thomas Loteyro³, Raisa Mikaela Lua², Nicole Gayle Manansala², Raphael Mosquito², Alexa Marie Octaviano², Alexandra Erika Orfanel², Anne Elizabeth Papa², Gheyanna Merly Pascual², Mark Nhel Peralta², Aubrey Joy Sale², Sophia Lorraine Tendenilla², Maria Sofia Lauren Trinidad², Nicole Jan Trinidad², Daphne Louise Verano², and Nicanor Austriaco^{2,4,5*} ¹Department of Psychology, Providence College, 1 Cunningham Square, Providence, Rhode Island 02918, U.S.A. ²Department of Biological Sciences, University of Santo Tomas, España Blvd., Manila 1015, PHILIPPINES ³Department of Advertising Arts, University of Santo Tomas, España Blvd., Manila 1015, PHILIPPINES ⁴Research Center for the Natural and Applied Sciences, University of Santo Tomas, España Blvd., Manila 1015, PHILIPPINES ⁵UST Laboratories for Vaccine Science, Molecular Biology, and Biotechnology, University of Santo Tomas, España Blvd., Manila 1015, PHILIPPINES Corresponding Author: Rev. Nicanor Austriaco, OP, UST Laboratories for Vaccine Science, Molecular Biology, and Biotechnology, University of Santo Tomas, España Blvd., Manila 1015, PHILIPPINES Email address: naustriaco@ust.edu.ph

Abstract

To mitigate the unprecedented health, social, and economic damage of COVID-19, the Philippines implemented a nation-wide vaccine program to mitigate the effects of the global pandemic. In a previous study, we interrogated COVID-19 vaccine intent in the country by deploying a nationwide open-access online survey, two months before the initial rollout of the national vaccination program that began in March of 2021. In this follow-up study, we explored the influence of the ongoing vaccine rollout on vaccine intent by deploying a similar survey six months after the initial rollout of vaccines had begun throughout the archipelago. Our data suggests that the presence of vaccines and vaccinated individuals in a Filipino community predicts vaccination intent. When directly asked if they were more willing to receive a vaccination than six months prior, 92.26% agreed to some extent that they were indeed more willing. Finally, despite the changes in the numbers of respondents who were more open to the COVID-19 vaccines, there were no significant changes in the predictive power of our Health Belief Model (HBM) constructs. For the most part, the HBM factors that predicted vaccination intent in our earlier study also predicted vaccination intent in this current study.

Introduction

On January 30, 2020, the Department of Health (DOH) of the Philippines reported its first case of COVID-19, a novel respiratory disease first identified in Wuhan, China, that is caused by the coronavirus, SARS-CoV-2 (Xie et al., 2020; Zhu et al., 2020; Guan et al., 2020; Reis et al., 2022). With widespread human-to-human transmission, the virus is highly contagious, and the COVID-19 pandemic became a global concern of historic proportions (van Ginneken et al., 2022; Siddiqui, Alhamdi & Alghamdi, 2022; Sachs et al., 2022; Cassell et al., 2022). As of April 5, 2023, there have been 4,082,418 confirmed cases and 66,390 confirmed deaths from COVID-19 reported by the DOH throughout the archipelago (www.covid19.gov.ph).

Vaccination has long been regarded as the most effective means for combating infectious disease (Rappuoli et al., 2014; Sathyanarayana et al., 2020). The Philippines began its national vaccine drive against COVID-19 on March 1, 2021, with the goal of vaccinating seventy million of its citizens by the end of the calendar year (Inter-Agency Task Force for the Management of Emerging Infectious Disease, 2021). One of the ongoing challenges for this campaign is the vaccine hesitancy among the Filipino people (Alfonso et al., 2021). Though immunization rates had been relatively high in the Philippines for many decades, the controversial 2016 rollout of the dengue vaccine, Dengvaxia, triggered significant drops in the rates of immunization as Filipino parents refused to have their children routinely vaccinated against polio, chicken pox, and tetanus (Fatima & Syed, 2018; Smith, 2018; Reñosa et al., 2022).

The Health Belief Model (HBM) posits that people are likely to adopt disease prevention behaviors and to accept medical interventions like vaccines if there is sufficient motivation and cues to action (Rosenstock, Strecher & Becker, 1988; Limbu, Gautam & Pham, 2022a; Limbu & Gautam, 2023). Motivational factors include perceived susceptibility to and severity of the disease and perceived benefits of the vaccine. Cues to action include information, people, and events that nudge the individual towards vaccination. The HBM has been adopted as a conceptual framework that has been used to evaluate the beliefs and attitudes toward a diversity of vaccines including the influenza, human papillomavirus, and hepatitis B vaccines (Teitler-Regev, Shahrabani & Benzion, 2011; Donadiki et al., 2014; Hu et al., 2017; Chen et al., 2019). Moreover, several studies have shown that the HBM constructs can serve as an important predictor of influenza vaccination uptake (Brewer et al., 2007; Shahrabani, Benzion & Yom Din, 2009; Shahrabani & Benzion, 2010; Tsutsui, Benzion & Shahrabani, 2012; Alsuwailem et al., 2023). During the COVID-19 pandemic, the HBM was used to assess the root causes of COVID-19 vaccine hesitancy in the Asia-Pacific region and beyond (Kabir et al., 2021; Alobaidi, 2021; Huynh et al., 2021; Zilhadia et al., 2022; Jian Ng et al., 2022; Shah et al., 2022; Seangpraw et al., 2022ba; Limbu, Gautam &

Pham, 2022b; Ghazy et al., 2022; Hidayana et al., 2022; Qin et al., 2022; Harutyunyan et al., 2023; Berger et al., 2023; Toledo-López et al., 2023).

In an earlier study, we interrogated COVID-19 vaccine intent in the Philippines by deploying a nationwide open-access online survey, two months *before* the rollout of the national vaccination program (Caple et al., 2022). Multivariable analysis of our data revealed then that HBM constructs were associated with vaccination intent in the Filipino population. In this follow-up study, we explored the influence of an ongoing vaccine rollout on vaccine intent by deploying a survey six months *after* the initial rollout of vaccines had begun throughout the Philippines. We wanted to determine if and how a vaccine rollout would predict public perceptions of the vaccines.

Results of the current study suggest that the presence of vaccines and vaccinated individuals in a Filipino community was associated with an increase in vaccination intent. When directly asked if individuals were more willing to receive a vaccination than six months prior, 92.26% agreed to some extent that they were indeed more willing. Finally, despite the changes in the numbers of respondents who were more open to the COVID-19 vaccines, there were no significant changes in the predictive power of the HBM constructs. For the most part, HBM factors that predicted vaccination intent in our earlier study also predicted vaccination intent in this current study.

Materials & Methods

Participants and Survey Design

The current study design was a cross-sectional, open-access, anonymous, web-based survey—developed using Qualtrics—conducted from August 14, 2021, to September 13, 2021. Our research team deployed an anonymous survey link via the social platforms of the University of Santo Tomas (UST) like Twitter and Facebook and university mailing lists including the UST School of Science and UST Student Council to distribute the survey. Participants were encouraged to distribute the survey link to their contacts throughout the country. The questionnaire was written in both Filipino and English. Responses used for data collection were limited to respondents who were at least 18 years old. An IRB-approved informed consent statement was included in the survey instrument to welcome respondents who had clicked on the anonymous survey link.

Survey Instrument

The survey consisted of questions and statements that assessed the following: (1) demographics, health status, and COVID-19 experience; (2) openness to a COVID-19 vaccine, and if participants received a COVID-19 vaccine; (3) perceived susceptibility to and severity of COVID-19; (4) perceived

benefits of a COVID-19 vaccine; and (5) barriers of a COVID-19 vaccine, and cues to action to obtain a COVID-19 vaccine.

Demographics, health status, and COVID-19 experience: Demographic information including age, sex, marital status, education level, occupation, and urban or rural location of residence were collected. Participants were also asked if they have an existing chronic condition, how they would rate their overall health, if they ever personally tested positive for COVID-19, to indicate if they know anyone who has tested positive for COVID-19, and if they know someone who has been vaccinated against COVID-19.

Openness to a COVID-19 vaccine and participant vaccination status: Openness to a COVID-19 vaccine was assessed using two items ("Would you take a COVID-19 vaccine?", "Are you more willing to take a COVID-19 vaccine now than you were six months ago?") on a five-point scale ranging from 1 = 'definitely no' to 5 = 'definitely yes'. Responses were further recoded into two distinct categories: vaccine resistance (responses included: 'definitely no', 'probably no', and 'unsure') and vaccine acceptance (responses included: 'probably yes' and 'definitely yes') for both scale items. Participant COVID-19 vaccination status was assessed using a one-item question ("Have you been vaccinated against COVID-19?"). Responses were additionally recoded into three distinct categories: not vaccinated, partially vaccinated, and fully vaccinated. Participants who indicated they had received either partial or full vaccination status were also asked to specify which vaccine they receive ("If you have been vaccinated against COVID-19, which vaccine did you receive?").

Perceived susceptibility to and severity of COVID-19: Health Belief Model (HBM) derived items were used to assess individual beliefs about a COVID-19 vaccine as well as individual beliefs about a COVID-19 vaccine prior to the national vaccine rollout. Questions posted to participants assessed perceived susceptibility of COVID-19 (three items), and perceived severity of COVID-19 (four items). All response items were on a four-point scale ranging from 'strongly disagree' to 'strongly agree'. For analysis purposes, all responses were coded as either *disagree* (responses included: 'strongly disagree' and 'disagree') or *agree* (responses included: 'strongly agree' and 'agree').

Perceived benefits of a COVID-19 vaccine: Perceived benefits were assessed using seven items, including statements that examined perceived benefits of a COVID-19 vaccine compared to before the national vaccine rollout. All response items were rated on a four-point scale ranging from 'strongly disagree' to 'strongly agree'. Similar to perceived susceptibility to and severity of COVID-19, all responses were coded as either *disagree* or *agree*.

Perceived barriers of a COVID-19 and cues to action: Perceived barriers surrounding a COVID-19 vaccine (e.g., "I worry about the possible side-effects of the COVID-19 vaccine"; "I worry about fake COVID-19 vaccines"). For analysis purposes, all responses were coded as either *disagree* or *agree*.

Additionally, cues to actions were assessed used eight items. All responses items were on a four-point scale ranging from 'strongly disagree' to 'strongly agree'. All responses were coded as either *disagree* (responses included: 'strongly disagree' and 'disagree') or *agree* (responses included: 'strongly agree' and 'agree').

Ethics Review & IRB Approval

Our study protocol (Protocol Number 21-026) was reviewed and approved by the Institutional Review Board of Providence College on January 15, 2021, with an amendment for the current study being reviewed and approved on July 15, 2021. Originally, we had sought ethical review at the University of Santo Tomas in the Philippines for our initial study but were advised by university authorities there to seek accelerated IRB approval in the United States due to the impact of the then ongoing- COVID-19 pandemic.

Statistical Analyses

All statistical analyses were conducted using Statistical Package for the Social Sciences (SPSS) version 28. A p-value of less than .05 was considered statistically significant. Frequency tables, charts, and proportions were used for data summarization—proportions and their respective 95% confidence intervals (CI) were calculated for each predictor variable. Participant responses to the one-item COVID-19 vaccine status question ("Have you been vaccinated against COVID-19?") was coded into two groups: not vaccinated and partially vaccinated/fully vaccinated responses. We ran univariate analyses followed by a multinomial logistic regression analysis, including all factors showing significance (p < .05), to determine which factors predicted individual COVID-19 vaccine intention. Only significant factors in the univariate analyses were included in the multinomial logistic regression analysis.

Results

Demographics

A total of 5,130 complete survey responses were received. Responses received represented participants with diverse demographics (Table 1) and from various regions of the Philippines (Figure 1). Demographic breakdowns were similar to our study of COVID-19 intent three months prior to the national vaccine rollout (Caple et al., 2022). However, the current study sample had a higher representation of younger adults aged 18 to 30 years old (67.3%). Additionally, the majority of participants identified as female (64.2%), single (75.7%), had obtained a college/university degree or above (77.2%), and lived in an urban area (79.7%). A small portion of our sample reported they have an

existing chronic condition (12.4%), while 11.9% of respondents reported having 'fair' health. Additionally, 64.3% of the sample reported knowing someone who has tested positive for COVID-19.

Health Beliefs

As displayed in Table 2, with regards to perceived susceptibility to COVID-19 vaccines, a portion of respondents (42.8%) agreed to some extent (which includes *strongly agreed/agreed*) there was a high chance of personally contracting COVID-19 in the next few months, which was not significantly different from the 31.5% reported previously (Caple et al., 2022). A significant majority of respondents (81.1%) also agreed to some extent that they were worried about the likelihood of personally contracting COVID-19. This was comparable to the number (84.1%) reported nine months earlier (Caple et al., 2022). Similarly, a large majority of respondents reported agreement to some extent when asked about their personal worry about the likelihood of someone in their family contracting COVID-19 (86.6%). A majority of participants reported COVID-19 is a serious disease with life-threatening complications (95.91%). A notable majority of respondents (92.7%) agreed to some extent that they were personally afraid of contracting COVID-19 and that they would personally be very sick if they contracted COVID-19 (72.1%). These were indistinguishable from the numbers, 93.1% and 75%, that reported previously from our survey completed nine months prior to this study (Caple et al., 2022).

With regards to the perceived benefits of vaccination, a majority of participants reported vaccination would decrease their chances of personally contracting COVID-19 (87.35%) as well as their worries about contracting COVID-19 (87.78%). While a majority of respondents reported they had worries about possible side-effects (68.1%), effectiveness (68.3%), safety (63.7%), and high cost (63.1%) of a COVID-19 vaccine, these numbers are overall lower compared to previous findings (Caple et al., 2022). Additionally, a majority of participants reported being worried about fake COVID-19 vaccines (88.5%).

Next, we assessed possible cues to action for COVID-19 vaccination in the course of a national vaccination campaign (Table 2). Our results suggest respondents' motivation to be immunized was predicted by whether they would be rewarded with perks offered by the private sector including discounts at restaurants and other service industries (80.0% agreed to some extent that they were motivated by this incentive), and rewarded with more freedom to move around and meet other people (78.8% agreed to some extent that they were motivated by this incentive). They were less enthusiastic about rewards that gave them freedom to remove masks in the company of fellow vaccinated individuals (58.69% *strongly agree/agree* that they were motivated by this incentive). Respondents also revealed that hearing from fully vaccinated people (87.5% agreed to some extent) and seeing family members and friends get

vaccinated (87.5% agreed to some extent) were predictive factors that would encourage one to seek vaccination.

Finally, as shown in Table 3, respondents were also prompted with statements about COVID-19 perceptions compared to six months prior. A majority of participants agreed to some extent (73%) that their chance of personally contracting COVID-19 was lower compared to six months prior. Similarly, 67.2% of participants agreed to some extent that they were less worried about the likelihood of contracting COVID-19. A majority (63.2%) also revealed that were also less worried about the likelihood of someone in their family contracting COVID-19 compared to six months prior.

With regards to perceived barriers to vaccination, majorities of respondents revealed that they were less worried about the possible side-effects of the vaccines (78.8%), less worried about vaccine effectiveness (76.7%), less worried about vaccine safety (78.64%), less worried about fake vaccines (64.4%), and less worried about vaccine testing (70.57%) compared to six months prior. Overall, our respondents were less worried about every aspect of possible concern associated with COVID-19 vaccination.

COVID-19 Vaccination Intent

Survey participants were asked whether they had been vaccinated against COVID-19, which was recoded into three groups: not vaccinated (16.7%), partially vaccinated (28.8%), or fully vaccinated (54.5%) (Figure 2A). We also included a specific question to interrogate vaccination intent. As shown in Figure 2B, when responses to the question, "Would you take a COVID-19 vaccine?", were recoded into two groups, a majority of respondents (95.3%) indicated vaccine acceptance (responses included 'probably yes' and 'definitely yes'). Only 242 of the 5,130 respondents (4.7%) were vaccine resistant (responses included were 'definitely no', 'probably no', and 'unsure'). A similar pattern emerged when participants were asked if they were more willing to take a COVID-19 vaccine now as compared to six months prior, with a vast majority (92.3%) identifying as open (responses included 'definitely yes' or 'probably yes') to the COVID-19 vaccines (Table 3).

Table 4 shows the univariate and multinomial analyses of factors associated with vaccination status by demographics and HBM constructs. Univariate analyses were conducted examining vaccine resistance and vaccine acceptance as a binary—not vaccinated and partially/fully vaccinated respectively. A total of 4,274 of respondents (83.3%) reported they were either partially (n = 1476, 28.8%) or fully vaccinated (n = 2798, 54.5%); demonstrating they were vaccine acceptant—while 856 of respondents (16.7%) reported they were not vaccinated suggesting that they were vaccine resistant. The majority of respondents who identified as being in the 18-30 age range, also reported being either partially or fully vaccinated (n = 2752, 79.7%).

Univariate analyses showed a high proportion of participants who identified as married (90.8%) noted they were either partially or fully vaccinated than compared to 80.9% of partially or fully vaccinated, single respondents. This association, however, was not significant in the multinomial analysis. When examining participant occupation, a higher proportion of respondents that identified as having vaccine acceptance (i.e., either partially or fully vaccinated) also identified as professional/white collar workers (91.2%). Additionally, significant differences were noted in vaccination status by geographic location, where individuals in urban locations (85.3%) reported being either partially or fully vaccinated, while 75.4% of participants in rural locations noted they were either partially or fully vaccination.

Based on demographics, univariate analyses revealed that females have greater odds of being either partially or fully vaccinated (OR = .653, 95% CI [.556, .768]) than males—a finding that also was statistically significant in the multinomial analysis. Furthermore, being a blue collar worker (OR = 1.918, 95% CI [1.294, 2.843]), a professional or white collar worker (OR = 1.694, 95% CI [1.195, 2.400]), a student (OR = .320, 95% CI [.233, .439]), having an existing chronic condition (OR = 1.425, 95% CI [1.115, 1.822]), and perceived overall health reported as 'fair', 'poor', or 'very poor' (OR = .494, 95% CI [.395, .617]) and as 'good' (OR = .712, 95% CI [.604, .840]), were significant predictors of vaccination status.

Most of the HBM factors were significantly associated with vaccination status. We observed a similar pattern in our previous study (Caple et al., 2022). While 50.4% of respondents noted they were worried COVID-19 vaccines would make them sick, disagreement with that notion (OR = 2.683, 95% CI [2.028, 3.551]) was the strongest predictor of vaccination status when comparing not vaccinated participants to fully vaccinated participants. Individual disagreement with the belief that COVID-19 vaccination would decrease participant worries about personally contracting COVID-19 for not vaccinated compared to partially vaccinated respondents (OR = 2.066, 95% CI [1.391, 3.068]) and for not vaccinated compared to fully vaccinated respondents (OR = 1.819, 95% CI [1.246, 2.654]) was also a strong predictor of vaccination status.

Finally, in Table 5, we compared the HBM constructs that were associated with vaccination intent in our earlier study (Caple et al., 2022), with the same constructs in this study, to determine if they had changed. Our analysis revealed that beliefs around the perceived susceptibility to getting COVID-19--whether the respondent believed that he or she would become sick in the next six months--which were predictive of vaccination intent in our earlier study, were now not able to forecast a respondent's openness to the vaccines. Strikingly, two more belief constructs regarding the perceived barriers of expensive vaccines and of fake vaccines, also ceased being predictive in this study when they were predictive in our earlier report.

Discussion

How does an ongoing vaccination campaign influence vaccination intent, especially during a pandemic? How do the presence of vaccines and vaccinated individuals shape and alter, if at all, vaccination intent? In our previous study, completed two months *before* the rollout of the national COVID-19 vaccination campaign in the Philippines, we showed that Health Belief Model (HBM) constructs were significantly associated with vaccination intent in the archipelago (Caple et al., 2022). Perceptions of high susceptibility, high severity, and significant benefits were all good predictors for vaccination intent. We also found that external cues to action were important.

Here, in a follow-up study completed six months *after* the beginning of the national vaccination campaign, we discovered that the presence of vaccines and vaccinated individuals in a Filipino community predicted vaccine intent. Most significantly, in our earlier study, 62.5% of our respondents indicated that they were willing to be vaccinated when the vaccines became available (Caple et al., 2022). This study revealed that at the time of our survey, 83.3% of our respondents had already begun the vaccination series (Figure 2). Strikingly, even more individuals (95.3%) indicated that they were willing to be vaccinated. When directly asked if they were more willing to receive a vaccination than six months prior, 92.3% agreed to some extent that they were indeed more willing. This significant increase in vaccination intent during a comparable time period has also been observed by others in the Philippines (Macaraan, 2021).

How do we explain the difference in numbers between those persons who were vaccinated (83.3%) and those who said that they were willing to be vaccinated (95.3%)? There are at least two possibilities. First, there could have been a group of individuals who were still waiting for vaccine access: These Filipinos were willing to be vaccinated, but there were no vaccines for them at the time of the survey. This scenario is likely because a significant minority (20.33%) of our respondents lived in rural areas. At the time of our survey, six months into the national vaccination campaign, the Philippines was still prioritizing individuals in the urban areas of the country in a bid to build a wall of immunization around cities that would prevent the transmission of the SARS-CoV2 virus into the rural countryside. This public policy was named the NCR+8 vaccination strategy (Casinas, 2021; Almajose et al., 2021).

Second, there could have been a group of individuals who were waiting to be vaccinated, allowing others to be vaccinated ahead of them, because of the shortage of vaccines. Though there were anecdotal stories of altruistic persons circulating on social media during the earliest months of the national vaccination campaign in the Philippines, they have never been confirmed.

Most notably, our study reveals that the presence of vaccines and vaccinated individuals in a Filipino community predicts vaccination intent (Figure 2). Note that it cannot simply be the presence of vaccines or vaccinated individuals that predicted vaccination intent, because the global vaccination

campaign had already begun at the time of our first study, unleashing numerous posts and reports on Filipino social media platforms of vaccinated citizens in the Global North. As our study revealed, what is significant for predicting vaccination intent is not whether or not the respondents knew that others were getting vaccinated. Rather what was most important is that our respondents personally knew loved ones and friends who had been vaccinated (Table 2). These were the most significant cues of action to COVID-19 vaccination in our study. Similarly, a COVID-19 study in Taiwan has also reported the effect of positive word-of-mouth testimony on vaccination intent (Wu & Chiang, 2023).

What is the significance of this finding? It suggests that Filipinos want to hear and to trust the testimonies of their *kababayans* ("fellow Filipinos, townmates, or countrymen"). Thus, future vaccination campaigns in the Philippines should showcase Filipinos who have been vaccinated, especially those who live in the same *barangay* as the target vaccination population. A *barangay* is the smallest administrative division in the Philippines and is the native Filipino term for a village, a district, or ward. The more familiar the testimony of the vaccinated *kababayan*, the better.

As for additional cues to action for COVID-19 vaccination, our study revealed that our Filipino respondents believed that concrete rewards for immunization like the private sector's *bakuna* benefits program which gave immunized individuals exclusive perks, discounts, and even freebies, would motivate them to get immunized. These *bakuna* benefits included free menu items, free upgrades on drink sizes, and "buy one, get one" promotional items (Ignacio, 2021). Other vaccination perks including more freedom to mingle with others also motivated our respondents (78.8% agreed to some extent that this would encourage them) though incentives that would give them more freedom to remove masks would not motivate them as much (58.69% agreed to some extent that this would encourage them). Why not masks? Surveys have shown that Filipinos value masking and that significant numbers of them intended to continue wearing their masks even years after the official end of the pandemic (De Vera-Ruiz, 2022).

Next, despite the changes in the numbers of respondents who became more open to the COVID-19 vaccines after the rollout of the national vaccination campaign, there were no significant changes in the predictive power of the HBM constructs during the same time period. For the most part, HBM factors that predicted vaccination intent in our earlier study also predicted vaccination intent in this current study (Table 5). Only one, age of the respondent, became predictive when it was not so in our previous work. One simple explanation for this is that the vaccination rollout in the Philippines prioritized the elderly. As such, at the time of our survey, it was significantly more likely for a senior citizen to be vaccinated than for a young adult to have been immunized (Rappler.com, 2021). This was not the case prior to the arrival of the vaccines.

Four of the twenty-four HBM constructs assessed in the current study no longer predicted vaccine intent once the national vaccination strategy was underway (Table 5). Two involved a respondent's

perceived susceptibility to COVID-19, and two were related to one's worries regarding the cost and the authenticity of the COVID-19 vaccines. In our earlier study, respondents who believed that there was a high risk that they would get infected with SARS-CoV2 were more likely to be open to the COVID-19 vaccines (Caple et al., 2022). In this study, perceived susceptibility ceased to be associated with a positive vaccine intent. One possible explanation is that a large number of our respondents in this study (67.2%) also reported that they had simply become less worried about COVID-19. These results suggest that fear of severe illness simply became less of a concern among our respondents as the vaccines were rolled out.

Similarly, worries surrounding the cost and authenticity of the COVID-19 vaccines lost their ability to predict vaccination intent. This should not be surprising since the availability of free and legitimate vaccines in the Philippines would have made these concerns moot in the minds of many Filipinos. Of note, the national government launched a widespread public advisory campaign against fake COVID-19 vaccines throughout the rollout of the national vaccination campaign. It was quick to announce investigations of companies purporting to distribute fake vaccines (CNN, 2021).

Finally, our study has several limitations. As we already noted with our earlier study, the use of an open-access online survey may result in sampling bias so we cannot generalize our findings to the entire Filipino population (Wyatt, 2000; Eysenbach & Wyatt, 2002). Furthermore, the responses were based on self-report and may be subject to self-reporting bias and a tendency to report socially desirable responses especially in a strongly collectivist society like the Philippines. Despite these shortcomings, we believe that our findings will provide insights to support future vaccine rollouts in the Philippines by helping public health authorities to understand vaccine demand and vaccine hesitancy in the country.

373 References

- 374 Alfonso C., Dayrit M., Mendoza R., Ong M. 2021. From Dengvaxia to Sinovac: Vaccine Hesitancy in the
- 375 Philippines. *The Diplomat*.
- 376 Almajose APL., White A., Diego MCE., Lazaro R., Nicanor Austriaco OP. 2021. 2SIR-VD Model to
- 377 Compare Idealized COVID-19 Vaccine Distribution Strategies in the Philippines. *Philippine*
- *Journal of Science* 150:1765–1778. DOI: 10.56899/150.6B.13.
- 379 Alobaidi S. 2021. Predictors of Intent to Receive the COVID-19 Vaccination Among the Population in
- the Kingdom of Saudi Arabia: A Survey Study. *Journal of multidisciplinary healthcare* 14:1119–
- 381 1128. DOI: 10.2147/JMDH.S306654.
- 382 Alsuwailem SI., Okmi EA., Alkhaldi EH., Almutairi KS., Alshamari WK. 2023. Determinants of Flu
- Vaccine Uptake Among the General Population in Saudi Arabia: A Study Based on the Health
- 384 Belief Model. *Cureus* 15:e41277. DOI: 10.7759/cureus.41277.
- Berger C., Ben-Shalom U., Tarant Z., Longo J., DeDonno M. 2023. The Influence of the Health Belief
- Model on the Decision to Get the COVID-19 Vaccine: An International Survey Study of College
- 387 Students. *Inquiry*: a journal of medical care organization, provision and financing
- 388 60:469580231164229. DOI: 10.1177/00469580231164229.
- 389 Brewer NT., Chapman GB., Gibbons FX., Gerrard M., McCaul KD., Weinstein ND. 2007. Meta-analysis
- of the relationship between risk perception and health behavior: The example of vaccination. *Health*
- 391 *Psychology* 26:136–145. DOI: 10.1037/0278-6133.26.2.136.
- 392 Caple A., Dimaano A., Sagolili MM., Uy AA., Aguirre PM., Alano DL., Camaya GS., Ciriaco BJ., Clavo
- PJM., Cuyugan D., Fermo CFG., Lanete PJ., La Torre AJ., Loteyro T., Lua RM., Manansala NG.,
- Mosquito RW., Octaviano A., Orfanel AE., Pascual GM., Sale AJ., Tendenilla SL., Trinidad MSL.,
- 395 Trinidad NJ., Verano DL., Austriaco N. 2022. Interrogating COVID-19 vaccine intent in the
- Philippines with a nationwide open-access online survey. *PeerJ* 10:e12887. DOI:
- 397 10.7717/PEERJ.12887.
- 398 Casinas JA. 2021.12 million workers from "NCR+8" prioritized in A4 COVID-19 vaccination DOH.
- 399 Available at https://mb.com.ph/2021/06/07/12-million-workers-from-ncr8-prioritized-in-a4-covid-
- 400 *19-vaccination-doh/* (accessed October 15, 2023).
- Cassell CH., Raghunathan PL., Henao O., Pappas-DeLuca KA., Rémy WL., Dokubo EK., Merrill RD.,
- Marston BJ. 2022. Global Responses to the COVID-19 Pandemic. *Emerging Infectious Diseases*
- 403 28:S4–S7. DOI: 10.3201/EID2813.221733.
- Chen IH., Hsu SM., Wu JSJ., Wang YT., Lin YK., Chung MH., Huang PH., Miao NF. 2019.
- Determinants of nurses' willingness to receive vaccines: Application of the health belief model.
- 406 *Journal of Clinical Nursing* 28:3430–3440. DOI: 10.1111/jocn.14934.

- 407 CNN. 2021.PH gov't investigating companies behind spread of fake vaccines. Available at
- 408 https://www.cnnphilippines.com/news/2021/3/30/Fake-vaccines-Philippines.html (accessed October
- 409 15, 2023).
- 410 Donadiki EM., Jiménez-García R., Hernández-Barrera V., Sourtzi P., Carrasco-Garrido P., López de
- Andrés A., Jimenez-Trujillo I., Velonakis EG. 2014. Health Belief Model applied to non-
- 412 compliance with HPV vaccine among female university students. *Public Health* 128:268–273. DOI:
- 413 10.1016/j.puhe.2013.12.004.
- 414 Eysenbach G., Wyatt J. 2002. Using the Internet for surveys and health research. *Journal of Medical*
- 415 *Internet Research* 4:76–94. DOI: 10.2196/JMIR.4.2.E13.
- 416 Fatima K., Syed NI. 2018. Dengvaxia controversy: impact on vaccine hesitancy. *Journal of Global*
- 417 *Health* 8. DOI: 10.7189/JOGH.08-020312.
- 418 Ghazy RM., Abdou MS., Awaidy S., Sallam M., Elbarazi I., Youssef N., Fiidow OA., Mehdad S.,
- Hussein MF., Adam MF., Abdullah FSA., Rebai WK., Raad EB., Hussein M., Shehata SF., Ismail
- 420 II., Salam AA., Samhouri D. 2022. Acceptance of COVID-19 Vaccine Booster Doses Using the
- Health Belief Model: A Cross-Sectional Study in Low-Middle- and High-Income Countries of the
- East Mediterranean Region. *International Journal of Environmental Research and Public Health* 19.
- 423 DOI: 10.3390/IJERPH191912136.
- van Ginneken E., Webb E., Maresso A., Cylus J. 2022. Lessons learned from the COVID-19 pandemic.
- 425 *Health Policy* 126:348–354. DOI: 10.1016/j.healthpol.2022.04.004.
- 426 Guan WJ., Ni ZY., Hu Y., Liang WH., Ou CQ., He JX., Liu L., Shan H., Lei CL., Hui DSC., Du B., Li
- LJ., Zeng G., Yuen KY., Chen RC., Tang CL., Wang T., Chen PY., Xiang J., Li SY., Wang JL.,
- Liang ZJ., Peng YX., Wei L., Liu Y., Hu YH., Peng P., Wang JM., Liu JY., Chen Z., Li G., Zheng
- 429 ZJ., Qiu SQ., Luo J., Ye CJ., Zhu SY., Zhong NS. 2020. Clinical Characteristics of Coronavirus
- Disease 2019 in China. *The New England journal of medicine* 382:1708–1720. DOI:
- 431 10.1056/NEJMoa2002032.
- Harutyunyan T., Sargsyan Z., Sahakyan S., Chiloyan A., Melkom Melkomian D., Khachadourian V.
- 433 2023. Predicting COVID-19 vaccination uptake in Armenia using the Health Belief Model: Results
- from a nationwide survey. *Human Vaccines and Immunotherapeutics* 19. DOI:
- 435 10.1080/21645515.2023.2165383.
- 436 Hidayana I., Amir S., Pelupessy DC., Rahvenia Z. 2022. Using a health belief model to assess COVID-19
- vaccine intention and hesitancy in Jakarta, Indonesia. *PLOS global public health* 2:e0000934. DOI:
- 438 10.1371/journal.pgph.0000934.
- 439 Hu Y., Wang Y., Liang H., Chen Y. 2017. Seasonal influenza vaccine acceptance among pregnant
- women in zhejiang province, china: Evidence based on health belief model. *International Journal of*

- 441 Environmental Research and Public Health 14. DOI: 10.3390/ijerph14121551. 442 Huynh G., Nguyen T Van., Nguyen DD., Lam QM., Pham TN., Nguyen HTN. 2021. Knowledge About 443 COVID-19, Beliefs and Vaccination Acceptance Against COVID-19 Among High-Risk People in 444 Ho Chi Minh City, Vietnam. *Infection and drug resistance* 14:1773–1780. DOI: 445 10.2147/IDR.S308446. 446 Ignacio R. 2021. "Bakuna" benefits: Perks for COVID-19-vaccinated foodies. Available at 447 https://www.philstar.com/lifestyle/food-and-leisure/2021/07/02/2109767/bakuna-benefits-perks-448 covid-19-vaccinatedfoodies (accessed October 15, 2023). 449 Inter-Agency Task Force for the Management of Emerging Infectious Disease. 2021. The Philippine 450 National COVID-19 Vaccination Deployment Plan | Department of Health website. 451 Jian Ng JW., Vaithilingam S., Nair M., Hwang LA., Musa KI. 2022. Key predictors of COVID-19 452 vaccine hesitancy in Malaysia: An integrated framework. *PLoS ONE* 17. DOI: 453 10.1371/journal.pone.0268926. 454 Kabir R., Mahmud I., Chowdhury MTH., Vinnakota D., Jahan SS., Siddika N., Isha SN., Nath SK., 455 Hoque Apu E. 2021. COVID-19 Vaccination Intent and Willingness to Pay in Bangladesh: A Cross-456 Sectional Study. Vaccines 9. DOI: 10.3390/vaccines9050416. 457 Limbu YB., Gautam RK. 2023. How Well the Constructs of Health Belief Model Predict Vaccination 458 Intention: A Systematic Review on COVID-19 Primary Series and Booster Vaccines. Vaccines 11. 459 DOI: 10.3390/VACCINES11040816. 460 Limbu YB., Gautam RK., Pham L. 2022a. The Health Belief Model Applied to COVID-19 Vaccine 461 Hesitancy: A Systematic Review. Vaccines 10. DOI: 10.3390/VACCINES10060973. 462 Limbu YB., Gautam RK., Pham L. 2022b. The Health Belief Model Applied to COVID-19 Vaccine 463 Hesitancy: A Systematic Review. Vaccines 10. DOI: 10.3390/vaccines10060973. 464 Macaraan WER. 2021. From hesitancy to confidence: Filipinos' shifting attitude toward COVID-19 465 vaccination program. Journal of public health (Oxford, England). DOI: 10.1093/pubmed/fdab233. 466 Qin C., Yan W., Du M., Liu Q., Tao L., Liu M., Liu J. 2022. Acceptance of the COVID-19 vaccine 467 booster dose and associated factors among the elderly in China based on the health belief model 468 (HBM): A national cross-sectional study. Frontiers in Public Health 10. DOI: 469 10.3389/fpubh.2022.986916. 470 Rappler.com. 2021.FAST FACTS: Prioritized groups, guidelines for COVID-19 vaccination. Available at 471 https://www.rappler.com/newsbreak/iq/things-to-know-prioritized-groups-guidelines-covid-19-
- Rappuoli R., Pizza M., Del Giudice G., De Gregorio E. 2014. Vaccines, new opportunities for a new society. *Proceedings of the National Academy of Sciences of the United States of America*


472

vaccination-philippines/


- 475 111:12288–12293. DOI: 10.1073/pnas.1402981111.
- 476 Reis J., Faou A Le., Buguet A., Sandner G., Spencer P. 2022. Covid-19: Early Cases and Disease Spread.
- 477 *Annals of Global Health* 88. DOI: 10.5334/aogh.3776.
- 478 Reñosa MDC., Wachinger J., Bärnighausen K., Endoma V., Landicho-Guevarra J., Landicho J., Bravo
- 479 TA., Aligato M., McMahon SA. 2022. Misinformation, infighting, backlash, and an 'endless'
- recovery; policymakers recount challenges and mitigating measures after a vaccine scare in the
- 481 Philippines. *Global Health Action* 15. DOI: 10.1080/16549716.2022.2077536.
- Rosenstock IM., Strecher VJ., Becker MH. 1988. Social Learning Theory and the Health Belief Model.
- 483 *Health Education & Behavior* 15:175–183. DOI: 10.1177/109019818801500203.
- 484 Sachs JD., Karim SSA., Aknin L., Allen J., Brosbøl K., Colombo F., Barron GC., Espinosa MF., Gaspar
- V., Gaviria A., Haines A., Hotez PJ., Koundouri P., Bascuñán FL., Lee JK., Pate MA., Ramos G.,
- Reddy KS., Serageldin I., Thwaites J., Vike-Freiberga V., Wang C., Were MK., Xue L., Bahadur C.,
- Bottazzi ME., Bullen C., Laryea-Adjei G., Amor Y Ben., Karadag O., Lafortune G., Torres E.,
- Barredo L., Bartels JGE., Joshi N., Hellard M., Huynh UK., Khandelwal S., Lazarus J V., Michie S.
- 489 2022. The Lancet Commission on lessons for the future from the COVID-19 pandemic. *The Lancet*
- 490 400:1224–1280. DOI: 10.1016/S0140-6736(22)01585-9.
- 491 Sathyanarayana SD., Fernandes SD., Castelino LJ., Vadakkepushpakath AN., Shriram RG. 2020.
- Vaccines in the United States: a systematic review on history of evolution, regulations, licensing,
- and future challenges. *Clinical and experimental vaccine research* 9:69–75. DOI:
- 494 10.7774/cevr.2020.9.2.69.
- Seangpraw K., Pothisa T., Boonyathee S., Ong-Artborirak P., Tonchoy P., Kantow S., Auttama N.,
- Choowanthanapakorn M. 2022a. Using the Health Belief Model to Predict Vaccination Intention
- Among COVID-19 Unvaccinated People in Thai Communities. Frontiers in Medicine 9. DOI:
- 498 10.3389/FMED.2022.890503.
- 499 Seangpraw K., Pothisa T., Boonyathee S., Ong-Artborirak P., Tonchoy P., Kantow S., Auttama N.,
- Choowanthanapakorn M. 2022b. Using the Health Belief Model to Predict Vaccination Intention
- Among COVID-19 Unvaccinated People in Thai Communities. Frontiers in medicine 9:890503.
- DOI: 10.3389/fmed.2022.890503.
- 503 Shah S., Gui H., Chua PEY., Tan JY (Benjamin)., Suen LK ping., Chan SW chi., Pang J. 2022. Factors
- associated with COVID-19 vaccination intent in Singapore, Australia and Hong Kong. Vaccine
- 505 40:2949–2959. DOI: 10.1016/J.VACCINE.2022.03.062.
- 506 Shahrabani S., Benzion U. 2010. Workplace vaccination and other factors impacting influenza
- vaccination decision among employees in Israel. *International Journal of Environmental Research*
- *and Public Health* 7:853–869. DOI: 10.3390/ijerph7030853.

- 509 Shahrabani S., Benzion U., Yom Din G. 2009. Factors affecting nurses' decision to get the flu vaccine.
- *European Journal of Health Economics* 10:227–231. DOI: 10.1007/s10198-008-0124-3.
- 511 Siddiqui S., Alhamdi HWS., Alghamdi HA. 2022. Recent Chronology of COVID-19 Pandemic. Frontiers
- *in Public Health* 10. DOI: 10.3389/fpubh.2022.778037.
- 513 Smith N. 2018. Philippines immunisation rates plummet amid Dengue vaccination scare. *The Telegraph*.
- Teitler-Regev S., Shahrabani S., Benzion U. 2011. Factors Affecting Intention among Students to Be
- Vaccinated against A/H1N1 Influenza: A Health Belief Model Approach. *Advances in Preventive*
- 516 *Medicine* 2011:1–8. DOI: 10.4061/2011/353207.
- Toledo-López A., Leyva-Hernández SN., Jiménez-Castañeda JC., Avendaño-Rito MDC. 2023.
- Determinants for COVID-19 vaccination intention in Mexico. *Heliyon* 9:e18079. DOI:
- 519 10.1016/j.heliyon.2023.e18079.
- Tsutsui Y., Benzion U., Shahrabani S. 2012. Economic and behavioral factors in an individual's decision
- to take the influenza vaccination in Japan. *Journal of Socio-Economics* 41:594–602. DOI:
- 522 10.1016/j.socec.2012.05.001.
- 523 De Vera-Ruiz E. 2022.Most Filipinos still prefer to wear face masks to fight off Covid-19 OCTA
- 524 survey. Available at https://mb.com.ph/2022/09/13/most-filipinos-still-prefer-to-wear-face-masks-
- *to-fight-off-covid-19-octa-survey/* (accessed October 15, 2023).
- Wu SW., Chiang PY. 2023. Exploring the Moderating Effect of Positive and Negative Word-of-Mouth on
- the Relationship between Health Belief Model and the Willingness to Receive COVID-19 Vaccine.
- 528 *Vaccines* 11. DOI: 10.3390/VACCINES11061027.
- Wyatt JC. 2000. When to use web-based surveys. Journal of the American Medical Informatics
- Association 7:426–430. DOI: 10.1136/jamia.2000.0070426.
- Xie J., Tong Z., Guan X., Du B., Qiu H. 2020. Clinical Characteristics of Patients Who Died of
- Coronavirus Disease 2019 in China. *JAMA network open* 3:e205619. DOI:
- 533 10.1001/jamanetworkopen.2020.5619.
- Zhu N., Zhang D., Wang W., al. et. 2020. A novel coronavirus from patients with pneumonia in China,
- 535 2019. *N Engl J Med* 382:727–733.
- 536 Zilhadia Z., Ariyanti F., Nurmansyah MI., Iriani DU., Dwirahmadi F. 2022. Factors Associated with
- 537 COVID-19 Vaccination Acceptance Among Muslim High School Students in Jakarta Metropolitan
- Area, Indonesia. *Journal of Multidisciplinary Healthcare* 15:2341–2352. DOI:
- 539 10.2147/JMDH.S380171.

REGION OF THE PHILIPPINES	Number of Responses (n)	Percent of Responses (%)
BARMM	8	0.2
CAR	25	0.5
NCR	2,357	45.9
Region I	58	1.1
Region II	30	0.6
Region III	1,261	24.6
Region IV-A	1,007	19.6
Region IV-B	17	0.3
Region IX	9	0.2
Region V	22	0.4
Region VI	62	1.2
Region VII	35	0.7
Region VIII	11	0.2
Region X	46	0.9
Region XI	135	2.6
Region XII	47	0.9
Region XIII	0	0
TOTAL	5,130	100







	F	lave you been vaccir	nated against COVII	D-19?
	Overall N (%)	Not Vaccinated n = 856 (16.7%)	Partially Vaccinated n = 1476 (28.8%)	Fully Vaccinated n = 2798 (54.5%)
Demographics				
Age group (years)				
18–30	3453 (67.31)	701 (20.3)	1223 (35.4)	1529 (44.3)
31–40	521 (10.16)	41 (7.9)	122 (23.4)	358 (68.7)
41–50	451 (8.79)	52 (11.5)	65 (14.4)	334 (74.1)
51–99	705 (13.74)	62 (8.8)	66 (9.4)	577 (81.8)
Gender				
Female	3293 (64.19)	616 (18.7)	890 (27)	1787 (54.3)
Male	1837 (35.81)	240 (13.1)	586 (31.9)	1011 (55)
Marital Status				
Single	3883 (75.69)	741 (19.1)	1292 (33.3)	1850 (47.6)
Married	1247 (24.31)	115 (9.2)	184 (14.8)	948 (76)
Highest Education Level				
Elementary school or below	3 (.06)	1 (33.3)	2 (66.7)	
Junior high school	132 (2.57)	14 (10.6)	55 (41.7)	63 (47.7)
Senior high school	1036 (20.19)	220 (21.2)	412 (39.8)	404 (39)
College/university of above	3959 (77.17)	621 (15.7)	1007 (25.4)	2331 (58.9)
Occupation				
Blue collar worker	878 (17.12)	64 (7.3)	397 (45.2)	417 (47.5)
Profession/white collar worker	1627 (31.72)	133 (8.2)	328 (20.2)	1166 (71.7)
Self-employed	541 (10.55)	62 (11.5)	209 (38.6)	270 (49.9)
Student	1710 (33.33)	548 (32)	465 (27.2)	697 (40.8)
Housewife/retired/unemployed/other	374 (7.29)	49 (13.1)	77 (20.6)	248 (66.3)
Location				
Rural	1043 (20.33)	257 (24.6)	334 (32)	452 (43.3)
Urban	4087 (79.67)	599 (14.7)	1142 (27.9)	2346 (57.4)

Experience with COVID-19				
Ever tested positive for COVID-19				
Yes	334 (6.51)	48 (14.4)	72 (21.6)	214 (64.1)
No	4796 (93.49)	808 (16.8)	1404 (29.3)	2584 (53.9)
Know anyone who has tested positive for COVID-19				
Yes	3299 (64.31)	617 (18.7)	649 (19.7)	2033 (61.6)
No	1831 (35.69)	239 (13.1)	827 (45.2)	765 (41.8)
Health Characteristics				
Have an existing chronic condition				
Yes	635 (12.38)	81 (12.8)	61 (9.6)	493 (77.6)
No	4495 (87.62)	775 (17.2)	1415 (31.5)	2305 (51.3)
Perceived overall health				
Very good	2021 (39.4)	268 (13.3)	703 (34.8)	1050 (52)
Good	2466 (48.07)	436 (17.7)	650 (26.4)	1380 (56)
Fair/Poor/Very poor	643 (12.53)	152 (23.6)	123 (19.1)	368 (57.2)

	Have you been vaccinated against COVID-19?					
	Overall N (%)	Not Vaccinated n = 856 (16.7%)	Partially Vaccinated & Fully Vaccinated n = 4274 (83.3%)			
Demographics						
Age group (years)						
18–30	3453 (67.31)	701 (20.3)	2752 (79.7)			
31–40	521 (10.16)	41 (7.9)	480 (92.1)			
41–50	451 (8.79)	52 (11.5)	399 (88.5)			
51–99	705 (13.74)	62 (8.8)	643 (91.2)			
Gender						
Female	3293 (64.19)	616 (18.7)	2677 (81.3)			
Male	1837 (35.81)	240 (13.1)	1597 (86.9)			
Marital Status						
Single	3883 (75.69)	741 (19.1)	3142 (80.9)			
Married	1247 (24.31)	115 (9.2)	1132 (90.8)			
Highest Education Level						
Elementary school or below	3 (.06)	1 (33.3)	2 (66.7)			
Junior high school	132 (2.57)	14 (10.6)	118 (89.4)			
Senior high school	1036 (20.19)	220 (21.2)	816 (78.8)			
College/university of above	3959 (77.17)	621 (15.7)	3338 (84.3)			
Occupation						
Blue collar worker	878 (17.12)	64 (7.3)	814 (92.7)			
Profession/white collar worker	1627 (31.72)	133 (8.2)	1494 (91.8)			
Self-employed	541 (10.55)	62 (11.5)	479 (88.5)			
Student	1710 (33.33)	548 (32)	1162 (68)			
Housewife/retired/unemployed/other	374 (7.29)	49 (13.1)	325 (86.9)			
Location						
Rural	1043 (20.33)	257 (24.6)	786 (75.4)			
Urban	4087 (79.67)	599 (14.7)	3488 (85.3)			

Experience with COVID-19			
Ever tested positive for COVID-19			
Yes	334 (6.51)	48 (14.4)	286 (85.6)
No	4796 (93.49)	808 (16.8)	3988 (83.2)
Know anyone who has tested positive for COVID-19			
Yes	3299 (64.31)	617 (18.7)	2682 (81.3)
No	1831 (35.69)	239 (13.1)	1592 (86.9)
Health Characteristics			
Have an existing chronic condition			
Yes	635 (12.38)	81 (12.8)	554 (87.2)
No	4495 (87.62)	775 (17.2)	3720 (82.8)
Perceived overall health			
Very good	2021 (39.4)	268 (13.3)	1753 (86.7)
Good	2466 (48.07)	436 (17.7)	2030 (82.3)
Fair/Poor/Very poor	643 (12.53)	152 (23.6)	491 (76.4)

Perceived Susceptibility to COVID-19	% Strongly Agree/Agree (This Study)	% Strongly Agree/Agree (Caple et al., 2022)
Chance of getting COVID-19 in the next few months is high	42.81	31.5
Worried about the likelihood of getting COVID-19	81.05	84.1
Worried about the likelihood of someone in my family getting COVID-19	86.61	NA
Perceived Severity of COVID-19		
COVID-19 is a serious disease with life-threatening complications	95.91	96.3
I will be very sick if I get COVID-19	72.12	75
I am afraid of getting COVID-19	92.73	93.1
I am afraid that someone in my family will get COVID-19	95.59	NA
Perceived Benefits of Vaccination		
Vaccination will decrease my chances of getting COVID-19	87.35	88.1
Vaccination will decrease my worries about catching COVID-19	87.78	84.5
Perceived Barriers to Vaccination		
I worry about the possible side-effects of the COVID-19 vaccine	68.13	89.6
I worry about the effectiveness of the COVID-19 vaccine	68.32	87.1
I worry about the safety of the COVID-19 vaccine	63.7	88.8
I worry about the high cost of the COVID-19 vaccine	63.14	78
I worry about fake COVID-19 vaccines	88.52	97.4
I worry that the COVID-19 vaccines were made too quickly	59.57	NA
I worry that the COVID-19 vaccines were not tested properly	55.13	NA
I worry that the COVID-19 vaccines will make me sick	50.37	68.4
I worry that the COVID-19 vaccines will not be effective against new variants of the virus	75.48	80
Cues to Action for COVID-19 Vaccination		
Rewarding fully vaccinated individuals with bakuna benefits would encourage me to get vaccinated against COVID-19	79.96	
Rewarding fully vaccinated individuals with more freedom to move around and to meet with people would encourage me to get vaccinated against COVID-19	78.75	
Rewarding fully vaccinated individuals by allowing them to remove masks in the company of other fully vaccinated individuals would	58.69	

encourage me to get vaccinated against COVID- 19		
Testimonies from fully vaccinated people have	87.5	
encouraged me to get vaccinated		
Seeing my family members and friends get	90.14	
vaccinated has encouraged me to get the vaccine		
Seeing famous people get vaccinated on social	62.83	
media has encouraged me to get the vaccine		

TABLE 2: RESPONDENTS' HEALTH BELIEFS REGARDING COVID-19 AND ITS VACCINES

TABLE 3: CHANGES IN RESPONDENTS' HEALTH BELIEFS REGARDING COVID-19 AND ITS VACCINES

Perceived Susceptibility to COVID-19	% Strongly Agree/Agree
Chance of getting COVID-19 now is lower than it	73
was six months ago	73
Less worried about the likelihood of getting	67.2
COVID-19 now than I was six months ago	07.2
Less worried about the likelihood of someone in	
my family getting COVID-19 now than I was six	63.2
months ago	
Less afraid of getting COVID-19 now than I was	63.8
six months ago	03.8
Less afraid that someone in my family will get	61.2
COVID-19 now than I was six months ago	01.2
Perceived Barriers to Vaccination	
I worry less about the possible side-effects of the	
COVID-19 vaccine now than I was six months	78.8
ago	
I worry less about the effectiveness of the	
COVID-19 vaccine now than I was six months	76.7
ago	
I worry less about the safety of the COVID-19	78.6
vaccine now than I was six months ago	76.0
I worry less about fake COVID-19 vaccines now	64.4
than I was six months ago	04.4
I worry less that the COVID-19 vaccines were not	70.6
tested properly now than I was six months ago	70.0
I worry less that the COVID-19 vaccines were	70.2
made too quickly now than I was six months ago	70.2
I worry less that the COVID-19 vaccines will	72.0
make me sick now than I was six months ago	72.0
I worry less that the COVID-19 vaccines will not	
be effective against new variants of the virus now	62.2
than I was six months ago	
Vaccination Intent	%Definitely Yes/Probably Yes
Are you more willing to take a COVID-19	92.3
vaccine now than you were 6 months ago?	72.3

			Univariab	le analysis		Multivariable logistic regression				
	Overall N (%)	Not Vaccinated n = 856 (16.7%)	Partially Vaccinated & Fully Vaccinated n = 4274 (83.3%)	Unadjusted OR [95% CI]	<i>p</i> -value	Not Vaccinated vs. Partially Vaccinated Adjusted OR [95% CI]	<i>p</i> -value	Not Vaccinated vs. Fully Vaccinated Adjusted OR [95% CI]	<i>p</i> -value	
Demographics										
Age group (years)										
18–30	3453 (67.31)	701 (20.3)	2752 (79.7)	.379 [.288, .498]	< .001	.782 [.406, 1.506]	.463	.251 [.137, .461]	< .001	
31–40	521 (10.16)	41 (7.9)	480 (92.1)	1.129 [.748, 1.704]	.564	1.105 [.562, 2.172]	.772	.500 [.268, .934]	.030	
41–50	451 (8.79)	52 (11.5)	399 (88.5)	.740 [.501, 1.092]	.129	.680 [.351, 1.315]	.680	.448 [.248, .809]	.008	
51–99	705 (13.74)	62 (8.8)	643 (91.2)	Reference		Reference		Reference		
Gender										
Female	3293 (64.19)	616 (18.7)	2677 (81.3)	.653 [.556, .768]	<.001	.774 [.612, .978]	.032	.775 [.618, .972]	.028	
Male	1837 (35.81)	240 (13.1)	1597 (86.9)	Reference		Reference		Reference		
Marital Status										
Single	3883 (75.69)	741 (19.1)	3142 (80.9)	.431 [.350, .530]	< .001	1.216 [.758, 1.950]	.418	.948 [.608, 1.477]	.814	
Married	1247 (24.31)	115 (9.2)	1132 (90.8)	Reference		Reference		Reference		
Highest Education Level										
Elementary school or below	3 (.06)	1 (33.3)	2 (66.7)	.372 [.034, 4.110]	.420	.432 [.036, 5.163]	.508	n too small to calcula	te	
Junior high school	132 (2.57)	14 (10.6)	118 (89.4)	1.568 [.895, 2.747]	.116	1.367 [.559, 3.343]	.494	1.463 [.596, 3.590]	.406	
Senior high school	1036 (20.19)	220 (21.2)	816 (78.8)	.690 [.581, .819]	< .001	1.199 [.929, 1.547]	.164	.914 [.710, 1.176]	.483	
College/university or above	3959 (77.17)	621 (15.7)	3338 (84.3)	Reference		Reference		Reference		
Occupation										

Blue collar worker	878 (17.12)	64 (7.3)	814 (92.7)	1.918 [1.294, 2.843]	.001	1.262 [.669, 2.380]	.472	1.414 [.760, 2.625]	.272
Professional/white collar worker	1627 (31.72)	133 (8.2)	1494 (91.8)	1.694 [1.195, 2.400]	.003	.951 [.543, 1.666]	.860	1.375 [.807, 2.343]	.241
Self-employed	541 (10.55)	62 (11.5)	479 (88.5)	1.165 [.781, 1.738]	.455	1.123 [.593, 2.129]	.721	1.101 [.594, 2.040]	.761
Student	1710 (33.33)	548 (32)	1162 (68)	.320 [.233, .439]	< .001	.247 [.140, .433]	< .001	.297 [.172, .511]	< .001
Housewife/retired/unemployed/other	374 (7.29)	49 (13.1)	325 (86.9)	Reference		Reference		Reference	
Location									
Rural	1043 (20.33)	257 (24.6)	786 (75.4)	.525 [.445, .620]	<.001	.876 [.689, 1.113]	.278	.661 [.523, .863]	< .001
Urban	4087 (79.67)	599 (14.7)	3488 (85.3)	Reference		Reference		Reference	
Experience with COVID-19									
Ever tested positive for COVID-19									
Yes	334 (6.51)	48 (14.4)	286 (85.6)	1.207 [.881, 1.654]	.241				
No	4796 (93.49)	808 (16.8)	3988 (83.2)	Reference					
Know anyone who has tested positive for COVID-19									
Yes	3299 (64.31)	617 (18.7)	2682 (81.3)	.653 [.555, .767]	< .001	.491 [.377, .640]	< .001	.861 [.662, 1.119]	.263
No	1831 (35.69)	239 (13.1)	1592 (86.9)	Reference		Reference			
Health Characteristics									
Have an existing chronic condition									
Yes	635 (12.38)	81 (12.8)	554 (87.3)	1.425 [1.115, 1.822]	.005	.770 [.498, 1.188]	.237	1.503 [1.036, 2.182]	.032
No	4495 (87.62)	775 (17.2)	3720 (82.2)	Reference		Reference		Reference	
Perceived overall health									
Very good	2021 (39.4)	268 (13.3)	1753 (86.7)	Reference		Reference		Reference	
Good	2466	436 (17.7)	2030	.712 [.604,	< .001	.915 [.718, 1.164]	.469	.900 [.711, 1.139]	.379

	(48.07)		(82.3)	.840]					
Fair/Poor/Very poor	643 (12.53)	152 (23.6)	491 (76.4)	.494 [.395, .617]	< .001	.835 [.587, 1.187]	.315	.930 [.669, 1.292]	.664
Openness to the COVID-19 Vaccine									
Would you take a COVID-19 vaccine									
Definitely yes/probably yes	4888 (95.28)	646 (13.2)	4242 (86.8)	Reference		Reference		Reference	
Unsure/probably no/definitely no	242 (4.72)	210 (86.8)	32 (13.2)	.023 [.016, .034]	< .001	.016 [.008, .032]	< .001	.012 [.006, .024]	< .001
More willing to take a COVID-19 vaccine now than you were 6 months ago									
Definitely yes/probably yes	4733 (92.26)	633 (13.4)	4100 (86.6)	Reference		Reference		Reference	
Unsure/probably no/definitely no	397 (7.74)	223 (56.2)	174 (43.8)	.120 [.097, .149]	< .001	.531 [.335, .842]	.007	.488 [.311, .766]	.002
Do you know someone who has been vaccinated against COVID-19									
Yes	4923 (95.96)	834 (16.9)	4089 (83.1)	.583 [.372, .913]	.018	.467 [.236, .922]	.028	.444 [.222, .886]	.021
No	207 (4.04)	22 (10.6)	185 (89.4)	Reference		Reference		Reference	
Health Belief									
Perceived susceptibility									
Chance of getting COVID-19 in the next few months is high									
Strongly agree/agree	2196 (42.81)	386 (17.6)	1810 (82.4)	Reference					
Disagree/strongly disagree	2934 (57.19)	470 (16)	2464 (84)	1.118 [.965, 1.296]	.139				
Worried about the likelihood of getting COVID-19									
Strongly agree/agree	4158 (81.05)	693 (16.7)	3465 (83.3)	Reference					
Disagree/strongly disagree	972 (18.95)	163 (16.8)	809 (83.2)	.993 [.823, 1.197]	.938				
Worried about the likelihood of someone in my family getting COVID-									

19									
Strongly agree/agree	4443 (86.61)	749 (16.9)	3694 (83.1)	Reference					
Disagree/strongly disagree	687 (13.39)	107 (15.6)	580 (84.4)	1.099 [.881, 1.370]	.401				
Perceived severity									
COVID-19 is a serious disease with life-threatening complications									
Strongly agree/agree	4920 (95.91)	800 (16.3)	4120 (83.7)	Reference		Reference		Reference	
Disagree/strongly disagree	210 (4.09)	56 (26.7)	154 (73.3)	.534 [.390, .732]	< .001	1.122 [.609, 2.064]	.713	.924 [.500, 1.711]	.802
I will be very sick if I get COVID-19									
Strongly agree/agree	3700 (72.12)	667 (18)	3033 (82)	Reference		Reference		Reference	
Disagree/strongly disagree	1430 (27.88)	189 (13.2)	1241 (86.8)	1.444 [1.213, 1.719]	< .001	1.635 [1.221, 2.191]	< .001	2.052 [1.549, 2.717]	< .001
I am afraid of getting COVID-19									
Strongly agree/agree	4757 92.73)	759 (16)	3998 (84)	Reference		Reference		Reference	
Disagree/strongly disagree	373 (7.27)	97 (26)	276 (745)	.540 [.423, .689]	< .001	.642 [.387, 1.063]	.085	.688 [.425, 1.115]	.129
I am afraid that someone in my family will get COVID-19									
Strongly agree/agree	4919 (95.59)	801 (16.3)	4118 (83.7)	Reference		Reference		Reference	
Disagree/strongly disagree	211 (4.11)	55 (26.1)	156 (73.9)	.552 [.402, .757]	< .001	1.033 [.534, 1.999]	.923	.714 [.372, 1.369]	.310
Perceived benefits									
Chance of getting COVID-19 now is lower than it was six months ago									
Strongly agree/agree	3745 (73)	356 (9.5)	3389 (90.5)	Reference		Reference		Reference	
Disagree/strongly disagree	1385 (27)	500 (36.1)	885 (63.9)	.186 [.159, .217]	< .001	.275 [.208, .364]	< .001	.224 [.171, .293]	<.001
Less worried about the likelihood of getting COVID-19 now than I was six months ago									

Strongly agree/agree	3447	331 (9.6)	3116	Reference		Reference		Reference	
Strongly agree/agree	(67.19)	331 (9.0)	(90.4)	Reference		Reference		Reference	
Disagree/strongly disagree	1683 (32.81)	525 (31.2)	1158 (68.8)	.234 [.201, .273]	< .001	.637 [.424, .957]	.030	.515 [.349, .762]	< .001
Less worried about the likelihood of someone in my family getting COVID-19 now than I was six months ago									
Strongly agree/agree	3241 (63.18)	330 (10.2)	2911 (89.8)	Reference		Reference		Reference	
Disagree/strongly disagree	1889 (36.82)	526 (27.8)	1363 (72.2)	.294 [.252, .342]	< .001	1.419 [.908, 2.217]	.125	1.750 [1.139, 2.687]	.011
Less afraid of getting COVID-19 now than I was six months ago									
Strongly agree/agree	3273 (63.8)	309 (9.4)	2964 (90.6)	Reference		Reference		Reference	
Disagree/strongly disagree	1857 (36.2)	547 (29.5)	1310 (70.5)	.250 [.214, .291]	<.001	.687 [.431, 1.095]	.114	.578 [.371, .902]	.016
Less afraid that someone in my family will get COVID-19 now than I was six months ago									
Strongly agree/agree	3141 (61.23)	307 (9.8)	2834 (90.2)	Reference		Reference		Reference	
Disagree/strongly disagree	1989 (38.77)	549 (27.6)	1440 (72.4)	.284 [.244, .331]	<.001	.967 [.592, 1.580]	.893	1.094 [.683, 1.751]	.709
Vaccination Belief Model									
Vaccination will decrease my chances of getting COVID-19									
Strongly agree/agree	4481 (87.35)	662 (14.8)	3819 (85.2)	Reference		Reference		Reference	
Disagree/strongly disagree	649 (12.65)	194 (29.9)	455 (70.1)	.407 [.337, .490]	< .001	1.145 [.787, 1.667]	.479	1.114 [.781, 1.590]	.551
Vaccination will decrease my worries about catching COVID-19									
Strongly agree/agree	4503 (87.78)	661 (14.7)	3842 (85.3)	Reference		Reference		Reference	
Disagree/strongly disagree	627 (12.22)	195 (31.1)	432 (68.9)	.381 [.316, .460]	< .001	2.066 [1.391, 3.068]	<.001	1.819 [1.246, 2.654]	.002
Perceived barriers									

I worry about the possible side-effects of the COVID-19 vaccine									
Strongly agree/agree	3495 (68.13)	652 (18.7)	2843 (81.3)	Reference		Reference		Reference	
Disagree/strongly disagree	1635 (31.87)	204 (12.5)	1431 (87.5)	1.609 [1.358, 1.906]	< .001	1.002 [.751, 1.338]	.988	1.666 [1.271, 2.184]	< .001
I worry about the effectiveness of the COVID-19 vaccine									
Strongly agree/agree	3505 (68.32)	632 (18)	2873 (82)	Reference		Reference		Reference	
Disagree/strongly disagree	1625 (31.68)	224 (13.8)	1401 (86.2)	1.376 [1.166, 1.623]	< .001	1.054 [.770, 1.443]	.744	.929 [.690, 1.250]	.625
I worry about the safety of the COVID-19 vaccine									
Strongly agree/agree	3268 (63.7)	598 (18.3)	2670 (81.7)	Reference		Reference		Reference	
Disagree/strongly disagree	1862 (36.3)	258 (13.9)	1604 (86.1)	1.392 [1.188, 1.632]	< .001	.838 [.604, 1.161]	.287	.910 [.667, 1.242]	.552
I worry about the high cost of the COVID-19 vaccine									
Strongly agree/agree	3239 (63.14)	534 (16.5)	2705 (83.5)	Reference					
Disagree/strongly disagree	1891 (36.86)	322 (17)	1569 (83)	.962 [.827, 1.119]	.616				
I worry about fake COVID-19 vaccines									
Strongly agree/agree	4541 (88.52)	773 (17)	3768 (83)	Reference					
Disagree/strongly disagree	589 (11.48)	83 (14.1)	506 (85.9)	1.251 [.979, 1.597	.073				
I worry that the COVID-19 vaccines were made too quickly									
Strongly agree/agree	3056 (59.57)	533 (17.4)	1751 (82.6)	Reference					
Disagree/strongly disagree	2074 (40.43)	323 (15.6)	2523 (84.4)	1.145 [.985, 1.332]	.078				
I worry that the COVID-19 vaccines were not tested properly									
Strongly agree/agree	2828	522 (18.5)	2306	Reference		Reference		Reference	

	(55.13)		(81.5)						
Disagree/strongly disagree	2302 (44.87)	334 (14.5)	1968 (85.5)	1.334 [1.148, 1.550]	<.001	.920 [.699, 1.210]	.550	1.034 [.795, 1.345]	.803
I worry that the COVID-19 vaccines will make me sick									
Strongly agree/agree	2584 (50.37)	542 (21)	2042 (79)	Reference		Reference		Reference	
Disagree/strongly disagree	2546 (49.63)	314 (12.3)	2232 (87.7)	1.887 [1.622, 2.195]	< .001	1.807 [1.348, 2.421]	<.001	2.683 [2.028, 3.551]	< .001
I worry that the COVID-19 vaccines will not be effective against new variants of the virus									
Strongly agree/agree	3872 (75.48)	672 (17.4)	3200 (82.6)	Reference		Reference		Reference	
Disagree/strongly disagree	1258 (24.52)	184 (14.6)	1074 (85.4)	1.226 [1.027, 1.463]	.024	.806 [.607, 1.069]	.134	.574 [.438, .753]	< .001
I worry less about the possible side- effects of the COVID-19 vaccine now than I was six months ago									
Strongly agree/agree	4040 (78.75)	496 (12.3)	3544 (87.7)	Reference		Reference		Reference	
Disagree/strongly disagree	1090 (21.25)	360 (33)	730 (67)	.284 [.242, .332]	< .001	.895 [.642, 1.249]	.514	.781 [.570, 1.070]	.124
I worry less about the effectiveness of the COVID-19 vaccine now than I was six months ago									
Strongly agree/agree	3932 (76.65)	488 (12.4)	3444 (87.6)	Reference		Reference		Reference	
Disagree/strongly disagree	1198 (23.35)	368 (30.7)	830 (69.3)	.320 [.274, .373]	< .001	1.198 [.809, 1.775]	.368	1.167 [.807, 1.689]	.412
I worry less about the safety of the COVID-19 vaccine now than I was six months ago									
Strongly agree/agree	4034 (78.64)	488 (12.1)	3546 (87.9)	Reference		Reference		Reference	
Disagree/strongly disagree	1096 (21.36)	368 (33.6)	728 (66.4)	.272 [.233, .319]	< .001	.816 [.538, 1.238]	.339	.853 [.576, 1.264]	.429
I worry less about fake COVID-19 vaccines now than I was six months									

ago									
Strongly agree/agree	3303 (64.39)	360 (10.9)	2943 (89.1)	Reference		Reference		Reference	
Disagree/strongly disagree	1827 (35.61)	496 (27.1)	1331 (72.9)	.328 [.282, .382]	< .001	.749 [.556, 1.008]	.056	.692 [.523, .916]	.010
I worry less that the COVID-19 vaccines were not tested properly now than I was six months ago									
Strongly agree/agree	3620 (70.57)	431 (11.9)	3189 (88.1)	Reference		Reference		Reference	
Disagree/strongly disagree	1510 (29.43)	425 (28.1)	1085 (71.9)	.345 [.297, .401]	< .001	1.013 [.659, 1.558]	.952	.843 [.557, 1.276]	.420
I worry less that the COVID-19 vaccines were made too quickly now than I was six months ago									
Strongly agree/agree	3603 (70.23)	428 (11.9)	3175 (88.1)	Reference		Reference		Reference	
Disagree/strongly disagree	1527 (29.77)	428 (28)	1099 (72)	.346 [.298, .402]	< .001	.901 [.597, 1.362]	.621	1.035 [.696, 1.539]	.865
I worry less that the COVID-19 vaccines will make me sick now than I was six months ago									
Strongly agree/agree	3692 (71.97)	437 (11.8)	3255 (88.2)	Reference		Reference		Reference	
Disagree/strongly disagree	1438 (28.03)	419 (29.1)	1019 (70.9)	.327 [.281, .380]	< .001	.840 [.569, 1.241]	.381	.988 [.683, 1.429]	.948
I worry less that the COVID-19 vaccines will not be effective against new variants of the virus now than I was six months ago									
Strongly agree/agree	3192 (62.22)	379 (11.9)	2813 (88.1)	Reference		Reference		Reference	
Disagree/strongly disagree	1938 (37.78)	477 (24.6)	1461 (75.4)	.413 [.356, .479]	<.001	1.232 [.906, 1.676]	.183	1.442 [1.079, 1.928]	.013
Cues to action									
Rewarding fully vaccinated individuals with bakuna benefits would encourage me to get vaccinated against COVID-									

Strongly agree/agree	4102 (79.96)	573 (14)	3529 (86)	Reference		Reference		Reference	
Disagree/strongly disagree	1028 (20.04)	283 (27.5)	745 (72.5)	.427 [.363, .503]	<.001	.909 [.664, 1.245]	.554	.808 [.600, 1.090]	.163
Rewarding fully vaccinated individuals with more freedom to move around and to meet with people would encourage me to get vaccinated against COVID-19									
Strongly agree/agree	4040 (78.75)	550 (13.6)	3490 (86.4)	Reference		Reference		Reference	
Disagree/strongly disagree	1090 (21.25)	306 (28.1)	784 (71.9)	.404 [.344, .474]	<.001	.926 [.668, 1.282]	.642	.864 [.636, 1.175]	.351
Rewarding fully vaccinated individuals by allowing them to remove masks in the company of other fully vaccinated individuals would encourage me to get vaccinated against COVID-19									
Strongly agree/agree	3011 (58.69)	415 (13.8)	2596 (86.2)	Reference		Reference		Reference	
Disagree/strongly disagree	2119 (41.31)	441 (20.8)	1678 (79.2)	.608 [.525, .705]	<.001	.897 [.680, 1.184]	.442	.932 [.716, 1.213]	.599
Testimonies from fully vaccinated people have encouraged me to get vaccinated									
Strongly agree/agree	4489 (87.5)	647 (14.4)	3842 (85.6)	Reference		Reference		Reference	
Disagree/strongly disagree	641 (12.5)	209 (32.6)	432 (67.4)	.348 [.289, .419]	< .001	.892 [.577, 1.380]	.608	.809 [.534, 1.226]	.317
Seeing my family members and friends get vaccinated has encouraged me to get the vaccine									
Strongly agree/agree	4624 (90.14)	691 (14.9)	3933 (85.1)	Reference		Reference		Reference	
Disagree/strongly disagree	506 (9.86)	165 (32.6)	341 (67.4)	.363 [.296, .445]	< .001	1.355 [.815, 2.253]	.242	1.211 [.739, 1.983]	.447
Seeing famous people get vaccinated on social media has encouraged me to get the vaccine				-					
Strongly agree/agree	3223 (62.83)	443 (13.7)	2780 (86.3)	Reference		Reference		Reference	

Disagree/strongly disagree	1907 (37.17)	413 (21.7)	1494 (78.3)	.576 [.497, .668]	< .001	1.320 [1.021, 1.707]	.034	1.279 [.999, 1.636] .051
I think the national government has managed this COVID-19 vaccination campaign								
Strongly agree/agree	2727 (53.16)	300 (11)	2427 (89)	Reference		Reference		Reference
Disagree/strongly disagree	2403 (46.84)	556 (23.1)	1847 (76.9)	.411 [.352, .478]	< .001	.957 [.723, 1.266]	.757	1.012 [.775, 1.321] .932
I think my LGU has effectively managed this COVID-19 vaccination campaign								
Strongly agree/agree	3503 (68.28)	390 (11.1)	3113 (88.9)	Reference		Reference		Reference
Disagree/strongly disagree	1627 (31.72)	466 (28.6)	1161 (71.4)	.312 [.269, .363]	< .001	.645 [.497, .839]	.001	.464 [.362, .593] < .001

			ssociated with C tus after nation rollout		Factors Associated with intention to take the COVID-19 vaccine before national vaccine rollout (Caple et al., 2022)					
	Overall N (%)	Not Vaccinated n = 856 (16.7%)	Partially Vaccinated & Fully Vaccinated n = 4274 (83.3%)	Individual Predictor of vaccine status?	Overall N (%)	Vaccine hesitant n = 2696 (%)	Not vaccine hesitant $n = 4497$ (%)	Individual Predictor of vaccine intent?		
Demographics										
Age group (years)				Yes				No		
18–30	3453 (67.31)	701 (20.3)	2752 (79.7)		3770 (52.4)	1405 (37.3)	2365 (62.7)			
31–40	521 (10.16)	41 (7.9)	480 (92.1)		815 (11.3)	306 (37.5)	509 (62.5)			
41–50	451 (8.79)	52 (11.5)	399 (88.5)		861 (12)	335 (38.9)	526 (61.1)			
51–99	705 (13.74)	62 (8.8)	643 (91.2)		1747 (24.29)	650 (37.21)	1097 (62.79)			
Gender				Yes			, , ,	Yes		
Female	3293 (64.19)	616 (18.7)	2677 (81.3)		4789 (66.6)	1955 (40.8)	2834 (59.2)			
Male	1837 (35.81)	240 (13.1)	1597 (86.9)		2404 (33.4)	741 (30.8)	1663 (69.2)			
Marital Status				Yes				Yes		
Single	3883 (75.69)	741 (19.1)	3142 (80.9)		4724 (65.7)	1719 (36.4)	3005 (63.6)			
Married	1247 (24.31)	115 (9.2)	1132 (90.8)		2469 (34.3)	977 (39.6)	1492 (60.4)			
Highest Education Level				Yes				Yes		
Elementary school or below	3 (.06)	1 (33.3)	2 (66.7)		1	1				
Junior high school	132 (2.57)	14 (10.6)	118 (89.4)		46 (.6)	29 (63)	17 (37)			
Senior high school	1036 (20.19)	220 (21.2)	816 (78.8)		1102 (15.3)	378 (34.3)	724 (65.7)			
College/university or above	3959 (77.17)	621 (15.7)	3338 (84.3)		6044 (84)	2288 (37.9)	3756 (62.1)			
Occupation				Yes				Yes		
Blue collar worker	878 (17.12)	64 (7.3)	814 (92.7)		91 (1.3)	42 (46.2)	49 (53.8)			
Professional/white collar worker	1627 (31.72)	133 (8.2)	1494 (91.8)		2980 (41.4)	1133 (38)	1847 (62)			
Self-employed	541 (10.55)	62 (11.5)	479 (88.5)		575 (8)	254 (44.2)	321 (55.8)			
Student	1710 (33.33)	548 (32)	1162 (68)		2831 (39.4)	976 (34.5)	1855 (65.5)			
Housewife/retired/unemployed/other	374 (7.29)	49 (13.1)	325 (86.9)		716 (10)	291 (40.6)	425 (59.4)			
Location			_	Yes			_	Yes		
Rural	1043 (20.33)	257 (24.6)	786 (75.4)		1517 (21.1)	711 (46.9)	806 (53.1)			
Urban	4087	599 (14.7)	3488 (85.3)		5676	1985 (35)	3691 (65)			

	(70.67)				(79.0)			
Experience with COVID-19	(79.67)				(78.9)			
Ever tested positive for COVID-19				No				No
Yes	334 (6.51)	48 (14.4)	286 (85.6)		211 (2.9)	68 (32.2)	143 (67.8)	
No	4796 (93.49)	808 (16.8)	3988 (83.2)		6982 (97.1)	2628 (37.6)	4354 (62.4)	
Know anyone who has tested positive for COVID-19				Yes				Yes
Yes	3299 (64.31)	617 (18.7)	2682 (81.3)		5234 (72.8)	1780 (34)	3454 (66)	
No	1831 (35.69)	239 (13.1)	1592 (86.9)		1959 (27.2)	916 (46.8)	1043 (53.2)	
Health Characteristics								
Have an existing chronic condition				Yes				Yes
Yes	635 (12.38)	81 (12.8)	554 (87.3)		1178 (16.4)	396 (33.6)	782 (66.4)	
No	4495 (87.62)	775 (17.2)	3720 (82.2)		6015 (83.6)	2300 (38.2)	3715 (61.8)	
Perceived overall health				Yes				Yes
Very good	2021 (39.4)	268 (13.3)	1753 (86.7)		1997 (27.8)	802 (40.2)	1195 (59.8)	
Good	2466 (48.07)	436 (17.7)	2030 (82.3)		3934 (54.7)	1376 (35)	2558 (65)	
Fair/Poor/Very poor	643 (12.53)	152 (23.6)	491 (76.4)		1262 (17.5)	518 (41)	744 (59)	
Health Belief								
Perceived susceptibility								
Chance of getting COVID-19 in the next few months is high				No				Yes
Strongly agree/agree	2196 (42.81)	386 (17.6)	1810 (82.4)		2264 (31.5)	1595 (70.5)	669 (29.5)	
Disagree/strongly disagree	2934 (57.19)	470 (16)	2464 (84)		4929 (68.5)	2027 (41.1)	2902 (58.9)	
Worried about the likelihood of getting COVID-19				No				Yes
Strongly agree/agree	4158 (81.05)	693 (16.7)	3465 (83.3)		6047 (84.1)	2105 (34.8)	3942 (65.2)	
Disagree/strongly disagree	972 (18.95)	163 (16.8)	809 (83.2)		1146 (15.9)	591 (51.6)	555 (48.4)	
Perceived severity								
COVID-19 is a serious disease with life-threatening complications				Yes				Yes
Strongly agree/agree	4920 (95.91)	800 (16.3)	4120 (83.7)		6928 (96.32)	2523 (36.4)	4405 (63.6)	
Disagree/strongly disagree	210 (4.09)	56 (26.7)	154 (73.3)		265 (3.68)	173 (65.3)	95 (34.7)	
I will be very sick if I get COVID-19				Yes				Yes
Strongly agree/agree	3700 (72.12)	667 (18)	3033 (82)		5394 (75)	1911 (35.4)	3483 (64.6)	
Disagree/strongly disagree	1430	189 (13.2)	1241 (86.8)		1799	785 (43.6)	1014	

T. (11.6 w) COVED 10	(27.88)			**	(25)		(56.4)	
I am afraid of getting COVID-19				Yes				Yes
Strongly agree/agree	4757 92.73)	759 (16)	3998 (84)		6700 (93.15)	2439 (36.4)	4261 (63.6)	
Disagree/strongly disagree	373 (7.27)	97 (26)	276 (745)		493 (6.85)	257 (52.1)	236 (47.9)	
Vaccination Belief Model								
Vaccination will decrease my chances of getting COVID-19				Yes				Yes
Strongly agree/agree	4481 (87.35)	662 (14.8)	3819 (85.2)		6339 (88.1)	2006 (31.6)	4333 (68.4)	
Disagree/strongly disagree	649 (12.65)	194 (29.9)	455 (70.1)		854 (11.9)	690 (80.8)	164 (19.2)	
Vaccination will decrease my worries about catching COVID-19				Yes				Yes
Strongly agree/agree	4503 (87.78)	661 (14.7)	3842 (85.3)		6078 (84.5)	1833 (30.2)	4245 (69.8)	
Disagree/strongly disagree	627 (12.22)	195 (31.1)	432 (68.9)		1115 (15.5)	863 (77.4)	252 (22.6)	
Perceived barriers								
I worry about the possible side-effects of the COVID-19 vaccine				Yes				Yes
Strongly agree/agree	3495 (68.13)	652 (18.7)	2843 (81.3)		6447 (89.6)	2647 (41.1)	3800 (58.9)	
Disagree/strongly disagree	1635 (31.87)	204 (12.5)	1431 (87.5)		746 (10.4)	49 (6.6)	697 (93.4)	
I worry about the effectiveness of the COVID-19 vaccine				Yes				Yes
Strongly agree/agree	3505 (68.32)	632 (18)	2873 (82)		6263 (87.1)	2581 (41.2)	3682 (58.8)	
Disagree/strongly disagree	1625 (31.68)	224 (13.8)	1401 (86.2)		930 (12.9)	115 (12.4)	815 (87.6)	
I worry about the safety of the COVID- 19 vaccine				Yes				Yes
Strongly agree/agree	3268 (63.7)	598 (18.3)	2670 (81.7)		6389 (88.8)	2626 (41.1)	3763 (58.9)	
Disagree/strongly disagree	1862 (36.3)	258 (13.9)	1604 (86.1)		804 (11.2)	70 (8.7)	734 (91.3)	
I worry about the high cost of the COVID-19 vaccine				No				Yes
Strongly agree/agree	3239 (63.14)	534 (16.5)	2705 (83.5)		5608 (78)	2301 (41)	3307 (59)	
Disagree/strongly disagree	1891 (36.86)	322 (17)	1569 (83)		1585 (22)	395 (24.9)	1190 (75.1)	
I worry about fake COVID-19 vaccines				No				Yes
Strongly agree/agree	4541 (88.52)	773 (17)	3768 (83)		7006 (97.4)	2646 (37.8)	4360 (62.2)	
Disagree/strongly disagree	589 (11.48)	83 (14.1)	506 (85.9)		187 (.03)	50 (26.7)	197 (73.3)	
I worry that the COVID-19 vaccines will make me sick				Yes				Yes
Strongly agree/agree	2584	542 (21)	2042 (79)		4918	2404 (48.9)	2514	

	(50.37)				(68.4)		(51.1)	
Disagree/strongly disagree	2546 (49.63)	314 (12.3)	2232 (87.7)		2275 (31.6)	292 (12.8)	1983 (87.2)	
I worry that the COVID-19 vaccines will not be effective against new variants of the virus				Yes				Yes
Strongly agree/agree	3872 (75.48)	672 (17.4)	3200 (82.6)		5757 (80)	2506 (43.5)	3251 (56.5)	
Disagree/strongly disagree	1258 (24.52)	184 (14.6)	1074 (85.4)		1436 (20)	190 (13.2)	1246 (86.8)	