

1 **Title** : A vagal influence on schizophrenia? A nationwide retrospective cohort of vagotomized
2 individuals

3

4 **Authors**: Cornelia F Richter¹, Karolina P Skibicka^{2,3}, Urs Meyer^{1,4}, Sabine Rohrmann⁵, Jean-
5 Philippe Krieger^{1,2}

6

7 **Affiliations**:

8 1 Institute of Veterinary Pharmacology and Toxicology, University of Zurich-Vetsuisse,
9 Switzerland

10 2 Institute of Neuroscience and Physiology, University of Gothenburg, Sweden

11 3 Department of Nutritional Sciences, Pennsylvania State University, State College, PA,
12 United States

13 4 Neuroscience Center Zurich, University of Zurich and ETH Zurich, Switzerland

14 5 Division of Chronic Disease Epidemiology, Epidemiology, Biostatistics and Prevention
15 Institute, University of Zurich, Switzerland

16

17

18 **Corresponding author**: Jean-Philippe Krieger. Institute of Neuroscience and Physiology,
19 University of Gothenburg, Sweden (jean-philippe.krieger@gu.se).

20

21 **Keywords**: vagus nerve, schizophrenia, gut-brain axis, ulcer, cohort study,

22

23 **Abstract:**

24 Emerging preclinical evidence suggests that vagal signals contribute to the development of
25 schizophrenia-related abnormalities in brain and behavior. Whether vagal communication in
26 general, and its impairment in particular, is a risk factor for schizophrenia in humans remains,
27 however, unclear.

28 Vagotomy, the surgical lesion of the vagus nerve, was routinely performed as a treatment for
29 peptic ulcer before modern treatment options were available. Hence, the primary aim of this
30 study was to investigate whether vagotomy modulates the subsequent risk of developing
31 schizophrenia. Moreover, given the existence of diverse vagotomy techniques (i.e., “truncal”
32 or “selective”), our secondary goal was to test whether the extent of denervation modulates
33 the risk of schizophrenia.

34 Using a nationwide retrospective matched cohort design, we identified 8,833 vagotomized
35 individuals from the Swedish National Patient Register during the period 1970-2020 and
36 17,666 non-vagotomized individuals matching for age, sex and type of peptic ulcer. The risk
37 of being diagnosed with schizophrenia and associated psychoses (ICD10 codes F20-29) was
38 analyzed using Cox proportional hazards regression models, including death as competing
39 risk.

40 Overall, vagotomy was not significantly associated with schizophrenia (HR: 0.86 [0.66; 1.12]).
41 When considering different vagotomy techniques, however, truncal vagotomy significantly
42 increased (HR: 1.60 [1.02; 2.51]), while selective vagotomy decreased (HR: 0.75 [0.56; 1.00]),
43 the risk of developing schizophrenia.

44 Our results provide epidemiological support for the hypothesis that impairments in vagal
45 functions can increase the risk of schizophrenia. Notably, the finding that truncal but not
46 selective vagotomy is associated with an increased risk of schizophrenia raises the possibility
47 that the activity of subdiaphragmatic non-gastric vagal branches may be of particular
48 relevance for the development of schizophrenia.

49 **1 Introduction**

50 The vagus nerve is a major neuronal route of communication between the periphery and the
51 brain (Neuhuber and Berthoud, 2021). Vagal afferent neurons, the ascending component of
52 this communication, sense peripheral signals and, thus, inform the brain about the state of the
53 periphery. On the other hand, vagal efferent neurons provide top-down parasympathetic
54 control over organ physiology.

55 Several lines of evidence suggest that this bidirectional communication is relevant for the
56 development of schizophrenia. First, vagal parasympathetic activity is decreased in patients
57 with schizophrenia, which contributes to overall autonomic nervous system dysfunction in
58 affected individuals (Stogios et al., 2021). This dysfunction underlies cardiovascular
59 complications associated with schizophrenia, but is also increasingly postulated to contribute
60 to psychological symptoms severity (Stogios et al., 2021). In addition, vagal efferent neurons
61 exert an anti-inflammatory action through the cholinergic anti-inflammatory pathway (Bonaz et
62 al., 2016). Hence, impaired vagal function has been proposed to contribute to systemic as well
63 as central inflammation seen in a subset of patients with schizophrenia (Corsi-Zuelli et al.,
64 2017).

65 In comparison to the descending vagal circuits, little focus has been given to ascending vagal
66 afferents in schizophrenia. Several lines of evidence suggest, however, that vagal afferents
67 may also play a significant role in pathophysiology. First, the activity of vagal afferent neurons
68 directly influences the descending pathways and, hence, can contribute to the effects
69 mentioned above. In addition, while recent studies emphasize the role of the gut microbiome
70 in schizophrenia (Thirion et al., 2023; Zhu et al., 2020), vagal sensory neurons are well-
71 positioned to be the main mediators of the relevant microbiome metabolites (Dinan et al.,
72 2014). Finally, pre-clinical rodent studies indicate that vagal afferent signals could contribute
73 to the development of schizophrenia-related abnormalities in brain and behavior. Indeed,
74 subdiaphragmatic vagal deafferentation in rats leads to brain transcriptional changes in
75 functional networks related to schizophrenia, increased sensitivity to dopamine-stimulating
76 drugs, and impairments in sensorimotor gating and the attentional control of associative
77 learning (Klarer et al., 2018).

78 Despite these findings, direct evidence for a vagal influence on the pathophysiology of
79 schizophrenia is lacking in humans. However, such evidence could foster further therapies
80 targeted at the vagus nerve in schizophrenia and open avenues for more mechanistic studies.

81 Vagotomy, the lesion of the vagus nerve, was performed to treat peptic ulcers before the
82 advent of more effective treatments, with the goal to reduce gastric acid secretion. Several
83 vagotomy techniques co-existed and can be categorized into “truncal vagotomy”, the section

84 of the vagal trunks below the diaphragm, and “selective vagotomy”, which aimed to selectively
85 severe vagal sub-branches to the stomach (Lundell, 2011).

86 Several epidemiological studies have taken advantage of cohorts of vagotomized individuals
87 to assess vagal influence on dementia (Lin et al., 2018), Parkinson’s disease (Liu et al., 2017),
88 inflammatory bowel disease (Liu et al., 2020) and mental disorders (Bunyozy et al., 2022).
89 While this last study included schizophrenia as a secondary outcome, a large study assessing
90 the risk of developing schizophrenia in vagotomized individuals as a primary outcome is
91 lacking. Therefore, we built a nationwide retrospective cohort of patients who underwent
92 vagotomy in Sweden and contrasted their risk of developing schizophrenia with that of control
93 individuals matched for age, sex and peptic ulcer diagnosis over a 50-year period. In addition,
94 we took advantage of the diverse vagotomy techniques to assess whether the extent of
95 denervation modulates the risk of developing schizophrenia.

96 **2 Methods**

97 **2.1 Ethics and reporting**

98 This study was approved by the Swedish Ethical review authority (decision 2019-04833 and
99 amendment 2021-05371-02). The current report is structured according to the STROBE
100 guidelines (Strengthening the Reporting of Observational Studies in Epidemiology; (von Elm
101 et al., 2007)).

102 **2.2 Study design**

103 This study is a register-based nationwide retrospective matched cohort covering the period
104 1970-2020 in Sweden.

105 **2.3 Data sources**

106 Anonymized data from different registers were provided by the Swedish National Board of
107 Health and Welfare and Statistics Sweden. Specifically, the National Patient Register (NPR),
108 the Swedish Cause of Death Register and the Total Population Register were used and linked
109 by the Swedish personal identification number, which exists for each resident in Sweden
110 (Socialstyrelsen, 2024). Since the early 1960s information from in-patients at public hospitals
111 is documented in the NPR, including reports of psychiatric care. Nationwide coverage exists
112 since 1987 (Ludvigsson et al., 2011). Furthermore, since 2001 outpatient information (doctor
113 visits, psychiatric care and surgery) from private and public institutions are covered.
114 Involuntary psychiatric care and forensic psychiatric care are also reported since 2008.
115 Primary health care doctor visits are not yet documented in the NPR (Ludvigsson et al., 2011).
116 The death register contains information on deaths from 1964. Regular, thorough quality and
117 validity checks are done on data submitted to the National Board of Health and Welfare.
118 Information about dates of immigration into (from 1969) and emigration out of (from 1961)
119 Sweden is recorded in the Total Population Register.

120 **2.4 Study population**

121 A total of 14,206 individuals who underwent vagotomy between 01-01-1970 and 31-12-2020
122 and did not have a diagnosis of schizophrenia prior to surgery were identified from the National
123 Patient Register based on the Swedish Classification of Operations and Major Procedures
124 (Table S1) as described in (Liu et al., 2020). The closest diagnosis of a peptic ulcer (ICD-10
125 codes K25-K28, see Table S2) prior to vagotomy was considered as indication for vagotomy.
126 Individuals identified from the NPR as vagotomized were excluded if surgery dates were not
127 plausible (ie, before admission at the hospital or after death dates), if vagotomy surgery code
128 was missing from the source file or if an ulcer diagnosis did not exist prior to or within 14 days
129 post surgery. Two controls per case were then selected from the NPR using a stepwise

130 procedures: first, all controls matching a vagotomy case for age at cohort entry, sex and
131 presenting the same peptic ulcer diagnosis (exact ICD codes) were selected. As a second
132 step, the two controls that received the peptic ulcer diagnosis at the closest date to the
133 vagotomy case were selected. The same control individuals could not be used as a control for
134 multiple cases. In case a vagotomized individual could not be matched with two controls, the
135 corresponding case-control group was excluded. Overall, 8,833 vagotomy patients and 17,666
136 controls, were included in the study, as described in the flow chart (Figure 1).

137 **2.5 Ascertainment of surgery types**

138 Vagotomy surgeries were categorized into “truncal” and “selective” as previously described in
139 (Liu et al., 2020) based on the Swedish classification of Operations and Major procedures
140 (Table S1). Briefly, truncal vagotomy lesions both vagal trunks around the oesophagus and
141 therefore denervate all subdiaphragmatic organs. Selective vagotomies cover multiple
142 techniques achieving a selective denervation of the stomach (Seeras et al., 2023). Five
143 individuals had surgery codes that cannot be assigned to truncal or selective and were
144 therefore only considered for the analysis of the overall vagotomy effects.

145 **2.6 Ascertainment of schizophrenia diagnosis**

146 Due to historical evolution in the diagnosis of schizophrenia (Jablensky, 2010) and the
147 heterogeneity of the disease (Tsuang et al., 2000), we opted for a broad definition of
148 schizophrenia for this long-term nationwide cohort. Hence, diagnoses of schizophrenia,
149 schizotypal disorders and non-mood psychotic disorders were identified by the presence of
150 the corresponding ICD codes in the NPR or the Death register (ICD10 codes F20-F29; see
151 Table S3 for earlier ICD versions). We refer to these disorders as “schizophrenia” in the rest
152 of the manuscript. In case of several diagnoses, the earliest diagnosis and diagnosis dates
153 were considered.

154 **2.7 Covariates**

155 Age and sex of participants were taken from the NPR. Overall comorbidity at cohort entry was
156 assessed with an adapted version (Deyo et al., 1992) of the weighted Charlson Comorbidity
157 Index (CCI) (Charlson et al., 1987; Ludvigsson et al., 2021). The CCI is a cumulative score
158 including diagnoses of vascular diseases, diabetes, chronic obstructive pulmonary disease,
159 mild, moderate and severe liver disease, cancer and chronic obesity. Relevant ICD codes can
160 be found in supplementary material (Table S4).

161 **2.8 Handling of missing data**

162 For 706 patients in the death registry, exact day of death was not indicated and was imputed
163 as the 15th of the corresponding month. Further, there were 125 cases for which month and

164 day of death were missing. In this case, we used December 15th of the respective year of
165 death.

166 **2.9 Statistical methods**

167 Cohort participants were followed up from the day of surgery, which was matched for the
168 reference population, to the date of a schizophrenia diagnosis, death, emigration of Sweden
169 or December 31st, 2020, whichever occurred first.

170 The first analysis assessed the association between overall vagotomy and the postoperative
171 risk of developing schizophrenia. In a secondary analysis, the association between vagotomy
172 types (truncal, selective) and schizophrenia was investigated. In a third analysis, we restricted
173 follow up to 5, 10, 15 and 20 years after cohort entry and analyzed the association between
174 overall vagotomy, as well as vagotomy types (truncal, selective) and schizophrenia.

175 For all analysis, Cox proportional hazard regression models and hazard ratios with
176 corresponding 95% confidence intervals were computed for the outcome schizophrenia,
177 including death as a competing risks. We considered a crude model, accounting for matching
178 variables age and sex at cohort entry and an adjusted model, additionally including the ICD
179 period and the adapted Charlson Comorbidity Index. All analyses were conducted with R
180 (version 4.3.1).

181 **3 Results**

182 Table 1 summarizes characteristics of the study participants. As a result of matching, the
183 distributions of sex (66% males), age (median age 54) and peptic ulcer types were the same
184 in vagotomized patients and controls. Most vagotomy patients (n = 6,524, 73.8 %), underwent
185 selective vagotomy. Individuals with truncal vagotomy had higher comorbidity (CCI) at
186 baseline and higher mortality during follow-up than reference individuals or individuals with
187 selective vagotomy.

188 The cumulative incidence of schizophrenia in vagotomized patients compared to the matched
189 reference individuals is shown in Figure 2 (overall vagotomy) and Figure 3 (truncal and
190 selective vagotomy). Overall, we found no statistically significant association of vagotomy with
191 schizophrenia (HR crude model: 0.88 [0.68; 1.13]; adjusted 0.86 [0.66; 1.12], Figure 4). When
192 considering vagotomy types, however, selective vagotomy decreased (HR: 0.75 [0.56; 1.00],
193 Figure 5), while truncal vagotomy increased (HR: 1.60 [1.02; 2.51]), the subsequent risk of
194 schizophrenia. In all analyses, the overall risk of developing schizophrenia was higher in
195 female participants (Table S5).

196 When the temporal relationship of vagotomy and schizophrenia was analysed by restricting
197 follow up to 5, 10, 15 and 20 years, we found no statistically significant effect of overall
198 vagotomy at any time point (Table S6). On the contrary, the protective effect of selective
199 vagotomy on the development of schizophrenia was statistically significant at all times (Figure
200 6 and Table S7), with the lowest adjusted HRs found at 5 and 10 years (0.43 [0.19, 0.95] and
201 0.34 [0.17, 0.68], respectively). The association of truncal vagotomy was statistically
202 significant at all times (Figure 6 and Table S7). The effect was highest after 5 years (adjusted
203 HR: 4.04 [2.17, 7.55]) and continuously decreased over time (20 years adjusted HR: 1.95
204 [1.17, 3.24]).

205

206 4 Discussion

207 We built a nationwide retrospective cohort of individuals who underwent vagotomy and
208 contrasted their risk of developing schizophrenia to that of control individuals. Our study
209 included a total of 26,499 individuals identified in the Swedish NPR over a 50-year period. We
210 used a novel approach where controls are matched with vagotomized patients for age, sex
211 and peptic ulcer diagnosis, enabling us to correct for confounding by indication, a challenge
212 faced by previous studies (Bunyozy et al., 2022; Lin et al., 2018; Liu et al., 2017; Liu et al.,
213 2020).

214 An earlier study assessed the risk of developing mental disorders in a nationwide sample of
215 vagotomized individuals in Denmark, and included the risk of developing schizophrenia as a
216 secondary outcome (Bunyozy et al., 2022). With 38 cases identified in the vagotomy group, this
217 study observed no effect of vagotomy on schizophrenia, although with large confidence
218 intervals. Here, we leveraged a larger national register, started the follow-up at an earlier date
219 (which is important given the temporal window in which vagotomies were performed) and used
220 a longer study period. These modifications in the study design yielded more observations and
221 cases than the Danish study (87 in the vagotomy group, 254 overall). Nevertheless, in
222 accordance with this study, we found no effect of vagotomy overall on the subsequent risk of
223 developing schizophrenia.

224 Vagotomy, however, covers several techniques (Seeras et al., 2023): truncal vagotomies
225 sever both vagal trunks at the level of the oesophagus and hence, remove vagal innervation
226 from all subdiaphragmatic organs. In contrast, selective vagotomies only interrupt vagal
227 innervation from the stomach. Our study is the first to distinguish between the two types of
228 vagotomies for the outcome of schizophrenia. Intriguingly, we found that truncal vagotomy
229 significantly increases, while selective vagotomy moderately decreases the risk of developing
230 schizophrenia compared to matched controls. Interestingly, Bunyozy et al. also found that
231 truncal vagotomy, but not selective vagotomy, was associated with an increase in mental
232 disorders (Bunyozy et al., 2022).

233 A first potential explanation of these differences may lie in the distinct indication between
234 selective and truncal vagotomy. Selective vagotomy is a more subtle lesion with less side
235 effects, but is associated with a higher recurrence rate than truncal vagotomy. Hence, truncal
236 vagotomy was primarily performed in patients where recurrence rate needed to be minimized
237 despite side effects, such as elderly patients or patients with comorbidities (Chan et al., 1994).
238 It has been argued that these patients are more likely to visit the hospital after surgery than
239 individuals with selective vagotomy and this could explain the different effects of vagotomy

240 types on the development of schizophrenia (surveillance bias) (Bunyoz et al., 2022). Here, we
241 matched controls for age, sex and ulcer diagnosis, which mitigates this bias. Moreover, the
242 effect of truncal vagotomy on other outcomes (inflammatory bowel disease, dementia) tends
243 to be similar to that of selective vagotomy (Lin et al., 2018; Liu et al., 2020). Truncal vagotomy
244 even reduces the risk of developing Parkinson's disease, while selective vagotomy had no
245 effect (Liu et al., 2017). Hence, the surveillance bias explanation seems unlikely and the
246 observed differences might be rooted in biology.

247 Importantly, the difference between selective and truncal vagotomy on the postoperative
248 development of schizophrenia does not only lie in the extent of the lesion. Both surgeries also
249 target distinct neuronal populations. Recent rodent findings confirmed the notion that distinct
250 vagal neuronal populations have diverse contributions to the same behavior (Tao et al., 2021).
251 For example, only a subset of vagal sensory neurons projecting throughout the gastrointestinal
252 tract contributes to eating behavior, or reward behavior modulation (Bai et al., 2019; Han et
253 al., 2018). Distinct vagal populations can even have opposing actions, as exemplified by two
254 vagal sensory subtypes that induce apnea or rapid breathing (Chang et al., 2015). Hence, it
255 is plausible that the distinct vagal neurons lesioned by truncal or selective vagotomies exert
256 differential effects on schizophrenia-relevant behaviors. Future studies could exploit this
257 emerging granularity in our understanding of vagal circuits to investigate the role of distinct
258 neuronal populations and projections on schizophrenia-like behaviors, for example using
259 rodent models.

260 Vagal neurons, specifically vagal afferent neurons, are well positioned to modulate central
261 pathways involved in the pathophysiology of schizophrenia. In the rat, a vagal lesion that
262 primarily affects subdiaphragmatic vagal afferent neurons, was associated with brain
263 transcriptomic changes annotating with schizophrenia-associated disease pathways and
264 increased striatal dopamine content (Klarer et al., 2018). Interestingly, increased dopamine
265 content within striatal structures is a core pathophysiological mechanism in schizophrenia
266 (Howes and Kapur, 2009). It is thought to underlie the disruption of attentional control of
267 associative learning (Murphy et al., 2000), sensorimotor gating (Swerdlow et al., 1990;
268 Swerdlow et al., 1992; Wan et al., 1995), and amphetamine hypersensitivity (Heidbreder and
269 Feldon, 1998). From a circuit perspective, ascending vagal inputs to the nucleus of the solitary
270 tract are synaptically connected to the ventral tegmental area (Han et al., 2018), which
271 contains the majority of dopaminergic cells projecting to the ventral striatum. Beyond
272 mesolimbic dopamine, future studies should examine the contribution of vagal neurons to
273 schizophrenia-relevant brain circuits in a more systematic manner.

274 Finally, the relevance of our findings extends beyond the specific case of vagotomized
275 individuals. Reduced parasympathetic activity, evidenced by a low heart rate variability, is a
276 common feature of many mental and metabolic disorders, suggesting that descending vagal
277 activity is also affected in a large population of patients (Ramesh et al., 2023). In addition,
278 vagal afferent neurons show reduced response to gastrointestinal hormones or stretch in
279 animal models of obesity (Kentish et al., 2011; Kentish et al., 2014; Kentish and Page, 2014;
280 Kentish et al., 2016) or animals with microbial dysbiosis (Jameson et al., 2023). Hence, if these
281 findings extend to humans, the number of individuals affected by impairments in vagal
282 ascending pathways is likely to be large.

283 Overall, our study confirms the absence of effect of overall vagotomy on schizophrenia seen
284 in a previous study. It opens, however, the possibility that distinct vagal circuits play a
285 differential role in the risk of developing schizophrenia. Notably, our findings suggest a role for
286 the non-gastric subdiaphragmatic branches of the vagus nerve in the development of
287 schizophrenia. While future epidemiological studies are warranted to strengthen this
288 conclusion, studies in rodents are well-suited to test this hypothesis and uncover the
289 underlying mechanisms.

290 **5 Conflicts of interests**

291 The authors have no conflicts to declare.

292 **6 Funding**

293 This research was funded by the Swedish Research Council (2018-00660 to KPS and 2021-
294 01549 to JPK), the National Institute of Health (R01DK129321 to KPS) and the Swiss National
295 Science Foundation (183899 to JPK).

296 **7 Author contributions**

297 CFR: Methodology, Formal analysis, Investigation, Visualization, Writing-Original draft
298 preparation

299 KPS: Conceptualization, Writing-Reviewing and Editing

300 SR: Conceptualization, Writing-Reviewing and Editing

301 UM: Writing-Reviewing and Editing

302 JPK: Conceptualization, Methodology, Writing-Original draft preparation, Writing-Reviewing
303 and Editing

304

305 **8 References**

- 306 Bai, L., Mesgarzadeh, S., Ramesh, K.S., Huey, E.L., Liu, Y., Gray, L.A., Aitken, T.J., Chen,
307 Y., Beutler, L.R., Ahn, J.S., Madisen, L., Zeng, H., Krasnow, M.A., Knight, Z.A., 2019.
308 Genetic Identification of Vagal Sensory Neurons That Control Feeding. *Cell* 179(5), 1129-
309 1143 e1123.
- 310 Bonaz, B., Sinniger, V., Pellissier, S., 2016. Anti-inflammatory properties of the vagus nerve:
311 potential therapeutic implications of vagus nerve stimulation. *J Physiol* 594(20), 5781-5790.
- 312 Bunyoz, A.H., Christensen, R.H.B., Orlovskaa-Waast, S., Nordentoft, M., Mortensen, P.B.,
313 Petersen, L.V., Benros, M.E., 2022. Vagotomy and the risk of mental disorders: A nationwide
314 population-based study. *Acta Psychiatr Scand* 145(1), 67-78.
- 315 Chan, V.M., Reznick, R.K., O'Rourke, K., Kitchens, J.M., Lossing, A.G., Detsky, A.S., 1994.
316 Meta-analysis of highly selective vagotomy versus truncal vagotomy and pyloroplasty in the
317 surgical treatment of uncomplicated duodenal ulcer. *Can J Surg* 37(6), 457-464.
- 318 Chang, R.B., Strohlic, D.E., Williams, E.K., Umans, B.D., Liberles, S.D., 2015. Vagal
319 Sensory Neuron Subtypes that Differentially Control Breathing. *Cell* 161(3), 622-633.
- 320 Charlson, M.E., Pompei, P., Ales, K.L., MacKenzie, C.R., 1987. A new method of classifying
321 prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*
322 40(5), 373-383.
- 323 Corsi-Zuelli, F., Brognara, F., Quirino, G., Hiroki, C.H., Fais, R.S., Del-Ben, C.M., Ulloa, L.,
324 Salgado, H.C., Kanashiro, A., Loureiro, C.M., 2017. Neuroimmune Interactions in
325 Schizophrenia: Focus on Vagus Nerve Stimulation and Activation of the Alpha-7 Nicotinic
326 Acetylcholine Receptor. *Front Immunol* 8, 618.
- 327 Deyo, R.A., Cherkin, D.C., Ciol, M.A., 1992. Adapting a clinical comorbidity index for use
328 with ICD-9-CM administrative databases. *J Clin Epidemiol* 45(6), 613-619.
- 329 Dinan, T.G., Borre, Y.E., Cryan, J.F., 2014. Genomics of schizophrenia: time to consider the
330 gut microbiome? *Mol Psychiatry* 19(12), 1252-1257.
- 331 Han, W., Tellez, L.A., Perkins, M.H., Perez, I.O., Qu, T., Ferreira, J., Ferreira, T.L., Quinn,
332 D., Liu, Z.-W., Gao, X.-B., Kaelberer, M.M., Bohórquez, D.V., Shammah-Lagnado, S.J., de
333 Lartigue, G., de Araujo, I.E., 2018. A Neural Circuit for Gut-Induced Reward. *Cell*.
- 334 Heidbreder, C., Feldon, J., 1998. Amphetamine-induced neurochemical and locomotor
335 responses are expressed differentially across the anteroposterior axis of the core and shell
336 subterritories of the nucleus accumbens. *Synapse* 29(4), 310-322.
- 337 Howes, O.D., Kapur, S., 2009. The dopamine hypothesis of schizophrenia: version III--the
338 final common pathway. *Schizophr Bull* 35(3), 549-562.
- 339 Jablensky, A., 2010. The diagnostic concept of schizophrenia: its history, evolution, and
340 future prospects. *Dialogues Clin Neurosci* 12(3), 271-287.
- 341 Jameson, K.G., Kazmi, S.A., Son, C., Mazdeyasnan, D., Leshan, E., Vuong, H.E., Paramo,
342 J., Lopez-Romero, A., Yang, L., Schweizer, F.E., Hsiao, E.Y., 2023. Vagal interoception of
343 microbial metabolites from the small intestinal lumen. *bioRxiv*.
- 344 Kentish, S., Li, H., Philp, L.K., O'Donnell, T.A., Isaacs, N.J., Young, R.L., Wittert, G.A.,
345 Blackshaw, L.A., Page, A.J., 2011. Diet induced adaptation of vagal afferent function. *The*
346 *Journal of Physiology*, no-no.
- 347 Kentish, S.J., O'Donnell, T.A., Frisby, C.L., Li, H., Wittert, G.A., Page, A.J., 2014. Altered
348 gastric vagal mechanosensitivity in diet-induced obesity persists on return to normal chow
349 and is accompanied by increased food intake. *Int J Obes (Lond)* 38(5), 636-642.

- 350 Kentish, S.J., Page, A.J., 2014. Plasticity of gastro-intestinal vagal afferent endings. *Physiol*
351 *Behav* 136, 170-178.
- 352 Kentish, S.J., Vincent, A.D., Kennaway, D.J., Wittert, G.A., Page, A.J., 2016. High-Fat Diet-
353 Induced Obesity Ablates Gastric Vagal Afferent Circadian Rhythms. *J Neurosci* 36(11),
354 3199-3207.
- 355 Klarer, M., Krieger, J.-P., Richetto, J., Weber-Stadlbauer, U., Günther, L., Winter, C., Arnold,
356 M., Langhans, W., Meyer, U., 2018. Abdominal Vagal Afferents Modulate the Brain
357 Transcriptome and Behaviors Relevant to Schizophrenia. *The Journal of neuroscience : the*
358 *official journal of the Society for Neuroscience*.
- 359 Lin, S.Y., Lin, C.L., Wang, I.K., Lin, C.C., Lin, C.H., Hsu, W.H., Kao, C.H., 2018. Dementia
360 and vagotomy in Taiwan: a population-based cohort study. *BMJ Open* 8(3), e019582.
- 361 Liu, B., Fang, F., Pedersen, N.L., Tillander, A., Ludvigsson, J.F., Ekbom, A., Svenningsson,
362 P., Chen, H., Wirdefeldt, K., 2017. Vagotomy and Parkinson disease: A Swedish register-
363 based matched-cohort study. *Neurology* 88(21), 1996-2002.
- 364 Liu, B., Wanders, A., Wirdefeldt, K., Sjolander, A., Sachs, M.C., Eberhardson, M., Ye, W.,
365 Ekbom, A., Olen, O., Ludvigsson, J.F., 2020. Vagotomy and subsequent risk of inflammatory
366 bowel disease: a nationwide register-based matched cohort study. *Aliment Pharmacol Ther*
367 51(11), 1022-1030.
- 368 Ludvigsson, J.F., Andersson, E., Ekbom, A., Feychting, M., Kim, J.L., Reuterwall, C.,
369 Heurgren, M., Olausson, P.O., 2011. External review and validation of the Swedish national
370 inpatient register. *BMC Public Health* 11, 450.
- 371 Ludvigsson, J.F., Appelros, P., Askling, J., Byberg, L., Carrero, J.J., Ekstrom, A.M., Ekstrom,
372 M., Smedby, K.E., Hagstrom, H., James, S., Jarvholm, B., Michaelsson, K., Pedersen, N.L.,
373 Sundelin, H., Sundquist, K., Sundstrom, J., 2021. Adaptation of the Charlson Comorbidity
374 Index for Register-Based Research in Sweden. *Clin Epidemiol* 13, 21-41.
- 375 Lundell, L., 2011. Acid secretion and gastric surgery. *Dig Dis* 29(5), 487-490.
- 376 Murphy, C.A., Pezze, M., Feldon, J., Heidbreder, C., 2000. Differential involvement of
377 dopamine in the shell and core of the nucleus accumbens in the expression of latent
378 inhibition to an aversively conditioned stimulus. *Neuroscience* 97(3), 469-477.
- 379 Neuhuber, W.L., Berthoud, H.R., 2021. Functional anatomy of the vagus system - Emphasis
380 on the somato-visceral interface. *Auton Neurosci* 236, 102887.
- 381 Ramesh, A., Nayak, T., Beestrup, M., Quer, G., Pandit, J.A., 2023. Heart Rate Variability in
382 Psychiatric Disorders: A Systematic Review. *Neuropsychiatr Dis Treat* 19, 2217-2239.
- 383 Seeras, K., Qasawa, R.N., Prakash, S., 2023. Truncal Vagotomy, StatPearls, Treasure
384 Island (FL) ineligible companies. Disclosure: Ryan Qasawa declares no relevant financial
385 relationships with ineligible companies. Disclosure: Shivana Prakash declares no relevant
386 financial relationships with ineligible companies.
- 387 Socialstyrelsen, 2024. National Patient Register; [https://www.socialstyrelsen.se/en/statistics-](https://www.socialstyrelsen.se/en/statistics-and-data/registers/national-patient-register)
388 [and-data/registers/national-patient-register](https://www.socialstyrelsen.se/en/statistics-and-data/registers/national-patient-register). Consulted January 2024.
- 389 Stogios, N., Gdanski, A., Gerretsen, P., Chintoh, A.F., Graff-Guerrero, A., Rajji, T.K.,
390 Remington, G., Hahn, M.K., Agarwal, S.M., 2021. Autonomic nervous system dysfunction in
391 schizophrenia: impact on cognitive and metabolic health. *NPJ Schizophr* 7(1), 22.
- 392 Swerdlow, N.R., Braff, D.L., Geyer, M.A., 1990. GABAergic projection from nucleus
393 accumbens to ventral pallidum mediates dopamine-induced sensorimotor gating deficits of
394 acoustic startle in rats. *Brain Res* 532(1-2), 146-150.

- 395 Swerdlow, N.R., Caine, S.B., Geyer, M.A., 1992. Regionally selective effects of intracerebral
396 dopamine infusion on sensorimotor gating of the startle reflex in rats. *Psychopharmacology*
397 (Berl) 108(1-2), 189-195.
- 398 Tao, J., Campbell, J.N., Tsai, L.T., Wu, C., Liberles, S.D., Lowell, B.B., 2021. Highly
399 selective brain-to-gut communication via genetically defined vagus neurons. *Neuron*
400 109(13), 2106-2115 e2104.
- 401 Thirion, F., Speyer, H., Hansen, T.H., Nielsen, T., Fan, Y., Le Chatelier, E., Fromentin, S.,
402 Berland, M., Plaza Onate, F., Pons, N., Galleron, N., Levenez, F., Marko, L., Birkner, T.,
403 Jorgensen, T., Forslund, S.K., Vestergaard, H., Hansen, T., Nordentoft, M., Mors, O.,
404 Benros, M.E., Pedersen, O., Ehrlich, S.D., 2023. Alteration of Gut Microbiome in Patients
405 With Schizophrenia Indicates Links Between Bacterial Tyrosine Biosynthesis and Cognitive
406 Dysfunction. *Biol Psychiatry Glob Open Sci* 3(2), 283-291.
- 407 Tsuang, M.T., Lyons, M.J., Toomey, R., 2000. Chapter 13 Heterogeneity of schizophrenia,
408 *Biological Psychiatry*, pp. 263-277.
- 409 von Elm, E., Altman, D.G., Egger, M., Pocock, S.J., Gotsche, P.C., Vandenbroucke, J.P.,
410 2007. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE)
411 statement: guidelines for reporting observational studies. *BMJ* 335(7624), 806-808.
- 412 Wan, F.J., Geyer, M.A., Swerdlow, N.R., 1995. Presynaptic dopamine-glutamate interactions
413 in the nucleus accumbens regulate sensorimotor gating. *Psychopharmacology (Berl)* 120(4),
414 433-441.
- 415 Zhu, F., Ju, Y., Wang, W., Wang, Q., Guo, R., Ma, Q., Sun, Q., Fan, Y., Xie, Y., Yang, Z.,
416 Jie, Z., Zhao, B., Xiao, L., Yang, L., Zhang, T., Feng, J., Guo, L., He, X., Chen, Y., Chen, C.,
417 Gao, C., Xu, X., Yang, H., Wang, J., Dang, Y., Madsen, L., Brix, S., Kristiansen, K., Jia, H.,
418 Ma, X., 2020. Metagenome-wide association of gut microbiome features for schizophrenia.
419 *Nat Commun* 11(1), 1612.
- 420

421 **Table 1. Patient characteristics**

	<i>Overall</i>	<i>No vagotomy</i>	<i>Vagotomy</i>	<i>Selective vagotomy</i>	<i>Truncal vagotomy</i>
n	26 499	17 666	8 833	6 524	2 304
Surgery Year Median [IQR]	1982 [1976, 1987]	1982 [1976,1987]	1982 [1976, 1987]	1980 [1975, 1986]	1984 [1981, 1989]
Age in years Median [IQR]	54 [43, 65]	54 [43, 65]	54 [43, 65]	51 [41, 60]	68 [56, 76]
Sex Females (n, %)	9 006 (34.0)	6 004 (34.0)	3 002 (34.0)	2 184 (33.4)	814 (35.3)
Death during study period (n, %)	21 952 (82.8)	15 101 (85.5)	6 851 (77.6)	4 724 (72.4)	2 123 (92.1)
Schizophrenia diagnosis (n, %)	254 (1.0)	167 (0.9)	87 (1.0)	65 (1.0)	22 (1.0)
Ulcer Diagnosis					
Gastric ulcer	8 451 (31.9)	5 634 (31.9)	2 817 (31.9)	1 903 (29.2)	910 (39.5)
Duodenal ulcer	15 771 (59.5)	10 514 (59.5)	5 257 (59.5)	4 068 (62.4)	1 188 (51.6)
Gastrojejunal ulcer	594 (2.2)	396 (2.2)	198 (2.2)	127 (2.0)	71 (3.1)
Not specified ulcer	1 683 (6.4)	1 122 (6.4)	561 (6.4)	426 (6.5)	135 (2.1)
Adapted CCI (n, %)					
0	25 875 (97.6)	17 478 (98.9)	8 379 (95.1)	6 351 (97.3)	2 041 (88.6)
1-2	323 (1.2)	94 (0.5)	229 (2.6)	85 (1.3)	144 (6.2)
>=3	301 (1.1)	94 (0.5)	207 (2.3)	88 (1.3)	119 (5.2)

422


423 *N = 5 vagotomy patients with vagotomy = not defined, not shown. IQR = Interquartile Range.*

424 *CCI = Charlson Comorbidity Index.*

425

426 **Figure 1. Flowchart**


427 *NPR = National Patient Registry*



428

429


430 **Figure 2. Cumulative incidence function of schizophrenia among patients with overall**
 431 **vagotomy and matched controls.**



432

433

434 **Figure 3. Cumulative incidence function of schizophrenia among patients with truncal**
 435 **or selective vagotomy and matched controls.**




436

437


438

439 **Figure 4. Regression Results of Cox proportional hazards model for overall vagotomy**
440 **and subsequent diagnosis of schizophrenia.**




441
442 *Crude model controlled age and sex. Adjusted model controlled for age, sex, weighted*
443 *Charlson Comorbidity Index, ICD-period. HR = Hazard Ratio, 95% CI = 95% Confidence*
444 *Interval. Vertical line: reference value of 1*
445

446 **Figure 5. Regression results of Cox proportional hazards model for selective or truncal**
 447 **vagotomy and subsequent diagnosis of schizophrenia.**



448
 449 *Crude model controlled age and sex. Adjusted model controlled for age, sex, weighted*
 450 *Charlson Comorbidity Index, ICD-period. HR = Hazard Ratio, 95% CI = 95% Confidence*
 451 *Interval. Vertical line: reference value of 1*
 452

453 **Figure 6: Regression results of Cox proportional hazards model for selective or truncal**
454 **vagotomy and schizophrenia, with follow-up restriction.**



455

456

457 *Crude model controlled age and sex. Adjusted model controlled for age, sex, weighted*
458 *Charlson Comorbidity Index, ICD-period. HR = Hazard Ratio, 95% CI = 95% Confidence*
459 *Interval. Vertical line: reference value of 1*

460

461