


APPENDICES



	PCU-H (N = 77)	Controls (N = 137)	P value
Maternal age (years)	25.6 [18.8-37.4]	27.8 [18.7-40.5]	0.0033
AMA	1 (1.3%)	12 (8.63%)	0.03
Teen	8 (10.4%)	6 (4.32%)	0.082
Maternal BMI at intake (kg/m ²)	28.7 [17.1-57.3]	33 [14.8-79.1]	0.0049
Education beyond high school	26 (33.8%)	66 (50%)	0.023
Marital status			0.495
Single	20 (60.6%)	32 (54.2%)	
Partnered/Married	11 (33.3%)	25 (42.4%)	
Divorced/Separated	1 (3%)	2 (3.4%)	
Widowed	0	0	
Other	1 (3%)	0	
ADI (national)	81.4 [31-100]	79.3 [17-100]	0.66
ADI >=75%ile			0.490
Race			0.59
Black or African American	66 (85.7%)	113 (81.3%)	
White	11 (14.3%)	23 (16.6%)	
American Indian/Alaska N	0	0	
Asian	0	2 (1.4%)	
Hawaiian N. & Pacific Is.	0	0	
Other	0	1 (0.7%)	
Ethnicity			0.059

Hispanic	1 (1.3%)	10 (7.2%)	
Non-Hispanic	76 (98.7%)	129 (92.8%)	
Gravity (median, [range])	2.7 [1-6]	3 [1-6]	0.4183
Prior live births (median, [range])	1 [0-4]	1 [0-5]	0.8915
Concurrent medical conditions			
Chronic hypertension	14 (18.4%)	36 (27.1%)	0.159
Chronic diabetes	1 (1.45%)	9 (6.7%)	0.083
Asthma	21 (27.6%)	42 (30.9%)	0.619
Anemia	25 (33.3%)	51 (37.5%)	0.546
Psychiatric diagnosis	32 (42.1%)	68 (50%)	0.269
Prior obstetric outcomes			
Preterm birth	15 (19.7%)	24 (17.7%)	0.706
Hypertensive disorders of pregnancy	17 (22.4%)	21 (15.6%)	0.216
Gestational diabetes	2 (2.7%)	5 (4%)	0.643
Cotinine positivity	42 (54.6%)	15 (10.8%)	<0.00
Gestational age at delivery (weeks)	37.5 [27-41]	37.5 [23-40]	0.6370
Attendance at 3 or more prenatal visits	75 (99.7%)	136 (99.3%)	0.671
Abbreviations: ADI, area of deprivation index; BMI, body mass index; kg, kilograms; m, meters; PCU-H, high prenatal cannabis use			
*Other indicates that the participant did not self-report a specific race			

	PCU-H (n=77)	Controls (n=139)	RR (95% CI)	*aRR (95% CI)
Antenatal Complications				
Gestational diabetes	0 (0%)	10 (8.0%)	N/A	N/A
Anemia	21 (87.5%)	40 (80%)	1.09 [0.89-1.34]	1.06 [0.68-1.20]
Gestational hypertension	19 (25%)	29 (21.6%)	1.16 [0.7-1.91]	1.05 [0.54-1.82]
Preeclampsia	11 (14.7%)	21 (15.7%)	0.93 [0.48-1.84]	1.19 [0.52-2.39]
Fetal growth restriction	8 (11%)	14 (10.5%)	1.04 [0.46-2.38]	1.22 [0.44-2.98]
Preterm delivery	17 (22.4%)	27 (19.4%)	1.15 [0.67-1.97]	1.54 [0.81-2.54]
Cesarean delivery	15 (20%)	37 (27.8%)	0.72 [0.42-1.22]	0.84 [0.43-1.46]
Breastfeeding at discharge	60 (80%)	97 (74.1%)	1.08 [0.93-1.28]	1.08 [0.87-1.21]
Attendance at 1 or more postpartum visits	42 (54.6%)	77 (55.4%)	0.98 [0.76-1.27]	0.92 [0.63-1.21]
Abbreviations: adjusted relative risk; CI, confidence interval; N/A, not applicable; PCU-H, high prenatal cannabis use; RR, relative risk				
*aRR for maternal outcomes were adjusted for area of deprivation index >= 75%ile, obesity, and cotinine positivity				

	PCU-H (n=77)	Controls (n=139)	RR (95% CI)	*aRR (95% CI)
SGA < 5ile	10 (13.2%)	6 (4.32%)	3.05 [1.15-8.06]	1.44 [0.39-4.68]
SGA < 10ile	23 (30.3%)	18 (13%)	2.34 [1.35-4.05]	1.53 [0.72-2.88]
LGA	1 (1.3%)	8 (5.8%)	0.23 [0.03-1.79]	0.16 [0.02-1.59]
5 min Agpar < 7	1 (1.3%)	9 (6.6%)	0.2 [0.03-1.54]	0.15 [0.02-1.34]

NICU admission	11 (14.9%)	24 (17.7%)	0.84 [0.44-1.62]	0.81 [0.35-1.71]
	PCU-H (n=77)	PCU-L (n=180)	RR (95% CI)	*aRR (95% CI)
SGA < 5ile	10 (13.2%)	21 (11.8%)	1.11 [0.55-2.25]	0.73 [0.29-1.71]
SGA < 10ile	23 (30.3%)	43 (24.2%)	1.25 [0.81-1.92]	1.05 [0.6-1.69]
LGA	1 (1.3%)	4 (2.3%)	0.59 [0.07-5.15]	0.77 [0.08-6.27]
5 min Agpar < 7	1 (1.3%)	8 (4.9%)	0.27 [0.03-2.12]	0.28 [0.03-2.12]
NICU admission	11 (14.9%)	32 (18.7%)	0.79 [0.42-1.49]	0.82 [0.39-1.61]
	PCU-L (n=180)	Controls (n=139)	RR (95% CI)	*aRR (95% CI)
SGA < 5ile	21 (11.8%)	6 (4.3%)	2.73 [1.13-6.59]	1.8 [0.69-4.31]
SGA < 10ile	43 (24.2%)	18 (13%)	1.87 [1.12-3.09]	1.31 [0.71-2.24]
LGA	4 (2.3%)	8 (5.8%)	0.39 [0.12-1.27]	0.45 [0.12-1.59]
5 min Agpar < 7	8 (4.9%)	9 (6.6%)	0.74 [0.29-1.86]	0.66 [0.24-1.71]
NICU admission	32 (18.7%)	24 (17.7%)	1.06 [0.66-1.71]	0.97 [0.5-1.74]
Abbreviations: aRR, adjusted relative risk ; CI, confidence interval; N/A, not applicable; LGA, large for gestational age; NICU, neonatal intensive care unit; PCU-H, high prenatal cannabis use; PCU-L, low prenatal cannabis use; RR, relative risk; SGA, small for gestational age				
*aRR for neonatal outcomes were adjusted for area of deprivation index >= 75%ile and cotinine positivity				

	PCU+c (N= 110)	PCU-c (N= 147)	P value
Maternal age in years (median, [range])	24.9 [18.8-38.6]	26.1 [19.1-39.2]	0.093
AMA	3 (2.7%)	7 (4.8%)	0.404
Teen	10 (9.1%)	11 (7.5%)	0.641
Maternal BMI at intake (kg/m ²) (median, [range])	28.2 [15.5-66.6]	27.1 [15.8-58.1]	0.929
Education beyond high school	26 (25.5%)	44 (32.4%)	0.25
Marital status			0.208
Single	20 (66.7%)	30 (51.7%)	
Partnered/Married	9 (30%)	27 (46.6%)	
Divorced/Separated	1 (3.3%)	0	
Widowed	0	0	
Other	0	1 (1.7%)	
ADI (national)	94 [31-100]	89 [34-100]	0.017
ADI >=75%ile	88 (86.3%)	100 (74.1%)	0.022
Race			0.413
Black or African American	100 (91.7%)	127 (87%)	
White	9 (8.3%)	16 (11%)	
American Indian/Alaska N	0	2 (1.4%)	
Asian	0	0	
Hawaiian N. & Pacific Is.	0	0	
Other	0	1 (0.7%)	
Ethnicity			
Hispanic	2 (1.8%)	3 (2.0%)	

Non-Hispanic	107 (98.2%)	144 (98%)	
Gravity (median, [range])	2 (1-6)	2 (1-6)	0.971
Prior live births (median, [range])	1 (0-5)	1 (0-5)	0.677
Concurrent medical conditions			
Chronic hypertension	21 (20.2%)	30 (21.6%)	0.792
Chronic diabetes	2 (1.9%)	5 (3.7%)	0.406
Asthma	33 (31.4%)	43 (31%)	0.934
Anemia	39 (37.1%)	56 (40.6%)	0.587
Psychiatric diagnosis	55 (52.4%)	68 (49%)	0.592
Prior obstetric outcomes			
Preterm birth	25 (23.4%)	28 (19.4%)	0.452
Hypertensive disorders of pregnancy	23 (22.1%)	15 (10.8%)	0.016
Gestational diabetes	4 (3.9%)	4 (3.1%)	0.745
Gestational age at delivery (weeks)	39 [24-40]	38 [27-41]	0.608
Attendance at 3 or more prenatal visits	104 (99.1%)	136 (98.6%)	0.728
Abbreviations: ADI, area of deprivation index; BMI, body mass index; kg, kilograms; m, meters; PCU+c, prenatal cannabis use with cotinine positivity; PCU-c, prenatal cannabis use with negative cotinine			
*Other indicates that the participant did not self-report a specific race			

Table 8: Birthing person outcomes among PCU patients by cotinine status

	PCU+cot (n=110)	PCU-cot (n=147)	RR (95% CI)	*aRR (95% CI)
Antenatal Complications				
Gestational diabetes	1 (1.0%)	6 (4.4%)	0.22 [0.02-1.8]	0.21 [0.02-1.78]
Anemia	32 (88.9%)	50 (89.3%)	1.0 [0.86-1.15]	0.93 [0.6-1.06]
Gestational hypertension	24 (23.3%)	22 (16.1%)	1.45 [0.86-1.44]	1.44 [0.83-2.31]
Preeclampsia	10 (9.8%)	21 (15.4%)	0.63 [0.31-1.29]	0.66 [0.30-1.33]
Fetal growth restriction	11 (10.5%)	13 (9.4%)	1.12 [0.52-2.4]	1.16 [0.49-2.51]
Preterm delivery	16 (14.7%)	26 (17.9%)	0.82 [0.46-1.45]	0.83 [0.43-1.52]
Cesarean delivery	20 (19.4%)	33 (24.3%)	0.80 [0.49-1.31]	0.81 [0.46-1.35]
Breastfeeding at discharge	87 (83.7%)	106 (80.9%)	1.03 [0.92-1.16]	1.03 [0.88-1.13]
Attendance at 1 or more postpartum visits	75 (68.2%)	80 (54.4%)	1.25 [1.03-1.52]	1.25 [1.00-1.45]
Abbreviations: adjusted relative risk; CI, confidence interval; N/A, not applicable; PCU+c, prenatal cannabis use with cotinine positivity; PCU-c, prenatal cannabis use with negative cotinine; RR, relative risk				
*aRR for maternal outcomes were adjusted for area of deprivation index >= 75%ile and obesity				

Table 9: Neonatal outcomes by cotinine status

	PCU+cot (n=110)	PCU-cot (n=147)	RR (95% CI)	*aRR (95% CI)
SGA < 5ile	19 (17.4%)	12 (8.3%)	2.1 [1.07-4.15]	2.7 [1.21-5.38]
SGA < 10ile	39 (35.8%)	27 (18.6%)	1.92 [1.26-2.93]	2.4 [1.51-3.48]
LGA	0 (0)	5 (3.5%)	N/A	N/A
5 min Agpar < 7	4 (3.9%)	5 (3.6%)	1.08 [0.3-3.93]	0.94 [0.25-3.27]
NICU admission	18 (17.4%)	25 (17.9%)	0.96 [0.55-1.66]	0.84 [0.43-1.53]

	PCU+cot (n=110)	Controls (n=124)	RR (95% CI)	*aRR (95% CI)
SGA < 5ile	19 (17.4%)	6 (4.8%)	3.60 [1.50-8.69]	3.29 [1.33-6.95]
SGA < 10ile	39 (35.8%)	16 (12.9%)	2.77 [1.64-4.67]	2.75 [1.64-4.13]
LGA	0 (0%)	6 (4.8%)	Unable to calc	Unable to calc
5 min Agpar < 7	4 (3.9%)	8 (6.6%)	0.59 [0.18-1.91]	0.53 [0.16-1.69]
NICU admission	18 (17.1%)	22 (18.2%)	0.94 [0.54-1.66]	0.78 [0.4-1.44]
	PCU-cot (n=110)	Controls (n=124)	RR (95% CI)	*aRR (95% CI)
SGA < 5ile	12 (8.3%)	6 (4.8%)	1.71 [0.66-4.42]	1.15 [0.40-3.05]
SGA < 10ile	27 (18.6%)	16 (12.9%)	1.44 [0.82-2.55]	1.15 [0.59-2.08]
LGA	5 (3.5%)	6 (4.8%)	0.71 [0.22-2.28]	0.87 [0.25-2.81]
5 min Agpar < 7	8 (6.6%)	5 (3.6%)	0.55 [0.18-1.63]	0.51 [0.16-1.49]
NICU admission	22 (18.2%)	25 (17.9%)	0.98 [0.58-1.65]	0.87-0.48]
Abbreviations: aRR, adjusted relative risk ; CI, confidence interval; N/A, not applicable; LGA, large for gestational age; NICU, neonatal intensive care unit; PCU+c, prenatal cannabis use with cotinine positivity; PCU-c, prenatal cannabis use with negative cotinine; RR, relative risk; SGA, small for gestational age				
*aRR for neonatal outcomes were adjusted for area of deprivation index >= 75%ile				