# 1 DRAFT ORIGINAL ARTICLE

# 2 THE INDONESIA HEALTH WORKFORCE QUANTITY AND DISTRIBUTION

- 3 Farizal Rizky Muharram<sup>1</sup>, Hanif Ardiansyah Sulistya<sup>2</sup>, Julian Benedict Swannjo<sup>2</sup>, Fikri Febrian
- 4 Firmansvah<sup>2</sup>, Muhammad Masrur Rizal<sup>2</sup>, Alifina Izza<sup>2</sup>, Muhammad Atoillah Isfandiari<sup>3</sup>, Ninuk
- 5 Dwi Ariningtyas<sup>4</sup>, Achmad Chusnu Romdhoni<sup>5\*</sup>.

# 6 AFFILIATION

- 7 1 Department of Global Health and Social Medicine, Harvard Medical School, Boston, MA
- 8 2 Advanced Research and Collaboration Center, Surabaya, Indonesia
- 9 3 Department of Epidemiology, Biostatistics and Demography, Health Promotion and Behavioral
- 10 Science Faculty of Public Health, Surabaya, Indonesia
- 11 4 Department of Medicinal Education, Faculty of Medicine, Universitas Muhammadiyah
- 12 Surabaya, Indonesia
- 13 5 Department of Otorhinolaryngology-Head-and-Neck Surgery, Airlangga University-Dr. Soetomo
- 14 General Academic Hospital, Surabaya, Indonesia
- 15
- 16 \*Corresponding Author: Achmad Chusnu Romdhoni
- 17
- 18 Short Title: Indonesia Health Workforce Quantity and Distribution

# 19 ABBREVIATION

- 20 HRH: Human Research for Health; HW: Health Worker, NHS: National Health Services; GP:
- 21 General Practitioner; UHC: Universal Health Coverage

# 22 ABSTRACT

Background: Indonesia, the world's largest archipelago, faces unique challenges in distributing its health workforce across its diverse geographic barriers, leading to disparities in health worker number and distribution. By dissecting the distribution patterns and identifying areas of critical need, the research seeks to inform policy interventions that can more effectively bridge the gap on health worker quantity and inequity.

Methods: We conducted a descriptive analysis of healthcare workforce data across all 514 districts in Indonesia. The study focused on five categories of health workers: General Practitioners (GPs), medical specialists, dentists, nurses, and midwives. We calculated the health worker ratio to determine the availability of healthcare workers relative to the population. To evaluate the distribution of these workers, we employed the Gini Index as a measure of distribution equality. Additionally, we conducted a comparative metric approach to assess both the quantity and the equity of healthcare worker distribution across the districts.

35 **Results:** In Indonesia, the current health worker ratio stands at 3.84 per 1000 population, falling 36 short of the WHO's threshold of 4.45 for achieving 80% Universal Health Coverage. This shortfall 37 translates to a need for an additional 166,000 health workers. Our analysis reveals a varied 38 distribution of health worker categories: while midwives show a relatively equitable distribution. 39 specialists and dentists exhibit significant inequality, especially at the district level. The Gini Index, 40 used to measure this inequality, indicates greater disparities at the district level compared to the 41 provincial level. There has been notable progress in the distribution of medical specialists across 42 provinces, with the between-provinces Gini Index for specialists decreasing from 0.57 in 1993 to 43 0.44 in 2022. However, the inter-district Gini Index remains high at 0.53 in 2022, signifying a 44 concentration of specialists in major cities and provincial capitals.

45 **Conclusion:** This study shows that human resources for health in Indonesia suffer not only in

46 quantity but also in distribution. Our finding underscores the importance of considering inter-

47 province and inter-district disparities to tailor policies to tackle unique problems each region faces.

48 **Keywords:** Health Workforce; Quantity; Distribution; Gini Index

#### **Evidence Before Study**

- Prior research has established that the quantity and distribution of health workers are critical factors in improving life expectancy and are fundamental components of the health system.
- Following Indonesia's constitutional changes in 2001, which included the autonomy and decentralization of healthcare services, assessing the number of health workers at the district level has gained significant importance for determining national healthcare needs.
- There has been a notable gap in studies analyzing the quantity and distribution of health workers in Indonesia's district level. Previous research often missed the nuances of district-level challenges, focusing instead on broader, national-level assessments.

#### What This Study Adds

- First National Study on Health Workforce: This is the first study of health workforce quantity and distribution at Indonesia's National level. so this paper serves as a basic reference for future research
- Quantitative Analysis of Distribution Equity: Utilizing the Gini Index, the study quantifies the level of inequality in the distribution of healthcare workers, offering a clear metric to guide policymakers in assessing and addressing regional disparities.
- Identification of Regional Variations: The study highlights significant regional variations, with some provinces showing an inequitable distribution of health workers, demonstrating a critical need for increased healthcare personnel and better distribution strategies.
- **Dual-dimensional assessment:** The study introduces a quadrant comparative approach that simultaneously evaluates the quantity of healthcare workers and the equity of their distribution across Indonesian provinces. This dual-dimensional analysis is a significant methodological advancement, providing a more holistic understanding of healthcare workforce allocation.

#### How This Study Might Affect Research, Practice, and/or Policy

This study has the potential to become the basis of policy-making related to the distribution of health workers and provide constructive feedback and strategical insights that could be utilized to decrease the gap between health workers and their maldistribution.

# 50 INTRODUCTION

The landscape of global health is continually evolving, marked by the dynamic interplay between 51 52 healthcare needs and the availability of healthcare professionals. At the heart of this landscape is the 53 pivotal role of health workers, whose quantity and distribution are essential for effective healthcare 54 delivery<sup>1</sup>. The World Health Organization (WHO) since 2005 and aligned with Sustainable 55 Development Goals (SDGs) indicators underscore this significance, particularly after adjusting the need of universal health coverage <sup>2,3</sup>. Human resources for health, serving as the backbone of health 56 57 systems, face the persistent challenge of achieving equitable distribution. With its unique 58 geographical and demographic profile, Indonesia exemplifies this struggle, presenting a complex 59 case for the distribution and availability of health workers.

Indonesia, the world's largest archipelago and fourth most populous nation, with its approximately 17,000 islands, embodies these challenges in distributing its health workforce<sup>4</sup>. The country's vast and varied geography, encompassing remote areas, borders, and small islands, complicates the equitable distribution of healthcare services<sup>5</sup>. Additionally, the decentralized migration of health workers towards urban centers exacerbates the issue, leading to a disparity in healthcare availability between urban and rural or remote regions<sup>6</sup>.

The number of health workers is crucial in determining the effectiveness of healthcare delivery<sup>7</sup>. 66 67 However, the issue extends beyond mere numbers; the distribution of these workers plays a critical 68 role in ensuring that all regions, irrespective of their geographic or demographic characteristics, have access to adequate healthcare services. This challenge is further compounded by the issue of 69 healthcare workforce production in Indonesia, which is not accompanied by equitable distribution<sup>8–</sup> 70 71 <sup>10</sup>. The imbalance in the distribution of health workers, particularly in a geographically diverse 72 country like Indonesia, presents a multifaceted challenge that needs comprehensive analysis and targeted interventions<sup>11</sup>. 73

74 The aim of this study is to provide an in-depth analysis of the current state of health workers in 75 Indonesia, with a particular focus on both their quantity and distribution at the district level. 76 Previous reports and research have often overlooked broader regional or national-level assessments<sup>12</sup>. This lack of detailed data has been a significant gap in understanding the true extent 77 78 of the challenge faced by the Indonesian health system. By delving into the district-level dynamics 79 of health worker distribution and quantity, this study seeks to bridge this knowledge gap and offer 80 insights that could inform policy and practice, ultimately contributing to reducing inequities in 81 healthcare access across Indonesia.

82

# 83 **METHODS**

We gather information from various sources, particularly formal government agencies and ministries that collect data on health resources, both published and unpublished. All the analysis needed will be done in the R using stats and the dineq package.

#### 87 Definition of Administrative Unit

88 Indonesia's administrative area is divided into 34 provinces and 514 districts. Districts are

- 89 categorized as cities (urban areas) or regencies(rural areas), with 416 districts and 98 cities reported
- 90 in 2022. Health worker data is also aggregated into five major islands: Sumatra, Java, Kalimantan,
- 91 Bali-Nusa, Sulawesi, and Maluku-Papua. This approach considers that health service coverage can
- 92 extend beyond district and province administrative boundaries.

#### 93 Definition and Source of Health Worker Data

- 94 Health workers are general practitioners, specialists, dentists, nurses, and midwives. Health workers 95 must have gone through higher education and professional training<sup>13,14</sup>. This study consolidates all 96 24 categories of specialists into a single group. General practitioners (GPs) practice both in primary 97 and referral care, and between private and public. while specialists primarily operate within 98 hospitals and private practices. Midwives are predominantly employed in primary care.
- 99 Number of health workers at the provincial-level data is derived from the "Annual Indonesian
- 100 Health Profile Book" from 1990 to 2021. District-level data were gathered from the "Minister of
- 101 Health's Health Resources Database (SISDMK)". We extracted SISDMK data on 31 December
- 102 2022. Data on doctors, dentists, and specialists are supplemented with Data on Members of the
- 103 Indonesian Doctors Association and Doctors' Registration Data by the Indonesian Medical Council.
- 104 The Health worker ratio will represent the number of health workers per 1,000 population within a
- 105 geographic unit. Population estimate data, used as the health worker ratio's denominator, were
- 106 retrieved from the Ministry of Internal Affairs' GIS data, with the most recent estimates dated 30
- 107 June 2022.

## **108 Benchmark of Health Worker Density**

The existing density of health workers will be compared with goals set by Indonesia's Directorate of Health Workforce Planning's Target Document, utilizing Data Envelopment Analysis (DEA) and Stochastic Frontier Analysis (SFA) for target ratio calculations (see Table 1). We will also compared to the World Health Organization (WHO) proposed threshold of health-worker ratio (including doctors, nurses, and midwives) of 4.45 per 1,000 people to meet 80% coverage of the Sustainable Development Goals (SDGs)<sup>15</sup>

Table 1. Recommendation of Health Worker density											
Health Worker	National Target	Others Recommendation	Source								
GP	1:1000	1:1000	Kumar <i>et.al.</i> , 2018 <sup>16</sup>								
Specialist	0.2:1000	1:1000	WHO								
Dentist	0.2:1000	0.13:1000	Pandya, et.al., 2020 <sup>17</sup>								
Nurse	2.4:1000	3.3:1000	WHO								
Midwifery	2:1000	11.4:1000 (female)	The White Ribbon Alliance. 2017 <sup>18</sup>								
All Health Worker	5.8:1000	4.45: 1000	WHO: Global strategy on human resources for health <sup>19</sup>								

#### 115 **Distribution Analysis**

To analyze distribution, we will directly compare ratios between administrative, urban, and rural areas and use the Gini Index, as previously recommended in WHO Measuring Health Worker inequalities<sup>20 21–23</sup>. The Gini coefficient is a statistical metric obtained from the Lorenz curve, which measures the level of inequality in a distribution. It is expressed as a value between 0 (indicating perfect equality) and 1 (indicating perfect inequality). We performed inter-provincial and within-province Gini Index, which has been prevalently used in prior research <sup>24–27</sup>. To further measure inequities between district, the inter-district Gini index will be used in this research<sup>28,29</sup>

#### 123 Quadrant Analysis Framework

124 We made a quadrant analysis to evaluate the two facets of quantity and distribution, as depicted in

- 125 Figure 1. This method involves plotting provinces on a two-dimensional plane defined by two axes:
- 126 the Gini Index and the Health Worker (HW) Ratio. The quadrant framework categorizes provinces
- 127 into four distinct groups based on their position relative to the median values of the Gini Index and
- 128 HW Ratio. The resulting quadrants are labeled as follows:
- Scarce and Inequitable Province: High Gini Index and Low HW Ratio, indicating a province
   with a shortage of health workers and a highly uneven distribution.
- 131 2. Inequitable Province: High Gini Index and High HW Ratio, suggesting a province with a
  132 sufficient number of health workers but an unequal distribution among the population.
- 3. Scarce Province: Low Gini Index and Low HW Ratio, pointing to a province with an overall
   scarcity of health workers but a relatively even distribution among the population.
- 135 4. **Ideal Province**: Low Gini Index and High HW Ratio, representing a province with both a high
- number of health workers and an equitable distribution.



Figure 1. A Quadrant Analysis of Gini Index vs. Health Worker Ratio

This framework is intended to identify and visualize the disparities in health worker distribution across provinces. In line with the nature of Indonesia, decentralization pushes the provincial government to achieve equal growth within the province. This approach will inform policymakers at the provincial and national levels by highlighting areas of need and guiding targeted interventions.

143

137

## 144 **RESULTS**

#### 145 **Quantity of Health Workers**

Indonesia's health workforce comprises 1,046,640, averaging 3.84 per 1,000 people nationally but
varying widely across districts (0.31 to 19.08). Nurses are the largest group (531,214; 1.95 per
1,000), followed by midwives (300,756; 1.10 per 1,000), general practitioners (145,602; 0.38 per
1,000), specialists (41,893; 0.15 per 1,000), and dentists (27,177; 0.1 per 1,000). The nurse-todoctor ratio is about 2.17 nationally.

151 In examining the distribution of health workers between urban and rural areas in Indonesia, clear 152 disparities are evident across different medical professions. General Practitioners (GPs) in rural 153 areas have a median ratio of 0.27 and an average of 0.31, contrasting sharply with urban areas, 154 where the median ratio is substantially higher at 1.10 and the average is 1.34. The disparity is even 155 more pronounced for dentists; in rural regions, they have a median ratio of 0.05 and an average of 156 0.06, whereas in urban areas, these figures significantly increase to a median of 0.20 and an average 157 of 0.27. Specialists present a similar pattern of distribution. In rural areas, the median ratio for 158 specialists is 0.06 with an average of 0.08, which is markedly lower than in urban areas, where the 159 median ratio rises to 0.35 and the average to 0.42. 160 Rural areas have median ratios of 0.87 for midwives and 1.26 for nurses, which are substantially

161 lower compared to urban areas, where the median ratios are 0.86 for midwives and a significantly

162 higher 3.18 for nurses. Overall, this data underscores a consistent trend of higher health worker

- 164 For most healthcare worker categories, the distribution is right-skewed, meaning that most rural and
- urban areas have lower ratios of healthcare workers, with a few areas having much higher ratios.
- 166 The kurtosis values suggest that these distributions are leptokurtic, with a higher peak and heavier
- tails than a normal distribution, particularly for GPs and dentists in rural areas and for urban nurses.
- 168

	Min	25%	Median	Mean	75%	Max	Skewness	Kurtosis
NATIONAL								
Population	25,015	425,613	1,056,965	1,283,218.61	1,865,647	5,385,219	2.73	10.41
Physician Ratio	0.02	0.22	0.31	0.53	0.63	5.42	4.28	26.7
Specialist Ratio	0	0.05	0.08	0.15	0.16	1.86	4.37	28.97
Dentist Ratio	0	0.03	0.05	0.10	0.10	1.52	6.42	63.73
Midwife Ratio	0.07	0.67	0.86	1.10	1.36	5.16	1.04	1.30
Nurse Ratio	0.22	1.07	1.44	1.95	2.54	12.03	1.63	4.53
Healthworker Ratio	0.31	2.21	3.09	3.84	4.97	19.08	1.45	3.91
RURAL (REGENCIES	) (n = 426)							
Population	25,015	417,509	1,024,842	1,231,418.51	1,679,267	5,385,219	2.98	13.47
Physician Ratio	0.02	0.21	0.27	0.31	0.35	2.27	3.6	22.57
Specialist Ratio	0	0.05	0.06	0.08	0.10	0.57	2.86	15.22
Dentist Ratio	0	0.03	0.05	0.05	0.06	0.52	3.98	28.01
Midwife Ratio	0.07	0.68	0.87	1.13	1.45	5.16	0.99	1.16
Nurse Ratio	0.22	0.98	1.26	1.50	1.8	7.16	1.18	1.79
Healthworker Ratio	0.31	2.04	2.6	3.08	3.68	12.48	0.9	0.94
URBAN (CITIES) (n=8	8)							
Population	34,096	531,570	1,266,194	1,472,195.8	2,470,972	3,274,716	1.94	2.97
Physician Ratio	0.21	0.82	1.10	1.34	1.58	5.42	2.61	8.41
Specialist Ratio	0.02	0.25	0.35	0.42	0.51	1.86	2.52	9.04
Dentist Ratio	0.02	0.13	0.20	0.27	0.35	1.52	3.98	20.64
Midwife Ratio	0.51	0.61	0.86	1.00	1.1	3.75	1.14	1.17
Nurse Ratio	1.55	2.54	3.18	3.61	4.16	12.03	1.46	3.01
Healthworker Ratio	3.1	4.97	5.94	6.64	7.62	19.08	1.53	2.88

Table 2. Descriptive Statistics of Health workers density in Indonesia

## 169 Island and Provincial Levels Distribution

Table 3 shows the distribution of health worker ratios across Indonesian islands and provinces. The data reveals several disparities, with Java having the lowest health worker ratio (3.06 per 1,000 population) and Sulawesi the highest (5.04 per 1,000 population). Each island presents a distinct health worker composition; notably, Bali Nusa Tenggara and Java have the highest GP ratios at 0.45 and 0.38, respectively. Java leads in specialist and dentist ratios (0.17 and 0.1) but has the lowest ratios for nurses and midwives (1.69 and 0.72), suggesting an imbalance in healthcare worker distribution.

- 177 Provincially, Aceh has the highest health worker ratio (7.68), followed by DKI Jakarta (6.23). The
- 178 lowest ratios are in West Java (2.44), Banten (2.69), and Central Java (2.83). Bali's GP ratio (0.98)
- 179 is close to the WHO-recommended physician ratio, and DKI Jakarta has the highest specialist ratio
- 180 (0.63). These figures indicate uneven healthcare resource allocation, with certain areas having a
- 181 high concentration of healthcare professionals and others facing severe shortages.



Figure 2. Regional Health Worker Distribution by City and District: A Density Plot Comparison. This set of density plots compares the distribution of health worker ratios across different Indonesian regions, distinguishing between city and district types. Peaks represent the most common ratios within each region, with red lines indicating city densities and blue lines for districts. The plots reveal variations in distribution patterns, with some regions showing distinct differences between city and district densities, potentially indicating disparities in health worker allocation.

The density plot at figure 2 presents a comparative analysis of health worker (HW) ratios across different islands and regions in Indonesia, categorized by city and district types. In Bali and Nusa, the HW ratio distribution for cities and districts demonstrates a bimodal pattern, indicating two distinct groupings of HW ratios within each category. Java's distribution exhibits a stark peak for cities at a lower HW ratio, suggesting a uniformity in lower ratios among cities, in contrast to districts that show a broader, more varied distribution. Kalimantan's data suggests a single, similar peak for both cities and districts, with cities having a

marginally higher peak, implying a slight concentration of HW ratios within city areas. Meanwhile,
Maluku & Papua reveal a significant contrast between cities and districts; cities have a flatter, more
varied HW ratio distribution, whereas districts show a distinct, narrow peak at higher ratios.

198 Sulawesi's density plot indicates a single peak for both city and district types, with the city peak

being more pronounced, pointing towards a more consistent HW ratio among Sulawesi cities.

200 Sumatera's analysis reveals a higher peak for districts compared to cities, suggesting a trend towards

201 higher HW ratios within district areas. Overall, the graph indicates significant regional variability in

202 HW ratios, with certain trends such as the bimodal distribution in Bali & Nusa and the contrasting

203 distributions in Maluku & Papua standing out.

REGIONAL	POP	GP	Specialist	Dentist	Nurse	Midwife	All HW	
GROUP		n Ratio	n Ratio	n Ratio	n Ratio	n Ratio	n	Ratio
NATIONAL	272,235,314	<b>145602</b> 0.53	<b>41891</b> 0.15	<b>27177</b> 0.10	<b>531214</b> 1.95	<b>300756</b> 1.10	1046640	3.84
CITY	58,567,886	78557 1.34	24772 0.42	15608 0.27	211326 3.61	58721 1.00	388984	6.64
DISTRICT	213,667,428	67045 0.31	17119 0.08	11569 0.05	319888 1.50	242035 1.13	657656	3.08
BALI NUSA	15,028,974	<b>9139</b> 0.61	<b>2326</b> 0.15	<b>1259</b> 0.08	<b>36937</b> 2.46	<b>22769</b> 1.52	72430	4.82
Bali	4,287,193	5754 1.34	1541 0.36	839 0.20	12261 2.86	5835 1.36	26230	6.12
West Nusa Tenggara	5,359,579	1657 0.31	413 0.08	212 0.04	10779 2.01	6393 1.19	19454	3.63
East Nusa Tenggara	5,382,202	1728 0.32	372 0.07	208 0.04	13897 2.58	10541 1.96	26746	4.97
JAVA	153,344,799	<b>84221</b> 0.55	<b>25848</b> 0.17	<b>17476</b> 0.11	259769 1.69	<b>110580</b> 0.72	497894	3.25
Banten	12,142,696	6220 0.51	1745 0.14	1136 0.09	16764 1.38	8719 0.72	34584	2.85
D.I. Yogyakarta	3,677,522	4343 1.18	997 0.27	975 0.27	10715 2.91	2890 0.79	19920	5.42
D.K.I. Jakarta	11,249,585	19493 1.73	7071 0.63	5002 0.44	38819 3.45	7582 0.67	77967	6.93
West Java	48,623,371	19863 0.41	5803 0.12	3357 0.07	65009 1.34	30922 0.64	124954	2.57
Central Java	37,323,519	15156 0.41	4528 0.12	2224 0.06	59749 1.60	28892 0.77	110549	2.96
East Java	40,328,106	19146 0.47	5704 0.14	4782 0.12	68713 1.70	31575 0.78	129920	3.22
KALIMANTAN	16,779,243	<b>8089</b> 0.48	<b>2240</b> 0.13	<b>1442</b> 0.09	<b>41324</b> 2.46	<b>21984</b> 1.31	75079	4.47
West Kalimantan	5,482,046	1632 0.30	404 0.07	236 0.04	10876 1.98	6160 1.12	19308	3.52
South Kalimantan	4,069,123	2087 0.51	668 0.16	504 0.12	9283 2.28	5292 1.30	17834	4.38
Central Kalimantan	2,672,790	1112 0.42	294 0.11	156 0.06	7637 2.86	4435 1.66	13634	5.10
East Kalimantan	3,847,792	2790 0.73	743 0.19	468 0.12	10943 2.84	4882 1.27	19826	5.15
North Kalimantan	707,492	468 0.66	131 0.19	78 0.11	2585 3.65	1215 1.72	4477	6.33
MALUKU & PAPUA	8,634,347	<b>3412</b> 0.40	717 0.08	<b>341</b> 0.04	<b>23000</b> 2.66	<b>11254</b> 1.30	38724	4.48
Maluku	1,886,735	920 0.49	171 0.09	90 0.05	6330 3.36	2575 1.36	10086	5.35
North Maluku	1,317,333	571 0.43	117 0.09	63 0.05	3365 2.55	2740 2.08	6856	5.20
Papua	4,272,108	1282 0.30	303 0.07	120 0.03	8961 2.10	3798 0.89	14464	3.39
West Papua	1,158,171	639 0.55	126 0.11	68 0.06	4344 3.75	2141 1.85	7318	6.32
SULAWESI	19,989,579	<b>10662</b> 0.53	<b>2981</b> 0.15	<b>1833</b> 0.09	<b>54455</b> 2.72	<b>33924</b> 1.70	103855	5.20
Gorontalo	1,203,921	531 0.44	148 0.12	72 0.06	3017 2.51	1815 1.51	5583	4.64
West Sulawesi	1,441,712	360 0.25	100 0.07	102 0.07	3382 2.35	3401 2.36	7345	5.09
South Sulawesi	8,980,741	4994 0.56	1611 0.18	1051 0.12	22365 2.49	13716 1.53	43737	4.87
Central Sulawesi	3,074,958	1239 0.40	315 0.10	174 0.06	8789 2.86	6195 2.01	16712	5.43
South-East Sulawesi	2,625,494	1017 0.39	250 0.10	238 0.09	7876 3.00	6195 2.36	15576	5.93
North Sulawesi	2,662,753	2521 0.95	557 0.21	196 0.07	9026 3.39	2602 0.98	14902	5.60
SUMATERA	58,458,372	<b>30079</b> 0.51	<b>7779</b> 0.13	<b>4826</b> 0.08	115729 1.98	100245 1.71	258658	4.42
Aceh	5,356,965	3469 0.65	889 0.17	489 0.09	18178 3.39	19017 3.55	42042	7.85
Bengkulu	2,041,265	903 0.44	211 0.10	154 0.08	5077 2.49	4294 2.10	10639	5.21
Jambi	3,600,625	1684 0.47	405 0.11	276 0.08	8252 2.29	6763 1.88	17380	4.83
Bangka Belitung Isl.	1,464,663	826 0.56	221 0.15	135 0.09	3860 2.64	1696 1.16	6738	4.60
Riau Island	2,072,922	1623 0.78	405 0.20	277 0.13	5019 2.42	2567 1.24	9891	4.77
Lampung	8,866,673	2803 0.32	693 0.08	282 0.03	12216 1.38	10458 1.18	26452	2.98
Riau	6,032,801	4015 0.67	917 0.15	626 0.10	11405 1.89	8736 1.45	25699	4.26
West Sumatera	5,481,484	3398 0.62	999 0.18	552 0.10	11291 2.06	7800 1.42	24040	4.39
South Sumatera	8,489,000	3625 0.43	967 0.11	431 0.05	16032 1.89	14512 1.71	35567	4.19
North Sumatera	15,051.94	7733 0.51	2072 0.14	1604 0.11	24399 1.62	24402 1.62	60210	4.00

# 205 District-Level Distribution

Figure 3 shows a distinct change in the mapping of health workers between the provincial-level aggregate ratio(1a) and district-level aggregate ratio(1b), especially in Java Island. Health workers are concentrated in cities (small areas with a high ratio of health workers), and lack of health workers in more rural areas in Java. Despite having the largest number of healthcare workers, by comparing the number to their population, the ratio in most districts in Java remained below Indonesia and WHO-recommended density (Figure 4).



213 Figure 3. Geographic Distribution of Health Worker Ratios in Indonesia: A Comparative View at Provincial and 214 District Levels. This map pair illustrates the health worker (HW) ratio across Indonesia's provinces (1a) and districts 215 216 (1b). Darker shades of blue indicate higher HW ratios, with the scale ranging from 0 to 4. The provincial-level map

shows broader trends, while the district-level map provides a more granified perspective, revealing significant

217 variations within provinces.



218 219 220 Figure 4. Population vs. Health Worker Ratio Across Indonesian Regions: Scaling and WHO Standards Compliance. The scatter plot displays the relationship between the logarithm of the population size and the health worker ratio for various Indonesian regions. Each point represents a region, with the WHO-recommended ratio as a 221 reference line. The plot highlights disparities in health worker distribution relative to population, with some regions 222 significantly above or below the WHO benchmark.

The Figure 4 scatter plot displays a wide variance in health worker-to-population ratios. The graph indicates that most regions have health worker ratios below the density target except for a few districts that are also the province's capital. Several regions, particularly in Papua, such as Yahukimo, are identified as lower outliers with a deficiency in health worker density.

Districts with a larger population tend to have fewer health workers. A logarithmic trend line fitted to the data demonstrates a decremental trend, suggesting that regions with larger populations tend to have a lower health-worker ratio. The Logarithmic trend line equation quantifies the relationship between the population size and the health worker ratio, Health Worker Ratio = -1.2749\*ln(Population) + 19.004. The coefficient of determination, R-squared (0.215958). Moreover, the distribution of health worker ratios exhibits right-skewness. This skewness indicates that while most regions are understaffed, a few are exceptionally well-staffed.

# 234 Inequity in Distribution

The Gini Index has demonstrated variable trends over the past three decades, as depicted in Figure 4

- and Table 4. For General Practitioners (GPs), the provincial Gini Index increased from 0.35 in 1992
- to a peak of 0.49 in 2003 before declining to 0.39 in 2012 and rising again to 0.42 in 2022. This
- 238 fluctuation suggests changes in the equality of GP distribution over time. Dentists and nurses also
- experienced an increase in provincial inequality, with their Gini Indexes rising to 0.49 and 0.40,
- respectively, in 2022. Midwives, however, showed an improvement in equality from 1992 to 2022,
- with the Gini Index decreasing from 0.38 to 0.35.

Table 4. Provincial and District-Level Inequality in Health Worker Distribution: Gini Index from 1992 to 2022. This table presents the Gini Index scores for different health worker categories, measuring the inequality in their distribution across provinces from 1992 to 2022 and at the district level in 2022. Data Unavailability is marked as '...'.

Hoolth worker	Betwe	een Provii	Between District Gini					
Health worker	1992	1997	2003	2008	2012	2017	2022	Index 2022
GP	0.35	0.45	0.49	0.42	0.39	0.39	0.42	0.46
Specialist	0.57	0.57			0.42	0.45	0.44	0.53
Dentist	0.38	0.43	0.41	0.42	0.40	0.43	0.49	0.52
Nurse	0.34	0.36	0.37		0.36	0.38	0.40	0.33
Midwife	0.38	0.38	0.39	0.39	0.41	0.36	0.35	0.31

242

Disparities at the district level, recorded for 2022, reveal a more pronounced inequality for health workers than at the provincial level. GPs saw their district-level Gini Index escalate to 0.46, dentists to 0.52, and specialists to the highest at 0.53. In contrast, Nurses and midwives exhibited a reduced inequality at the district level, with a Gini Index of 0.33 and 0.31, respectively, signifying a more between-province inequality rather than relatively equal distribution across districts.





Supported by the Lorenz curves, it suggests that the bottom percentage of districts have significantly fewer health workers than would be expected in an equal distribution, especially for GPs, specialists, and dentists. The middle of the distribution for these categories also shows a considerable gap from the line of equality, indicating that the Gini index's sensitivity to the middle distribution reflects the reality of the distribution of health workers. The top district(s) seem to have a disproportionately high number of health workers, especially specialists and dentists, as indicated by the curves' steep rise as they approach 100%.

#### 260 Scarcity and Inquity

Figure 6 offers a visual evaluation of health worker distribution across Indonesian regions. The yaxis is the Gini Index to assess equity in distribution against the x-axis as health worker ratios for various categories, including general practitioners, specialists, dentists, midwives, and nurses.

The health worker scatterplot illustrates a widespread distribution across the provinces, with a noticeable lean towards lower ratios, which suggests a general scarcity of health workers in many areas. North Sumatra, in particular, registers a high Gini Index coupled with a low health-worker ratio, underscoring an urgent need for redistribution and an increase in the health workforce.

The scatterplots for physicians, specialists, and dentists display a similar pattern, with most provinces failing to meet minimum health worker targets. With exceptions such as Jakarta, Yogyakarta, and Bali that exceed these targets. With its five districts and a substantial physician population, Jakarta is uniquely situated in the bottom-right quadrant, highlighting an oversupply of physicians compared to other regions clustered to the left. Specialists are predominantly located in the upper-left quadrant, indicating a skewed distribution that favors urban centers or provincial

274 capitals and suggests a concentration of specialists in fewer, more developed areas. with

- 275 Nurses exhibit the most equitable distribution, with only Papua showing significant disparities.
- 276 However, the range between the highest and lowest ratios remains considerable, from 1.2 to 3.6 per
- 277 1,000. The Midwives ratio plot identifies Aceh as the only province exceeding the equity threshold.
- 278 Conversely, Jakarta, typically lower-right in other health worker categories, deviates in the case of
- 279 Midwive, showing a surprisingly low ratio and positioning it towards the left quadrant.



Figure 6. Comparative Analysis of Health Worker Distribution Across Regions: Gini Index vs. Various Health Worker Ratios. (Each scatter plot represents a different category of health workers, with the Gini index on the vertical axis and the respective health worker ratio on the horizontal axis. Dot represents a province, with blue shades indicating regions with a higher Gini of health. The dashed line for Gini referenced 0.35 Gini Ratio, the target for equality. Dashed line for Ratio referenced to national Target for each health worker)

# 286 **DISCUSSION**

## 287 Quantity of Health Workers

288 Indonesia's archipelagic geography, with its varied terrain and areas isolated by mountains and 289 oceans, adds to the distribution challenges. Among its 80,000 villages, over 20,000 face 290 geographical difficulties that obstruct access to healthcare. As of 2022, Indonesia's health-worker 291 ratio is 3.84. The threshold set by WHO to achieve 80% Universal Health Coverage is 4.45 requiring an additional 166,000 health workers to surpass the 4.45 threshold.<sup>30</sup> Indonesia's lack of 292 293 active GPs and Specialists requires attention (0.64 physicians per 1,000 population). Low compared 294 to other LMIC Nations such as India (0.73 per 1000), China (2.4 doctors per 1,000), and other ASEAN Countries (Singapore (2.29), the Philippines (0.6), and Malaysia (1.54).)<sup>31</sup>. Indonesia must 295

add up to 100,000 physicians to meet the threshold.<sup>17–20</sup>

## 298 **Decomposition of Inequity**

299 Contrary to the general perception of Java being saturated with health workers. By numbers, Java 300 has the most healthcare workers, but Java has the lesser density of total health workers, mainly 301 caused by the low number of nurses and midwives. Java is the only island with a midwives ratio 302 lower than 1 and a nurse ratio lower than 2 per 1000 population. However, Java is also the island 303 with the highest ratio of specialists (0.17), while Maluku and Papua had a ratio lower than half of 304 Java's (0.8).

Most islands have nearly reached the WHO threshold, with only Java (3.52) and Sumatra (4.42) still

below it. However, the high population on these islands significantly affects the national average of

health workers. None of the islands have yet met the Indonesian Government's threshold of 5.8
health workers per 1000 population, with Sulawesi coming closest at 5.2.

The key issue lies in the composition of health workers. Java faces a shortage of nurses and midwives despite having a relatively high number of doctors, though still below the threshold. In

311 contrast, islands outside of Java have a relative abundance of nurses and midwives but a scarcity of

312 doctors.<sup>32</sup> This imbalance in the distribution of health workers can be addressed through aggressive

313 policy measures and enhanced nurse education, as demonstrated by China's experience. In China,

around 2003, the ratio of nurses to physicians was quite low. However, recent data indicates a

significant improvement, with the ratio rising from 1 nurse per 1000 population in 2003 to 4.1 nurses per 1000 population in 2018.<sup>33</sup>

317 There is a profound disparity between urban and rural areas. Up to 50% of the total GP was in the

City, despite the city population being only 20% of the total Population. That's also happening in

319 India, where Out of all qualified workers, 77.4% were located in urban areas, even though the urban

population is only 31% of the country's total population. Cities exhibit higher mean ratios across all

321 healthcare workforce categories. Like other low- and lower-middle-income nations, the city tends to

have a more significant health worker density.<sup>23,34-42</sup> While still skewed, the midwife and nurse

323 ratios are less severe.<sup>43</sup> This finding is consistent with other research that found an uneven

distribution of the total health workforce between rural and urban areas.<sup>44–46</sup>

#### 325 Inequity Indicator

The overall trend indicates that while some health worker categories, like midwives, have achieved an equal distribution, others, particularly specialists and dentists, face higher levels of inequality, especially considering the district level. This discrepancy is highlighted by the Gini Index differences between the provincial and district levels, with an apparent increase in inequality at the finer scale of districts for most health worker categories.

331 There has been some progress in equalizing the distribution of medical specialists across provinces.

332 The between-provinces Gini Index For Specialists remarkably dropped from 0.57 in 1993 to 0.44 in

333 2022. However, the Inter-district Gini Index is very high at 0.53 in 2022. This implies that

specialists are still saturated in the preferable district, huge cities, and the province's capital.<sup>34,47,48</sup>
Measuring only the between-province Gini Index will mask the disparities within provinces and
interdistricts.

Comparative studies from western China, a less affluent region compared to the eastern and southern parts of the country, exhibit similar inequality trends. The Physician Gini Index here stands at 0.46. The World Health Organization's last published data from 2010 shows relatively more minor interdistrict disparities in China, with a Gini Index of 0.362 for doctors and 0.471 for nurses. In India, these figures are 0.436 for doctors and 0.527 for nurses<sup>49,50</sup>. The situation in Indonesia contrasts with both China and India, as it exhibits a higher Gini Index for general physicians and specialists than nurses.

## 344 Two Faces of Health Worker Problem in Indonesia: Inequity and Scarcity

The quadrant analysis provides a multidimensional view of healthcare workforce distribution across regions. It highlights the areas of need and the complexity of issues like scarcity and inequality that often overlap, making it difficult to discern the underlying causes and appropriate interventions.

348 The upper-right quadrant has a high ratio of health workers, but resources are not evenly spread 349 across the province. Bali serves as an illustrative case within this analysis. For instance, urban 350 centers like Denpasar City might be well-staffed, while more remote areas suffer from a dearth of 351 healthcare professionals. Addressing this imbalance requires policies incentivizing healthcare 352 workers to relocate to underserved areas or increasing the training and recruitment of health 353 workers specifically for these regions.

The upper-left quadrant represents a more critical scenario, as seen with East Java and Papua regions. The inequality in their distribution worsens the simultaneously general scarcity of health workers. Such provinces struggle with redistribution because the overall number of health workers is too low to meet even the minimum needs. To mitigate this, a two-pronged approach is essential: boosting the healthcare workforce numbers through training and recruitment and ensuring retention in underserved areas, perhaps through enhanced benefits, better working conditions, or other incentives.

From these findings, we can assume that Indonesia faces a challenge in evenly distributing healthcare professionals, with rural areas experiencing more pronounced shortages. To date, health equity hasn't been prioritized as a main criterion for evaluating the success of health programs.<sup>51</sup>

Policy interventions to incentivize healthcare professionals to work in underserved areas, scaling up healthcare infrastructure<sup>35</sup> and education programs to meet the demand in regions with lower healthcare workforce ratios.<sup>32,52,53</sup>

368 WHO emphasizes the necessity of district-level reporting of health worker data. While provincial-369 level data offers a general overview, it often fails to represent the situation at the district level. 370 High-level aggregated data can obscure crucial specifics about the distribution and availability of health workers at the subregional level<sup>51</sup>. The strain from a lack of healthcare workers was apparent 371 372 during the COVID-19 pandemic. Healthcare services almost collapsed during the pandemic peak. 373 Any decrease in healthcare workers led to longer working hours for the remaining staff. This created a domino effect of increasing health workers' loads<sup>54</sup>. In maternity care, World Bank 374 375 statistics reveal that maternal mortality rates in Indonesia have persistently remained relatively high 376 from 2017 to 2021 compared to other Southeast Asian countries. A shortage of healthcare 377 professionals is one critical factor influencing the heightened maternal mortality rate in Indonesia<sup>55,56</sup>. NCD demands a workforce for every step, especially preventive care <sup>57</sup>. 378 379 Hypertension in Indonesia increased from 25.8% of the population in 2013 to 34.1% in 2018. 380 Notably, many hypertension cases remain asymptomatic, leading to a high prevalence of 381 undiagnosed.

#### 382 Limitation

383 During data collection, we found notable discrepancies between the numbers and distribution of 384 General Physicians and Specialists as reported by the medical association, Medical Council, and 385 Ministry of Health. These inconsistencies likely stem from the council's reliance on correspondence 386 addresses rather than the actual practice locations of medical doctors. Ministry of Health data is 387 much more focused on public health center workers. We acknowledge limitations in the IDI data, 388 primarily due to a five-year delay from a patient's death to the data's release.

# 389 CONCLUSIONS

390 The imbalance in the distribution of health workers in Indonesia is a burden for this country

- 391 Government policy factors, security, economic conditions, individual characteristics, income access
- to children's education, knowledge development, and areas with a better socioeconomic scope will
- be more attractive to health workers, especially specialist doctors.

The health workforce improvement agenda must focus on both increasing the number of health worker and distribute it equally to all people. Central government play a huge role in improving health worker production and distribution to area needed, and Local governments play a vital role in healthcare worker retention. Tailor strategies to local contexts, improve working conditions, offer professional development, and create supportive environments. Engage communities to foster belonging, boosting job satisfaction and retention.

## 400 **DECLARATION**

- 401 **Ethics approval an consent to participate**
- 402 Not applicable

- 403
- 404 Availability of data and materials
- 405 Data is available by requesting an application through email to the corresponding author.
- 406 **Competiting interests**
- 407 All authors and contributors to the study stated that they had no conflict of interest.
- 408 Funding
- 409 The study was conducted with Research Grant from General Directorate of Higher Education,
- 410 Ministry of Education and Culture of Indonesia (DIPA-023.17.1.690523/2023).
- 411

# 412 Authors' contributions

- 413 Conceptualization: Farizal Rizky Muharram, Achmad Chusnu Romdhoni
- 414 Formal analysis: Farizal Rizky Muharram, Sudhir Anand, Hanif Ardiansyah Sulistya, Fikri
- 415 Febrian Firmansyah, Julian Benedict Swannjo
- 416 Methodology: Farizal Rizky Muharram, Hanif Ardiansyah Sulistya, Fikri Febrian Firmansyah,
- 417 Julian Benedict Swannjo, Muhammad Masrur Rizal
- 418 Project administration: Alifina Izza, Muhammad Atoillah Isfandiari, Ninuk Dwi Ariningtyas
- 419 Resources: Farizal Rizky Muharram, Hanif Ardiansyah Sulistya, Fikri Febrian Firmansyah, Julian
- 420 Benedict Swannjo, Muhammad Masrur Rizal, Alifina Izza
- 421 Supervision: Achmad Chusnu Romdhoni, Muhammad Atoillah Isfandiari, Ninuk Dwi Ariningtyas
- 422 Writing original draft: Farizal Rizky Muharram, Hanif Ardiansyah Sulistya, Julian Benedict
- 423 Swannjo, Muhammad Masrur Rizal, Alifina Izza
- 424 Writing review & editing: Farizal Rizky Muharram, Sudhir Anand, Hanif Ardiansyah Sulistya, ,
- 425 Julian Benedict Swannjo, Alifina Izza, Muhammad Atoillah Isfandiari, Ninuk Dwi Ariningtyas

426

427 Acknowledgments

429	REFEREN	CES
430	1.	Jaeger FN, Bechir M, Harouna M, Moto DD, Utzinger J. Challenges and opportunities
431		for healthcare workers in a rural district of Chad. BMC Health Serv Res [Internet].
432		2018 Jan 8 [cited 2023 Nov 22];18(1):1–11. Available from:
433		https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-017-2799-6
434	2.	Jacobs B, Ir P, Bigdeli M, Annear PL, Van Damme W. Addressing access barriers to
435		health services: an analytical framework for selecting appropriate interventions in low-
436		income Asian countries. Health Policy Plan [Internet]. 2012 Jul 1 [cited 2023 Nov
437		22];27(4):288–300. Available from: https://dx.doi.org/10.1093/heapol/czr038
438	3.	Singh Thakur J, Nangia R, Singh S. Progress and challenges in achieving
439		noncommunicable diseases targets for the sustainable development goals. FASEB
440		Bioadv [Internet]. 2021 Aug 1 [cited 2023 Nov 22];3(8):563–8. Available from:
441		https://onlinelibrary.wiley.com/doi/full/10.1096/fba.2020-00117
442	4.	Leosari Id Y, Uelmen JA, Marc R, Id C. Spatial evaluation of healthcare accessibility
443		across archipelagic communities of Maluku Province, Indonesia. PLOS Global Public
444		Health [Internet]. 2023 Mar 9 [cited 2023 Nov 22];3(3):e0001600. Available from:
445		https://journals.plos.org/globalpublichealth/article?id=10.1371/journal.pgph.0001600
446	5.	Dwi Laksono A, Dwi Wulandari R, Rohmah N, Rukmini R, Tumaji T. Basic Health
447		Survey. BMJ Open [Internet]. 2023 [cited 2023 Nov 22];13:64532. Available from:
448		http://bmjopen.bmj.com/
449	б.	Meliala A, Hort K, Trisnantoro L. Addressing the unequal geographic distribution of
450		specialist doctors in Indonesia: The role of the private sector and effectiveness of
451		current regulations. Soc Sci Med. 2013 Apr 1;82:30–4.
452	7.	Crisp N, Gawanas B, Sharp I. Training the health workforce: scaling up, saving lives.
453		The Lancet [Internet]. 2008 Feb 23 [cited 2023 Nov 22];371(9613):689–91. Available
454		from: http://www.thelancet.com/article/S0140673608603098/fulltext
455	8.	Efendi F, Haryanto J, Indarwati R, Kuswanto H, Ulfiana E, Has EMM, et al. Going
456		Global: Insights of Indonesian Policymakers on International Migration of Nurses. J
457		Multidiscip Healthc [Internet]. 2021 Nov 26 [cited 2023 Nov 22];14:3285–93.
458		Available from: https://www.dovepress.com/going-global-insights-of-indonesian-
459		policymakers-on-international-migr-peer-reviewed-fulltext-article-JMDH
460	9.	Meilianti S, Smith F, Kristianto F, Himawan R, Ernawati DK, Naya R, et al. A
461		national analysis of the pharmacy workforce in Indonesia. Hum Resour Health
462		[Internet]. 2022 Dec 1 [cited 2023 Nov 22];20(1):1–12. Available from:
463		https://human-resources-health.biomedcentral.com/articles/10.1186/s12960-022-
464		00767-4
465	10.	Anderson I, Meliala A, Marzoeki P, Pambudi E. The Production, Distribution, and
466		Performance of Physicians, Nurses, and Midwives in Indonesia: An Update. Health,
467		Nutrition and Population (HNP) Discussion Paper Series [Internet]. 2014 [cited 2023
468		Nov 22]; Available from: https://ideas.repec.org/p/wbk/hnpdps/91324.html
469	11.	Homepage J, Sutrisno S. Shortages of Medical Doctors in Indonesia, Is It True? Asian
470		Journal of Health Research [Internet]. 2023 Aug 11 [cited 2023 Nov 22];2(2):1–2.
471	10	Available from: https://a-jhr.com/a-jhr/article/view/121
472	12.	Laksono AD, Ridlo IA, Ernawaty. DISTRIBUTION ANALYSIS OF DOCTORS IN
4/3		INDONESIA. Indonesian Journal of Health Administration [Internet]. 2020 Mar 28
4/4		[cited 2023 Nov 22];8(1):29–39. Available from: https://e-
4/5	12	Journal.unair.ac.id/JAKI/article/view/14351
4/0	15.	indonesia. Law no. 30 of 2014 on Health Care Personnel. 2014. 2014 [cited 2023 Nov
4// 170	1 /	22]; Available from: www.indolaw.org
470 470	14.	Available from: https://paraturan.bak.go.id/Datails/252022/viv. no. 17 tabun 2022
419 100	15	Avanable Holli: https://peraturali.opk.go.lu/Details/258028/uu-no-1/-tanun-2023
40U /121	15.	Direktorat refericanaan Tenaga Kesenatan KKI. Dokumen Target Kasio Tenaga Kesehatan 2022:
401 187	16	Kumar P. Dal P. India achieves WUO recommanded destor population ratio: A call
402 183	10.	for paradigm shift in public health discourse! I Family Med Drim Care 2018:7(5):841
-0J		for paradigin sint in public nearly useourse; J ranny Meu Filli Cale, 2010,7(J).041.

484 485	17.	Pandya VS, Sampath N, Yadav R, Mahuli A V, Kapadiya JD. Dental Manpower in India: changing trends upto 2020. Journal of Xidian University. 2021 Jul 3;15(7).
486	18.	The White Ribbon Alliance. A COUNT OF BEDSIDE MIDWIVES IN MALAWI.
487		2017.
488	19.	World Health Organization. Global strategy on human resources for health: Workforce
489		2030. 2016.
490	20.	Achim Zeileis, Christian Kleiber. ineq: Measuring Inequality, Concentration, and
491		Poverty [Internet]. 2014 [cited 2022 Dec 2]. Available from: https://CRAN.R-
492		project.org/package=ineq
493	21.	Jin J, Wang J, Ma X, Wang Y, Li R. Equality of Medical Health Resource Allocation
494		in China Based on the Gini Coefficient Method [Internet]. Vol. 44, Iran J Public
495		Health. 2015. Available from: http://ijph.tums.ac.ir
496	22.	Panzera D, Postiglione P. Measuring the Spatial Dimension of Regional Inequality:
497		An Approach Based on the Gini Correlation Measure. Soc Indic Res. 2020 Apr
498		1;148(2):379–94.
499	23.	Hermawan A. Analisis Distribusi Tenaga Kesehatan (Dokter Perawat Dan Bidan) Di
500		Indonesia Pada 2013 Dengan Menggunakan Gini Index. Buletin Penelitian Sistem
501		Kesehatan. 2019;22(3):167–75.
502	24.	Saito E, Gilmour S, Yoneoka D, Gautam GS, Rahman MM, Shrestha PK, et al.
503		Inequality and inequity in healthcare utilization in urban Nepal: A cross-sectional
504		observational study. Health Policy Plan. 2016 Sep 1;31(7):817–24.
505	25.	Anyangwe SCE, Mtonga C. Inequities in the Global Health Workforce: The Greatest
506		Impediment to Health in Sub-Saharan Africa [Internet]. Vol. 4, Int. J. Environ. Res.
507		Public Health. 2007. Available from: www.ijerph.org
508	26.	Boniol M, McCarthy C, Lawani D, Guillot G, McIsaac M, Diallo K. Inequal
509		distribution of nursing personnel: a subnational analysis of the distribution of nurses
510		across 58 countries. Hum Resour Health. 2022 Dec 1;20(1).
511	27.	Boniol M, McCarthy C, Lawani D, Guillot G, McIsaac M, Diallo K. Inequal
512		distribution of nursing personnel: a subnational analysis of the distribution of nurses
513		across 58 countries. Hum Resour Health. 2022 Dec 1;20(1).
514	28.	Global strategy on human resources for health: Workforce 2030.
515	29.	WHO - Guideline on health policy and system support to optimize community health
516		worker programmes.
517	30.	Global strategy on human resources for health: workforce 2030 Reaffirming the
518		continuing importance of the application of the WHO Global Code of Practice on the
519		International Recruitment of Health Personnel (hereinafter "WHO Global Code"); 2
520		[Internet]. 2016. Available from: https://sustainabledevelopment.un.org/?menu=1300
521	31.	Bank W, Penh P, Kanchanachitra C, Lindelow M, Johnston T, Hanvoravongchai P, et
522		al. Health in Southeast Asia 5 Human resources for health in southeast Asia:
523		shortages, distributional challenges, and international trade in health services. Lancet
524		[Internet]. 2011:377:769–81. Available from: www.thelancet.com
525	32.	Afenyadu GY, Adegoke AA, Findley S, Improving Human Resources for Health
526		means Retaining Health-Workers: Application of the WHO-Recommendations for the
527		Retention of Health-Workers in Rural Northern-Nigeria. J Health Care Poor
528		Underserved [Internet], 2017 [cited 2023 Nov 18]:28(3):1066–86. Available from:
529		https://pubmed.ncbi.nlm.nih.gov/28804079/
530	33.	Lu H. Hou L. Zhou W. Shen L. Jin S. Wang M. et al. Original research: Trends.
531	001	composition and distribution of nurse workforce in China: a secondary analysis of
532		national data from 2003 to 2018. BMJ Open [Internet] 2021 Oct 27 [cited 2023 Nov
533		18]:11(10). Available from: /pmc/articles/PMC8552175/
534	34.	Karan A. Negandhi H. Hussain S. Zapata T. Mairembam D. De Graeve H. et al. Size
535	0	composition and distribution of health workforce in India <sup>•</sup> why and where to invest?
536		Hum Resour Health [Internet], 2021 Dec 1 [cited 2023 Nov 18]:19(1):39 Available
537		from: /pmc/articles/PMC7983088/

538	35	Polym PI Shroff 7 Zang OP Kingua S Dijanouassi S Kouontahou C at al
530	55.	Addressing health word-force distribution concernes a discrete choice experiment to
539		Addressing health workforce distribution concerns: a discrete choice experiment to
540		develop fural relention strategies in Cameroon. Int J Health Poncy Manag [Internet].
541	26	2015 [cited 2023 Nov 18];4(3):169. Available from: /pmc/articles/PMC435/984/
542	36.	Hsu YHE, Lin W, Tien JJ, Tzeng LY. Measuring inequality in physician distributions
543		using spatially adjusted Gini coefficients. International Journal for Quality in Health
544		Care. 2016;28(6):657–64.
545	37.	Ismail M. Regional disparities in the distribution of Sudan's health resources. Eastern
546		Mediterranean Health Journal. 2020 Sep 1;26(9):1105–14.
547	38.	Spatial distribution and associated factors of community based health insurance
548		coverage in Ethiopia: further analysis of Ethiopian demography and health survey,
549		2019   BMC Public Health   Full Text [Internet]. [cited 2022 Nov 23]. Available from:
550		https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-022-13950-y
551	39.	Zhang T, Xu Y, Ren J, Sun L, Liu C. Inequality in the distribution of health resources
552		and health services in China: Hospitals versus primary care institutions. Int J Equity
553		Health. 2017 Mar 3;16(1).
554	40.	Mirmoeini SM, Shooshtari SSM, Battineni G, Amenta F, Tayebati SK. Policies and
555		challenges on the distribution of specialists and subspecialists in rural areas of Iran.
556		Vol. 55. Medicina (Lithuania), MDPI AG: 2019.
557	41.	Anyangwe SCE, Mtonga C, Inequities in the Global Health Workforce: The Greatest
558		Impediment to Health in Sub-Saharan Africa [Internet] Vol 4 Int J Environ Res
559		Public Health 2007 Available from: www.jjernh.org
560	42	Saito F. Gilmour S. Yoneoka D. Gautam GS. Rahman MM. Shrestha PK. et al.
561	72.	Inequality and inequity in healthcare utilization in urban Nepal: A cross-sectional
562		observational study, Health Dolicy Dian, 2016 Sep 1:21(7):817–24
562	12	Laksono AD Didlo IA. Emousty E. Distribution Analysis of Destors in Indonesia
505	45.	Laksono AD, Kidio IA, Elliawaty E. Distribution Analysis of Doctors in Indonesia.
504		Indonesian Journal of Health Administration [Internet]. $2020$ Mar $26;6(1):29-39$ .
505	4.4	Available from: https://e-journal.unair.ac.id/JAKI/article/view/14551
566	44.	Heywood PF, Haranap NP. Human Resources for Health Human resources for health
567		at the district level in Indonesia: the smoke and mirrors of decentralization. 2009 [cited
568	4.5	2023 Nov 18]; Available from: http://www.numan-resources-nealth.com/content///1/6
569	45.	Samli A, Hadju V, Soma AS. Spatial analysis of health facilities in Mamuju City,
570		West Sulawesi. Enferm Clin [Internet]. 2020 Oct 1 [cited 2023 Nov 18];30:71–5.
571		Available from: https://www.elsevier.es/es-revista-enfermeria-clinica-35-articulo-
572		spatial-analysis-health-facilities-in-S1130862120303570
573	46.	Sumber Daya Manusia K, Sumber Daya Manusia Kesehatan pada Fasilitas Kesehatan
574		Tingkat Pertama dalam Era Jaminan K, Yuyun Yuniar Pusat Penelitian dan
575		Pengembangan Sumber Daya dan Pelayanan Kesehatan dan. Ketersediaan Sumber
576		Daya Manusia Kesehatan pada Fasilitas Kesehatan Tingkat Pertama dalam Era
577		Jaminan Kesehatan Nasional di Delapan Kabupaten-Kota di Indonesia. Media
578		Penelitian dan Pengembangan Kesehatan [Internet]. 2016 [cited 2023 Nov
579		18];26(4):201–10. Available from: https://www.neliti.com/publications/179261/
580	47.	Rao KD, Shahrawat R, Bhatnagar A. Composition and distribution of the health
581		workforce in India: estimates based on data from the National Sample Survey. WHO
582		South East Asia J Public Health [Internet]. 2016 Sep 1 [cited 2023 Nov 18];5(2):133–
583		40. Available from: https://pubmed.ncbi.nlm.nih.gov/28607241/
584	48.	Meliala A, Hort K, Trisnantoro L. Addressing the unequal geographic distribution of
585		specialist doctors in Indonesia: The role of the private sector and effectiveness of
586		current regulations. Soc Sci Med. 2013 Apr:82:30–4.
587	49.	Anand S. Methods for measuring health workforce inequalities: methods and
588	• • •	application to China and India. Human Resources for Health Observer [Internet] 2010
589		[cited 2023 Nov 21]:5:1–32. Available from:
590		http://www.who.int/hrh/resources/observer/en/

591	50.	Organization WH, Anand S, Fan V. The health workforce in India [Internet]. Geneva:
592		World Health Organization; 2016. (Human Resources for Health Observer Series, 16).
593		Available from: https://iris.who.int/handle/10665/250369
594	51.	Utomo B, Sucahya PK, Utami FR. Priorities and realities: Addressing the rich-poor
595		gaps in health status and service access in Indonesia. Vol. 10, International Journal for
596		Equity in Health. 2011.
597	52.	World Health Organization. Global strategy on human resources for health: Workforce
598		2030. 2016.
599	53.	Pagaiya N, Kongkam L, Sriratana S. Rural retention of doctors graduating from the
600		rural medical education project to increase rural doctors in Thailand: A cohort study.
601		Hum Resour Health [Internet]. 2015 Mar 1 [cited 2023 Nov 18];13(1):1-8. Available
602		from: https://human-resources-health.biomedcentral.com/articles/10.1186/s12960-015-
603		0001-y
604	54.	Mahendradhata Y, Andayani NLPE, Hasri ET, Arifi MD, Siahaan RGM, Solikha DA,
605		et al. The Capacity of the Indonesian Healthcare System to Respond to COVID-19.
606		Front Public Health. 2021 Jul 7;9.
607	55.	Mahmood MA, Hendarto H, Laksana MAC, Damayanti HE, Suhargono MH,
608		Pranadyan R, et al. Health system and quality of care factors contributing to maternal
609		deaths in East Java, Indonesia. PLoS One. 2021 Feb 1;16(2 February).
610	56.	Sejati EN, Rosa EM, Pramesona BA. Trends and Determinants of the Maternal
611		Mortality Ratio Based on Healthcare Resources. Unnes Journal of Public Health. 2023
612		Mar 1;12(1):1–11.
613	57.	Suparmi, Kusumawardani N, Nambiar D, Trihono, Hosseinpoor AR. Subnational
614		regional inequality in the public health development index in Indonesia. Glob Health
615		Action. 2018 Dec 3;11(sup1).
616		-





Figure 1. A Quadrant Analysis of Gini Index vs. Health Worker Ratio





Figure 2. Regional Health Worker Distribution by City and District: A Density Plot Comparison. This set of density plots compares the distribution of health worker ratios across different Indonesian regions, distinguishing between city and district types. Peaks represent the most common ratios within each region, with red lines indicating city densities and blue lines for districts. The plots reveal variations in distribution patterns, with some regions showing distinct differences between city and district densities, potentially indicating disparities in health worker allocation.

9



11 12

Figure 3. Geographic Distribution of Health Worker Ratios in Indonesia: A Comparative View at Provincial 13 and District Levels. This map pair illustrates the health worker (HW) ratio across Indonesia's provinces (1a) and 14 districts (1b). Darker shades of blue indicate higher HW ratios, with the scale ranging from 0 to 4. The provincial-15 level map shows broader trends, while the district-level map provides a more granified perspective, revealing 16 significant variations within provinces.

17 18

.



- 19 Figure 4. Population vs. Health Worker Ratio Across Indonesian Regions: Scaling and WHO Standards
- 20 Compliance. The scatter plot displays the relationship between the logarithm of the population size and the health
- 21 worker ratio for various Indonesian regions. Each point represents a region, with the WHO-recommended ratio as a
- 22 reference line. The plot highlights disparities in health worker distribution relative to population, with some regions 23
- significantly above or below the WHO benchmark.
- 24

medRxiv preprint doi: https://doi.org/10.1101/2024.03.31.24305126; this version posted April 1, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.





31

25



33 Figure 6. Comparative Analysis of Health Worker Distribution Across Regions: Gini Index vs.

34 **Various Health Worker Ratios.** (Each scatter plot represents a different category of health workers, with 35 the Gini index on the vertical axis and the respective health worker ratio on the horizontal axis. Dot

represents a province, with blue shades indicating regions with a higher Gini of health. The dashed line for

37 Gini referenced 0.35 Gini Ratio, the target for equality. Dashed line for Ratio referenced to national Target

- 38 for each health worker)
- 39

1

Table 1. Recommended Health Worker to Population Ratios: A Comparative Table of Global and National Standards									
Health Worker	National Target Ratio	Others Recommended Ratio	Source						
GP	1:1000	1:1000	Kumar <i>et.al.</i> , 2018 <sup>16</sup>						
Specialist	0.2:1000	1:1000	WHO						
Dentist	0.2:1000	0.13:1000	Pandya, et.al., 2020 <sup>17</sup>						
Nurse	2.4:1000	3.3:1000	WHO						
Midwifery	2:1000	11.4:1000 (female)	The White Ribbon Alliance. 2017 <sup>18</sup>						
All Health Worker	5.8:1000	4.45: 1000	WHO: Global strategy on human resources for health <sup>19</sup>						

2

4

#### Table 2. Descriptive Statistics of Healthcare Economic Indicators and Workforce Ratios in Indonesia

	Min	25%	Median	Mean	75%	Max	Skewness	Kurtosis
NATIONAL	1		l			I	1	U
Population	25015	425613	1056965	1283218.61	1865647	5385219	2.73	10.41
Fiscal Capacity	243	850	1297	2256.74	2510	11391	5.22	32.59
GNI Per Capita PPP	4190	9840	10913	11694.83	12522	24221	0.74	1.78
Physician Ratio	0.02	0.22	0.31	0.53	0.63	5.42	4.28	26.7
Specialist Ratio	0	0.05	0.08	0.15	0.16	1.86	4.37	28.97
Dentist Ratio	0	0.03	0.05	0.10	0.10	1.52	6.42	63.73
Midwife Ratio	0.07	0.67	0.86	1.10	1.36	5.16	1.04	1.30
Nurse Ratio	0.22	1.07	1.44	1.95	2.54	12.03	1.63	4.53
Healthworker Ratio	0.31	2.21	3.09	3.84	4.97	19.08	1.45	3.91
REGENCIES (n = 426)		t			t			
Population	25015	417509	1024842	1231418.51	1679267	5385219	2.98	13.47
Fiscal Capacity	243	803	1187	1575.09	1857	6209	3.29	15.03
GNI Per Capita PPP	4190	9446	10588	10530.34	11609	18927	0.06	0.97
Physician Ratio	0.02	0.21	0.27	0.31	0.35	2.27	3.6	22.57
Specialist Ratio	0	0.05	0.06	0.08	0.10	0.57	2.86	15.22
Dentist Ratio	0	0.03	0.05	0.05	0.06	0.52	3.98	28.01
Midwife Ratio	0.07	0.68	0.87	1.13	1.45	5.16	0.99	1.16
Nurse Ratio	0.22	0.98	1.26	1.50	1.8	7.16	1.18	1.79
Healthworker Ratio	0.31	2.04	2.6	3.08	3.68	12.48	0.9	0.94
CITIES (n=88)								·
Population	34096	531570	1266194	1472195.8	2470972	3274716	1.94	2.97
Fiscal Capacity	374	1188	3518	4743.54	6448	11391	2.62	6.02
GNI Per Capita PPP	7519	13803	15503	15943.14	18199	24221	0.63	1.32
Physician Ratio	0.21	0.82	1.10	1.34	1.58	5.42	2.61	8.41
Specialist Ratio	0.02	0.25	0.35	0.42	0.51	1.86	2.52	9.04
Dentist Ratio	0.02	0.13	0.20	0.27	0.35	1.52	3.98	20.64
Midwife Ratio	0.51	0.61	0.86	1.00	1.1	3.75	1.14	1.17
Nurse Ratio	1.55	2.54	3.18	3.61	4.16	12.03	1.46	3.01
Healthworker Ratio	3.1	4.97	5.94	6.64	7.62	19.08	1.53	2.88

5

 Table 3. Healthcare Workforce Distribution by Region: Population-to-Health Worker Ratios in Indonesia.

 The ratios are calculated per 1,000 population.

REGIONAL	POP	GP	Specialist	Dentist	Nurse	Midwife	All HW	
GROUP	-	n Ratio	n Ratio	n Ratio	n Ratio	n Ratio	n	Ratio
NATIONAL	272,235,314	<b>145602</b> 0.53	<b>41891</b> 0.15	<b>27177</b> 0.10	<b>531214</b> 1.95	<b>300756</b> 1.10	1046640	3.84
СІТҮ	58,567,886	78557 1.34	24772 0.42	15608 0.27	211326 3.61	58721 1.00	388984	6.64
DISTRICT	213,667,428	67045 0.31	17119 0.08	11569 0.05	319888 1.50	242035 1.13	657656	3.08
BALI NUSA	15,028,974	<b>9139</b> 0.61	<b>2326</b> 0.15	<b>1259</b> 0.08	<b>36937</b> 2.46	<b>22769</b> 1.52	72430	4.82
Bali	4,287,193	5754 1.34	1541 0.36	839 0.20	12261 2.86	5835 1.36	26230	6.12
West Nusa Tenggara	5,359,579	1657 0.31	413 0.08	212 0.04	10779 2.01	6393 1.19	19454	3.63
East Nusa Tenggara	5,382,202	1728 0.32	372 0.07	208 0.04	13897 2.58	10541 1.96	26746	4.97
JAVA	153,344,799	<b>84221</b> 0.55	<b>25848</b> 0.17	<b>17476</b> 0.11	<b>259769</b> 1.69	<b>110580</b> 0.72	497894	3.25
Banten	12,142,696	6220 0.51	1745 0.14	1136 0.09	16764 1.38	8719 0.72	34584	2.85
D.I. Yogyakarta	3,677,522	4343 1.18	997 0.27	975 0.27	10715 2.91	2890 0.79	19920	5.42
D.K.I. Jakarta	11,249,585	19493 1.73	7071 0.63	5002 0.44	38819 3.45	7582 0.67	77967	6.93
West Java	48,623,371	19863 0.41	5803 0.12	3357 0.07	65009 1.34	30922 0.64	124954	2.57
Central Java	37,323,519	15156 0.41	4528 0.12	2224 0.06	59749 1.60	28892 0.77	110549	2.96
East Java	40,328,106	19146 0.47	5704 0.14	4782 0.12	68713 1.70	31575 0.78	129920	3.22
KALIMANTAN	16,779,243	<b>8089</b> 0.48	<b>2240</b> 0.13	1442 0.09	<b>41324</b> 2.46	<b>21984</b> 1.31	75079	4.47
West Kalimantan	5,482,046	1632 0.30	404 0.07	236 0.04	10876 1.98	6160 1.12	19308	3.52
South Kalimantan	4,069,123	2087 0.51	668 0.16	504 0.12	9283 2.28	5292 1.30	17834	4.38
Central Kalimantan	2,672,790	1112 0.42	294 0.11	156 0.06	7637 2.86	4435 1.66	13634	5.10
East Kalimantan	3,847,792	2790 0.73	743 0.19	468 0.12	10943 2.84	4882 1.27	19826	5.15
North Kalimantan	707,492	468 0.66	131 0.19	78 0.11	2585 3.65	1215 1.72	4477	6.33
MALUKU & PAPUA	8,634,347	<b>3412</b> 0.40	717 0.08	<b>341</b> 0.04	<b>23000</b> 2.66	<b>11254</b> 1.30	38724	4.48
Maluku	1,886,735	920 0.49	171 0.09	90 0.05	6330 3.36	2575 1.36	10086	5.35
North Maluku	1,317,333	571 0.43	117 0.09	63 0.05	3365 2.55	2740 2.08	6856	5.20
Papua	4,272,108	1282 0.30	303 0.07	120 0.03	8961 2.10	3798 0.89	14464	3.39
West Papua	1,158,171	639 0.55	126 0.11	68 0.06	4344 3.75	2141 1.85	7318	6.32
SULAWESI	19,989,579	<b>10662</b> 0.53	<b>2981</b> 0.15	<b>1833</b> 0.09	<b>54455</b> 2.72	<b>33924</b> 1.70	103855	5.20
Gorontalo	1,203,921	531 0.44	148 0.12	72 0.06	3017 2.51	1815 1.51	5583	4.64
West Sulawesi	1,441,712	360 0.25	100 0.07	102 0.07	3382 2.35	3401 2.36	7345	5.09
South Sulawesi	8,980,741	4994 0.56	1611 0.18	1051 0.12	22365 2.49	13716 1.53	43737	4.87
Central Sulawesi	3,074,958	1239 0.40	315 0.10	174 0.06	8789 2.86	6195 2.01	16712	5.43
South-East Sulawesi	2,625,494	1017 0.39	250 0.10	238 0.09	7876 3.00	6195 2.36	15576	5.93
North Sulawesi	2,662,753	2521 0.95	557 0.21	196 0.07	9026 3.39	2602 0.98	14902	5.60
SUMATERA	58,458,372	<b>30079</b> 0.51	<b>7779</b> 0.13	<b>4826</b> 0.08	115729 1.98	100245 1.71	258658	4.42
Aceh	5,356,965	3469 0.65	889 0.17	489 0.09	18178 3.39	19017 3.55	42042	7.85
Bengkulu	2,041,265	903 0.44	211 0.10	154 0.08	5077 2.49	4294 2.10	10639	5.21
Jambi	3,600,625	1684 0.47	405 0.11	276 0.08	8252 2.29	6763 1.88	17380	4.83
Bangka Belitung Isl.	1,464,663	826 0.56	221 0.15	135 0.09	3860 2.64	1696 1.16	6738	4.60
Riau Island	2,072,922	1623 0.78	405 0.20	277 0.13	5019 2.42	2567 1.24	9891	4.77
Lampung	8,866,673	2803 0.32	693 0.08	282 0.03	12216 1.38	10458 1.18	26452	2.98
Riau	6,032,801	4015 0.67	917 0.15	626 0.10	11405 1.89	8736 1.45	25699	4.26
West Sumatera	5,481,484	3398 0.62	999 0.18	552 0.10	11291 2.06	7800 1.42	24040	4.39
South Sumatera	8,489,000	3625 0.43	967 0.11	431 0.05	16032 1.89	14512 1.71	35567	4.19
North Sumatera	15 051 94	7733 0 51	2072 0 14	1604 0.11	24399 1.62	24402 1 62	60210	4 00

9

Table 4. Provincial and District-Level Inequality in Health Worker Distribution: Gini Index from 1992 to 2022. This table presents the Gini Index scores for different health worker categories, measuring the inequality in their distribution across provinces from 1992 to 2022 and at the district level in 2022. Data Unavailability is marked as '...'.

Health worker	Betwe	een Provir	Between District Gini					
Health worker	1992	1997	2003	2008	2012	2017	2022	Index 2022
GP	0.35	0.45	0.49	0.42	0.39	0.39	0.42	0.46
Specialist	0.57	0.57			0.42	0.45	0.44	0.53
Dentist	0.38	0.43	0.41	0.42	0.40	0.43	0.49	0.52
Nurse	0.34	0.36	0.37		0.36	0.38	0.40	0.33
Midwife	0.38	0.38	0.39	0.39	0.41	0.36	0.35	0.31