

Video Observed Therapy (VOT) for People with Tuberculosis (TB): A Scoping Review

Authors:

An Du Think¹, Eleanor Morgan², Kassia Pereira³, *Celso Khosa^{1,3,4}, *Tom Wingfield^{1,2,5}

Affiliations:

1. Departments of Clinical Science and International Public Health, Liverpool School of Tropical Medicine, Liverpool, United Kingdom
2. Royal Liverpool University Hospital, Liverpool, United Kingdom
3. Instituto Nacional de Saúde, Marracuene, Mozambique
4. Faculdade de Medicina, Eduardo Mondlane University, Maputo, Mozambique
5. World Health Organization Collaborating Centre on Tuberculosis and Social Medicine, Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

*Contributed equally

Contributions

An Du Think (ADT): Conceptualization, Methodology, Investigation, Data Curation, Formal analysis, Writing - Original Draft, Review & Editing, Visualization

Eleanor Morgan (EM): Conceptualization, Methodology, Investigation, Data Curation, Formal analysis, Writing – supported Original Draft, Review & Editing, Visualization

Kassia Pereira (KP): Conceptualization, Methodology, Investigation, Data Curation, Formal analysis, Writing - Review & Editing, Visualization

Celso Khosa (CK): Conceptualization, Methodology, Investigation, Formal analysis, Writing - Review & Editing, Visualization

Tom Wingfield (TW): Conceptualization, Methodology, Investigation, Formal analysis, Writing - Review & Editing, Visualization

All authors read and approved the final manuscript.

Corresponding Author:

Dr Tom Wingfield

Email: tom.wingfield@lstm.ac.uk

X: @drtomwingfield

ORCID: <https://orcid.org/0000-0001-8433-6887>

Room W-01-34 Wolfson Building, Liverpool School of Tropical Medicine, Liverpool, L3 5QA, United Kingdom

Key words:

video observed therapy; digital health; tuberculosis; feasibility; acceptability; effectiveness; cost-effectiveness; scoping review

Abstract

Background: Tuberculosis (TB) remains a significant cause of morbidity and mortality globally, disproportionately affecting low- and middle-income countries (LMIC). Accessing Directly Observed Therapy (DOT) is associated with out-of-pocket costs and stigma. Video-observed therapy (VOT) is an alternative to DOT but evidence from LMIC with high TB burden is limited. To make recommendations for future VOT-related research and inform the design of a pilot implementation of VOT for people with multi-drug resistant TB (MDR-TB) in Mozambique, we did a scoping review of the evidence on VOT for people with TB.

Methodology: We systematically searched five scientific databases and key grey literature repositories to identify eligible abstracts. Abstracts were reviewed and full-text records were identified and evaluated. Data from full-text records were extracted into four implementation theme categories: Feasibility, Acceptability, Cost, and Effectiveness (FACE). Content analysis was used to describe implementation successes and challenges, comparing VOT versus DOT where possible. The Crowe Critical Appraisal Tool (CCAT) was used to evaluate the quality of studies.

Results: In total, 66 records were identified: 47 primary research studies, 13 reviews, and six grey literature documents. All studies were graded as moderate to high quality and reported against at least one FACE category. Studies from urban settings (n=34) and HIC (n=33) predominated. Where measured and described, VOT implementation was reported to be feasible (43/43, 100%) and acceptable (43/44, 97%). Of the 18/20 (90%) studies describing cost data, VOT was reported to offer savings to the health system compared to DOT. Patient costs were under-reported. Of the 21/23 (91%) studies describing effectiveness to improve adherence to TB treatment, VOT was reported to be non-inferior to DOT.

Conclusion: In HIC settings, VOT was reported as feasible, acceptable, and similarly effective alternative to DOT. Further evidence on VOT is needed from LMIC with high TB burden.

Author Summary

Directly Observed Therapy (DOT) remains widely used to monitor treatment adherence of people living with tuberculosis (TB). Since COVID-19, there has been a massive increase in digital health technologies, mobile phone ownership, and internet connectivity, including in high TB burden low- and middle-income countries (LMIC), highlighting the potential for Video Observed Therapy (VOT) as a suitable alternative to DOT. We did a scoping review of studies reporting the implementation and evaluation of VOT for people with TB using a novel thematic framework consisting of feasibility, acceptability, cost and effectiveness (FACE). We identified 20 VOT applications (45% freely available) used by studies. Most VOT studies were conducted in urban settings of low TB burden, High Income Countries (HIC), and did not include cost-effectiveness analyses from a patient perspective or participants from key, underserved groups such as pregnant women or people with TB/HIV. The available evidence suggested that, in mainly HIC with low TB burden, VOT was a feasible, acceptable and effective alternative to DOT which could provide cost savings to the healthcare provider. There is an urgent need for robust evidence of effectiveness, cost-effectiveness, and equity of VOT in LMICs with high TB burden and among key, underserved populations.

Introduction

Tuberculosis (TB) remains a major global public health challenge. In 2023, TB regained, from COVID-19, the position of the leading cause of death from a single infectious disease worldwide^{1,2}. Most high TB burden countries are lower- and lower-middle-income countries (LMIC), which are disproportionately affected by TB-related morbidity and mortality, especially where HIV prevalence is high^{2,3}. Without treatment, 10-year mortality from TB is approximately 70%³. Although shorter TB treatment regimens are effective in treating TB, most countries are still using antimicrobial regimens lasting up to 18 months to treat multi-drug resistant TB (MDR-TB), a leading cause of death from antimicrobial resistance (AMR). The high pill-burden and long treatment duration has clinical, psychological, and socio-economic consequences, challenging adherence. Treatment success rates for MDR-TB are estimated to be approximately 60%³. Suboptimal adherence increases the likelihood of adverse TB treatment outcomes including treatment failure, AMR, and onward transmission.

Providing support to people with TB to facilitate adherence to treatment can minimize adverse outcomes⁴. Directly Observed Therapy (DOT) has been recommended by the World Health Organization (WHO) as a method to improve TB treatment adherence. While some studies have suggested that DOT can increase adherence when compared to self-administered therapy (SAT), a Cochrane systematic review and meta-analysis found no significant differences in treatment success between DOT and SAT⁴. Facility-based DOT involves observation of medication doses by healthcare workers (HCW) at healthcare centres. This requires daily travel by people with TB and DOT may not be accessible for communities in remote locations, where travel to healthcare centres is challenging or unaffordable⁵. In many cases this results in high levels of SAT, especially during weekends when

healthcare centres may be closed. Moreover, DOT may be associated with stigma and a loss of autonomy among people with TB⁶.

Video Observed therapy (VOT) is a potential alternative to DOT. In VOT, videos are live-streamed or uploaded for review by HCWs, either *synchronously* (at the time of medication ingestion) or *asynchronously* (the recording is reviewed after medication ingestion). VOT has been highlighted as an essential digital health tool to achieve the 2015 WHO End TB Strategy aims to eliminate TB⁷. In some studies^{8,9}, VOT has shown promise in increasing treatment adherence. This may relate to VOT being more convenient than DOT because it allows people with TB to take their treatment from home, work, or another location, using their mobile device to record ingestion. Moreover, compared to an in-person DOT appointment at a healthcare centre, VOT can reduce travel time, health system costs, patient costs, and patients' perceived stigma^{7,10,11}.

Despite its potential, there is limited up-to-date evidence on the implementation and effectiveness of VOT in LMIC and amongst vulnerable groups, including people with MDR-TB or comorbidities. This evidence gap is particularly pertinent given the expansion of digital health technologies since the COVID-19 pandemic and the increase in mobile phone ownership, usage, and internet coverage in LMICs¹². This scoping review aimed to gather, synthesise and critically appraise the latest evidence on VOT for people with TB globally in order to inform the design of a pilot implementation of VOT for people with MDR-TB in Mozambique (the "SAFEST-1 MDR-TB" study), and make recommendations for future research.

Methods

Study Design

A scoping review was considered the most suitable method to comprehensively gather evidence from both the scientific and grey literature, provide details on VOT platforms used, and encompass heterogeneous study types and resources^{13,14}. The scoping review was conducted in accordance with the Joanna Briggs Institute (JBI) guidance¹⁴.

The JBI and Cochrane recommended PICO (Population, Intervention, Comparison, and Outcome) was used to help inform the search strategy and inclusion criteria¹⁵. Population was adults, pregnant women, adolescents, and children with TB. Intervention was VOT for TB treatment adherence monitoring. Comparison was DOT, DOTS, or SAT vs VOT. Outcomes included feasibility, acceptability, costs, effectiveness, and reported success and challenges of implementation.

Inclusion Criteria

The inclusion criteria included: i) Accessible peer-reviewed scientific journal articles of any study design or published reports with the primary or co-primary aim of describing and/or evaluating and/or reviewing Video Observed Therapy (VOT) for people (adults and children) with tuberculosis (TB) of any type and in any country, and assessed to be of sufficient quality to facilitate suitable data extraction for meaningful content analysis and interpretation (detailed further below); ii) Reports, working papers, technical guidance, and other non-scientific literature appearing in the grey literature of key databases (detailed further below); iii) Publication between 1st of January 2000 and 1st of January 2023 – a timescale chosen because of the paucity of digital adherence technologies, video-phone applications, and mobile phone coverage prior to the year 2000; and iv) Publications in English, Portuguese, French, and

Spanish languages, in which the research team collectively had suitable reading comprehension skills.

Search Strategy

We systematically searched Medline, Embase, PubMed, Google Scholar and Web of Science databases for relevant articles, supplemented by a grey literature search of key databases to identify records published between 1st January 2000 and 1st January 2023. Grey literature was identified from a pre-specified convenience sample of databases selected through the consensus of the research team. These included: the World Health Organisation (WHO), Centers for Disease Control and Prevention (CDC), European Centre for Disease Prevention and Control (ECDC), and WHO International Clinical Trials Registry Platform (ICTRP). Reference snowballing and citation tracking were used to complement and expand the search.

The search for records was conducted between 9th February 2023 and 31st March 2023. Duplicates were identified and removed. Titles and abstracts were reviewed by two team members independently, but not blinded, according to the inclusion criteria. Where there was disagreement, a third opinion was sought from a third reviewer. The full-text of selected records were uploaded to a password-protected online, “live” shared Excel spreadsheet stored on an LSTM cloud-based server. Only deidentified and anonymised secondary data was stored and used and there was no confidential participant information.

The quality of records selected for full-text review was evaluated using the Crowe Critical Appraisal Tool (CCAT)¹⁶. Based on previous research by our study team⁶, a decision was taken to pragmatically categorise CCAT scores (percentage score out of a total of 40 points) as follows: ≥ 30 points, 75%-100%, high quality; 20 to 29 points, 50%-74%, moderate quality; and < 20 points, $< 50\%$, low quality.

Data Analysis

From included full-text records, we extracted aggregate sociodemographic data of participants, geographic and World Bank economic classification data (low, middle, high income) of study sites, data related to the VOT platforms used, and implementation data for content analysis on the challenges and success of VOT with relation to Feasibility, Acceptability, Cost, and Effectiveness (FACE) – categories decided a priori through study team consensus. Data were presented in narrative and tabular form.

Results

Study Selection

The search yielded 429 records: PubMed (n=138), Medline (n=79), Web of Science (n=100), Cochrane (n=7) and Google Scholar (n=105)(Figure 1). Of these, 160 were removed, mainly due to duplication, leaving 269 records to be screened. Following eligibility screening, 60 records were selected for full-text review. The complementary search of the pre-specified convenience sample of grey literature databases yielded 22 registered clinical trials and citation “snowballing” searching by manually reviewing references of the above 60 included studies found another six relevant studies; these two sources combined to identify a further six studies for full-text review (Figure 1).

[The authors suggest inserting Figure 1 here]

Study Characteristics

The 66 records selected consisted of: 47 primary research studies (Table 1), 13 review-type studies including systematic reviews, scoping reviews, and

narrative/non-systematic reviews (Table 2), and six grey literature documents (Table 3).

[The authors suggest inserting Tables 1, 2, and 3 here]

Most studies (n=34) were conducted at study sites in urban settings, 16 were in suburban/peri-urban and only one study³³ assessed VOT in the rural context. Of the 55 studies from which a country and income level could be inferred, 33/55 (60%) of the studies were from high-income countries, 8/55 (15%) were from upper-middle-income and middle-income countries, and 14/55 (25%) were from LMIC. Most high-income studies came from the USA (n=25, Supplementary Figure 1), which has a low TB incidence of 2.5 cases per 100,000⁷⁷.

[The authors suggest inserting VOT world map here]

Intervention Types

Of the 47 primary research studies, the majority (n=29) were single-arm intervention studies of VOT with no comparison group. A minority of studies (n=15) were VOT versus DOT head-to-head comparisons, and the remaining three were VOT compared with other adherence methods such as SMS and medication monitors⁵¹, comparison of two different VOT apps¹⁸, and one study examining VOT in a paediatric population³¹. Four studies (4/66, 8%)^{9,36,55,57} were Randomised Control Trials (RCT) and, of the five systematic reviews, two^{4,71} had an accompanying meta-analysis. One study²⁶ specifically evaluated VOT for supporting people with MDR-TB. No studies were identified that assessed VOT interventions amongst pregnant women or specifically amongst people with TB/HIV co-infection.

A Comprehensive Summary of VOT Applications

In the 66 records identified, a wide variety of applications and technologies were mentioned. Of the 20 applications identified, 12/20 (60%), including the top five most cited or used, were developed in the USA. The only VOT application developed in an LMIC context was NIMCURE from Nigeria, which has the highest TB burden in Africa accounting for 4.6% of the global TB cases⁷⁸. Supplementary Table 1 summarises the applications' characteristics, costs, and which records cited or used these.

[The authors suggest inserting Supplementary Table 1 here]

Critical Appraisal using CCAT

Across the primary studies and review-type studies where the CCAT was applicable, the range of scores was from 21 (52.5%) to 39 (97.5%) out of 40, with a mean score of 32.38 (81%). This meant that all studies were rated as moderate to high quality on the CCAT scoring system. See Supplementary Table 2 for the disaggregation of CCAT scores attained.

Feasibility

A total of 43/66 (65%) records reported on assessments of the feasibility of VOT as a modality to measure TB adherence. Of these, 35 were primary studies, five were review-type studies and three were grey literature documents. All studies reported that VOT was feasible regardless of urbanicity (although most were conducted in urban/peri-urban settings), country income level or population context. However, the most frequently reported challenges to feasibility were lack of or unreliable internet coverage^{5,25,26,28,36,39,42,48,50,56,57,61,75}, low quality of videos or incomplete dose verification^{17,26,28,34,35,40–42,60,70} and other technical issues including battery problems^{18,22,25,28,31,33,49,50,58}.

Studies identified the need to loan a VOT-compatible smartphone to people with TB^{33,34,44,50} and consider the cost of packages of mobile phone data⁵. One study⁵³ found that prior ownership of a smartphone was not associated with the ability to learn how to use VOT. Potential lack of rapport between HCWs and people with TB was described^{5,35,62} and challenges to adherence including not attending scheduled VOT appointments or being lost to follow-up were perceived to be similar to standard DOT^{25,26,28,33}.

One retrospective study in Baltimore (urban), USA⁴⁶ reported usage of VOT during the COVID-19 pandemic when DOT was not possible. It highlighted age as the main barrier to the use of VOT, with older participants finding the technology more challenging. Two studies^{23,52} also reported that VOT uptake, was more common in younger people. Fewer than 10% of participants in Guo *et al.*³⁷ experienced issues with uploading videos for VOT and Garfein *et al.*³² reported that healthcare providers were able to verify most VOT-delivered doses. Holzman *et al.*⁴⁰ showed that VOT was utilised successfully when participants travelled to another country, which the authors interpreted as potentially facilitating increased acceptability of VOT. Ravenscroft *et al.*⁵⁵ highlighted that people with TB using VOT were more likely to report side effects than people with TB using DOT.

VOT was found to be technically feasible in the conflict setting of Syria⁶³ but its utility was limited by intermittent supply of TB medications. Donahue *et al.*³¹ found at a military health facility for families of service members found that VOT was feasible in a paediatric population with parental support. A WHO report¹⁰ called “Digital health for the end TB strategy: An Agenda for action” considered that VOT could become more feasible as coverage of mobile smartphones with internet capability increased.

In its guidance on TB control in Europe, the ECDC scientific panel deemed that VOT was “Likely” feasible to use for vulnerable and hard-to-reach populations⁷⁴.

Acceptability

A total of 44/66 (66%) records, mostly primary research studies, evaluated and reported on the acceptability of VOT. Except for a single study in the Philippines²⁶, in which uptake of VOT was low (36%), all found VOT acceptable to people with TB and healthcare providers. A preference amongst people with TB for VOT over DOT was reported to relate to increased convenience, time-saving, and flexibility^{22–25,32–34,36,37,40–42,48,54}. Amongst healthcare workers, two studies from China and The Philippines, reported a preference for VOT over DOT^{26,37}. In studies which measured satisfaction levels amongst people with TB using VOT versus people with TB using DOT, people using VOT reported higher levels of satisfaction^{27,55,72}.

Both people with TB and healthcare workers or providers raised privacy, security, and confidentiality as potential concerns with VOT across several studies^{5,33,34,36,41,47,48,50,57,58}. Specifically, Garfein *et al.*⁵ identified cultural concerns of videos of females with TB being observed by male care providers in certain countries or contexts. However, this was balanced by studies reporting that both people with TB and HCWs perceived VOT to be more confidential than DOT^{17,22,32,34,41,72}. Rabinovich *et al.*'s concept study⁵³ stated that although clients were receptive to the prospect of a mobile application for VOT purposes, there were concerns surrounding usability of the technology, especially for people with low literacy or who were illiterate.

In Vietnam, Nguyen *et al.*⁵⁰ noted the most common reason (34.2%) for unwillingness to engage in VOT was not wanting to use a smartphone, but this was not mirrored by other studies. In other studies, people with TB reported appreciated

ad hoc communication offered by VOT^{39,41} and nurses valued the professional development gained from IT training for VOT.

Costs

A total of 20/66 (33%) records, including 15 primary studies, examined costs associated with VOT. A further nine studies commented subjectively on the cost-saving attributes of VOT but did not conduct formal costing analyses. Most studies reported that, compared to healthcare facility-based or community-based DOT, VOT was associated with cost savings for people with TB, the health system, or both. There were, however, *two* exceptions: Asay *et al.*¹⁹ evaluated three TB programs offering VOT and found that, in Rhode Island, USA, a high-income context, VOT costs were not different from community-based DOT. This may have related to power because the same study suggested costs to people with TB and the health system were lower with VOT in larger cities (and cohorts) including in New York City and San Francisco. In a systematic review, Wong *et al.*⁷² found that, in the context of tuberculosis infection, SMS-reminder technology was the most cost-effective method for promoting adherence to TB preventive therapy, quoting a cost-analysis simulation study⁵¹, based in Brazil, a middle-income, high TB and TB/HIV country context.

Cost-saving attributes of VOT over DOT were due to reduced travel costs and reduced time spent observing doses in-person, with studies commenting specifically on personnel (staff) costs^{5,9,17,24,33,43,48,61}, hours and costs including transport saved by people with TB^{5,37,48,55}. Chuck *et al.*²⁸ found that staff could observe twice the number of people with TB taking their medications per allotted appointment slot duration when using VOT compared with DOT. Asynchronous VOT was found by Lam *et al.*⁴⁵ to have

lower costs to the health system than synchronous VOT but a mobile phone, software purchase, installation costs, and costs to people were not evaluated.

Effectiveness

A total of 39/66 (59%) records commented on VOT effectiveness, 29 of which were primary research studies, nine were review-type studies, and one was a grey literature article. Versus DOT, VOT was reported to be associated with better adherence across various settings^{8,9,21,27,28,30,33,37,46,52,55,68,71}. This was often interpreted as relating to VOT, especially asynchronous VOT, being provided 7-days per week and being less affected by staffing shortages, strikes, and national holidays. Several studies reported no difference in adherence rates between VOT and DOT^{5,17,22,35,40,47,57,64}. Perry *et al.*⁵² reported more missing doses (average 3.4% vs. 1.3%, respectively; $P < 0.001$) over a median period of 27 weeks but similar adherence rates (96% vs. 90%; $P=0.326$) in VOT vs DOT. Several studies reported high adherence rates with VOT but without comparison to a DOT control group^{18,24,31,32,34,41,48,50,58}.

Another common measure of effectiveness was treatment completion as defined by the WHO⁷⁶, which was reported as higher with VOT than DOT in various studies^{8,27,44}. Comparable treatment completion rates with VOT versus DOT were reported in other studies^{4,26,28,36,47,68,71,76}. No study reported treatment completion rates to be worse amongst people using VOT compared to DOT but, notably, none were specifically designed to demonstrate treatment completion inferiority or superiority. Mortality was infrequently reported but was found to be comparable in three studies^{4,36,76} and, in the study by Casalme *et al.*²⁶ amongst people with MDR-TB, mortality was found to be lower with VOT compared with DOT. In some studies, loss to follow-up was either lower²⁶ or the same³⁶ with VOT versus DOT. No evidence

of effectiveness to reduce other adverse TB treatment outcomes including treatment failure were identified.

Treatment success was higher amongst people with TB using VOT than DOT in one study²⁶, but comparable between VOT and DOT in other studies^{36,52,55,64}. Culture conversion rates for VOT were similar to DOT in three studies^{37,52,71}. TB recurrence (i.e. return of symptoms and signs including radiological signs of clinical TB and/or microbiologically-confirmed TB) was reported in Doltu *et al.*³⁰ and Guo *et al.*³⁷ in the DOT control groups but not VOT groups. No other studies reported TB recurrence in either the VOT or DOT groups.

The key success, challenges, and potential solutions to VOT implementation identified above are summarised and disaggregated by country income in Table 4.

[The authors suggest inserting Table 4 here]

Discussion

The findings from this scoping review illustrate a lack of evidence, especially from randomised controlled trials, on the feasibility, acceptability, cost and effectiveness of VOT for people with TB in LMIC. In predominantly HIC, VOT was shown to contribute to high adherence levels and to have the potential to be more effective and cost-saving than DOT. A wide variety of VOT applications were used, most of which were developed in HICs. International consensus guidelines relating to VOT from WHO, ECDC, and CDC, were largely based on best practice and with low-grade evidence.

Availability of VOT

This scoping review was able to build on the findings of a scoping review from 2016⁷⁹ to update and summarise the current landscape and diversity of VOT applications used and their pricing, availability, and user interface/platform. It was identified that costs associated with VOT applications, such as user licenses, downloads, and recurrent subscription costs, are not transparent or easily available. In addition, some study trials provided smartphones for participants which may not be sustainable or suitable for upscaling within the health systems of many LMIC high TB burden settings beyond trial conditions.

Feasibility

This scoping review found that VOT was uniformly reported to be *feasible* in HIC but more evidence is needed from LMIC. VOT is now an established alternative to DOT in New York City, USA and London, UK^{9,28}. Shared challenges to feasibility were identified that commonly related to technical aspects such as internet access, submissions of poor-quality videos, and initial problems with navigating the VOT applications, but most studies were able to overcome these without significant loss to follow-up reported. Personal access to a smartphone or loan of a smartphone to participants by the study was an accessibility issue raised^{11,23}. For TB programmes to sustain VOT delivery, a pre-intervention baseline analysis of the use of smartphones and network coverage and consideration of underserved groups, including those without access, extremely impoverished, or with literacy and digital literacy issues, is essential to ensure equitable delivery of VOT, echoing recommendations from Peru⁸⁰ and WHO⁷⁵.

Acceptability

People living with TB and healthcare workers were widely receptive to VOT as an acceptable alternative to DOT. The WHO “Handbook for the use of digital technologies to support tuberculosis medication adherence”⁷⁵ suggested the need to evaluate acceptability of VOT in subpopulations such as adolescent girls and women in high TB burden LMIC to ensure equitable impact. The ECDC scientific panel rated VOT as “likely acceptable” for adherence monitoring of underserved populations in Europe⁷⁴. Future VOT implementations must assure participants that their privacy will be maintained and ensure data security in line with local legislative requirements, such as the UK’s Data Protection Act 2018. In addition, differentiating the process and impact outcomes of asynchronous versus synchronous VOT remains a knowledge gap.

To enhance acceptability for HCWs, VOT training should be used as an opportunity for continued professional development of IT skills. From the perspective of people with TB, it must be recognised that VOT is not a blanket solution for all and TB programmes should retain the option for DOT for those who prefer or require face-to-face interactions.

Costs

Of the minority of studies that formally collected and analysed cost data, most reported cost savings of VOT versus DOT for people with TB and the health system. The actual and proportional cost savings may vary between HIC and LMIC settings due to differential personnel, equipment, and logistics costs. For future studies, it was noted that other costs such as incentivising adherence, for example through weekly food vouchers, must also be included for accurate cost calculations. Our findings were similar to a previous scoping review⁶⁷, which also found cost-savings of up to 58% for

the healthcare system amongst 19 studies and in agreement with the ECDC scientific panel contribution⁷⁴ that VOT was “Likely” cost-effective.

For people with TB, the most widely reported cost saving of VOT avoiding travel costs to DOT clinics, was also noted to have potential positive public health implications⁸¹. For HCWs, VOT was reported to improve time-efficiency and, in some cases, reduce their own travel costs compared to DOT.

Some important gaps in the literature relating to cost of VOT include within trial cost-effectiveness analyses, differential costs between *synchronous* and *asynchronous* VOT, and incorporating the impact of ownership, loans, or access to internet-enabled smartphone devices.

Effectiveness

The overarching evidence from diverse studies including early trials and post-implementation evaluations suggests that treatment outcomes in VOT are likely to be at least non-inferior to DOT. Other outcomes including treatment completion, bacteriological resolution, and mortality rate were similar to standard DOT. In line with these findings, WHO previously concluded that VOT was at least non-inferior to DOT in terms of TB treatment completion and mortality⁷⁶. However, it is clear that large-scale, randomised-controlled trials in diverse settings, especially in LMIC or with vulnerable populations including those with MDR-TB, are required.

Strengths & Limitations

This scoping review has updated, consolidated and synthesised the existing evidence base on the use of VOT – including a novel table summarising the VOT applications available - to support people with TB globally. The methods are repeatable and the quality scoring using CCAT is transparent.

The review had several limitations. First, reviewers were not blinded, risking bias, and the search criteria did not include studies prior to 1st January 2000. However, the subsequent snowballing of citations did not find relevant studies earlier than this date. We also found multiple international clinical trial protocols for VOT evaluation on trials registration sites that have yet to publish their results. Second, several studies acknowledged selection bias in their recruitment with few involving randomisation, which limits their generalizability and warrants caution in the interpretation of their findings. Third, there was limited evidence evaluating forms of VOT such as synchronous versus asynchronous, the platforms or applications used to implement VOT, and vulnerable, underserved groups were neglected with only one study of VOT in children and none focused on pregnant women or people with TB and HIV or other comorbidities. These are notable gaps in the literature and understanding how VOT could be adapted for these populations is crucial for a comprehensive TB programme service delivery.

Research Recommendations

There is significant potential for future research to move the field of VOT for TB forward, especially in high TB burden LMIC. Randomised-controlled trials including those with multi-arm multi-stage or hybrid effectiveness-implementation designs and incorporating distributional or extended cost-effectiveness analyses and recognised WHO TB Patient Cost Survey methods⁷⁷ could be of great value to evaluate not only effectiveness but also feasibility, acceptability, cost-effectiveness and equity of VOT. However, RCTs can be expensive and time-consuming and, programmatically, other studies or implementation projects including the use of quasi-experimental designs or interrupted time-series analysis may be more suitable.

Wider research including ecological studies and secondary analysis of existing household survey data, including data on access to phones and internet, could examine how increased penetrance of smartphone coverage and usage in LMIC could contribute to VOT being integrated within standard of care. Moreover, there has been minimal exploration of sociocultural factors associated with mobile phone usage and ownership, particularly in LMIC where there may be a gender differential in household-level economic agency. Understanding such sociocultural factors can impact VOT feasibility and hence be crucial for optimising its effectiveness. Future studies implementing and evaluating VOT would benefit from considerations of equity within their analyses, including disaggregating data by gender, urban versus rural location, socioeconomic position, educational level, literacy, and digital literacy.

Finally, broader potentials of internet-enabled smartphone technology in the future could be, alongside VOT, screening for TB in households, for example via acoustic cough analysis,⁸² monitoring TB preventive therapy adherence and completion, reviewing symptoms of chronic respiratory diseases in people with suspected TB but no formal diagnosis, and adjunctive features to monitor side-effects over the course of a treatment regimen.

Conclusion

The COVID-19 pandemic has accelerated acceptance of digital health technologies, their implementation, and utility. This scoping review found that VOT was reported to be a feasible, acceptable, and effective alternative to DOT, which was likely cost-saving to people with TB and health systems in HIC. More evidence is needed to inform its implementation and potential scale-up in LMICs with high TB burden.

References

1. World Health Organisation. Tuberculosis. *World Health Organisation* <https://www.who.int/news-room/fact> (2023). Last accessed: 01/24/25 02:01pm.
2. WHO. The top 10 causes of death. *WHO* <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death> (2020). Last accessed: 01/24/25 02:03pm.
3. WHO. Global Tuberculosis Report 2022. *World Health Organisation* <https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2022> (2022). Last accessed: 01/24/25 02:04pm.
4. Alipanah, N. et al. Adherence interventions and outcomes of tuberculosis treatment: A systematic review and meta-analysis of trials and observational studies. *PLoS Med* **15**, (2018). Last accessed: 01/24/25 02:06pm.
5. Garfein, R. S. & Doshi, R. P. Synchronous and asynchronous video observed therapy (VOT) for tuberculosis treatment adherence monitoring and support. *J Clin Tuberc Other Mycobact Dis* **17**, (2019). Last accessed 01/24/25 02:07pm.
6. Nuttall, C. et al. Interventions pathways to reduce tuberculosis-related stigma: a literature review and conceptual framework. *Infect Dis Poverty* **11**, (2022). Last accessed: 01/24/25 02:08pm.
7. WHO Regional Office for Europe. Quick guide to video-supported treatment of tuberculosis. *World Health Organization*. <https://iris.who.int/bitstream/handle/10665/335900/9789289055222-eng.pdf> (2020). Last accessed: 01/24/25 02:10pm.
8. Stoner, M. C. D., Maragh-Bass, A. C., Sukhija-Cohen, A. C. & Saberi, P. Digital directly observed therapy to monitor adherence to medications: a scoping review. *HIV Res Clin Pract* **23**, 47–60 (2022). Last accessed: 02:10pm.
9. Story, A. et al. Smartphone-enabled video-observed versus directly observed treatment for tuberculosis: a multicentre, analyst-blinded, randomised, controlled superiority trial. *LANCET* **393**, 1216–1224 (2019). Last accessed: 02:11pm.
10. Falzon, D., Jaramillo, E. & World Health Organisation. Digital health for the end TB strategy: An agenda for action. *European Respiratory Society* (2015). Last accessed: 01/24/25 02:12pm.
11. Asay, G. R. B. et al. Cost of Tuberculosis Therapy Directly Observed on Video For Health Departments and Patients in New York City; San Francisco, California; and Rhode Island (2017-2018). *Am J Public Health* **110**, 1696–1703 (2020). Last accessed: 01/24/25 02:13pm.
12. Anne Delaporte, K. B. *The State of Mobile Internet Connectivity 2022*. <https://www.gsma.com/r/wp-content/uploads/2022/12/The-State-of-Mobile-Internet-Connectivity-Report-2022.pdf> (2022). Last accessed: 01/24/25 02:13pm.
13. Peters, M. D. J. et al. Scoping reviews: reinforcing and advancing the methodology and application. *Syst Rev* **10**, 263 (2021). Last accessed: 01/24/25 02:14pm.
14. Munn, Z. et al. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* **18**, 143 (2018). Last accessed: 01/24/25 02:16.
15. Cochrane. PICO ontology. *Cochrane* <https://linkeddata.cochrane.org/pico-ontology> (2023). Last accessed: 01/24/25 02:16.

16. Crowe, M., Sheppard, L. & Campbell, A. Comparison of the effects of using the Crowe Critical Appraisal Tool versus informal appraisal in assessing health research: a randomised trial. *Int J Evid Based Healthc* **9**, 444–449 (2011).
17. DeMaio, J., Schwartz, L., Cooley, P. & Tice, A. The Application of Telemedicine Technology to a Directly Observed Therapy Program for Tuberculosis: A Pilot Project. *Clinical Infectious Diseases* **33**, [2082–2084 \(2001\)](#). Last accessed: 01/25/24 02:17pm.
18. Rajalahti, I., Kreivi, H.-R., Ollgren, J. & Vasankari, T. Asynchronous video supported treatment of tuberculosis is well adopted in a real-world setting - an observational study comparing two distinct applications. *Infectious diseases (London, England)* 1–4 Preprint at <https://doi.org/10.1080/23744235.2022.2160492> (2022). Last accessed: 01/25/24, 02:17pm.
19. Aisyah, D. N. *et al.* Knowledge, Attitudes, and Behaviors on Utilizing Mobile Health Technology for TB in Indonesia: A Qualitative Pilot Study. *Front Public Health* **8**, [\(2020\)](#). Last accessed: 01/25/24 02:17pm.
20. Beeler Asay, G. R. *et al.* Cost of Tuberculosis Therapy Directly Observed on Video for Health Departments and Patients in New York City; San Francisco, California; and Rhode Island (2017–2018). *Am J Public Health* **110**, [1696–1703 \(2020\)](#). Last accessed: 01/25/24 02:18.
21. Bachina, P. *et al.* Programmatic Adoption and Implementation of Video-Observed Therapy in Minnesota: Prospective Observational Cohort Study. *JMIR Form Res* **6**, [\(2022\)](#). Last accessed: 02:24pm.
22. Bendiksen, R., Ovesen, T., Asfeldt, A.-M., Halvorsen, D. S. & Gravningen, K. Use of video directly observed treatment for tuberculosis in Northern Norway. *Tidsskrift for den Norske lægeforening : tidsskrift for praktisk medicin, ny række* vol. 140 Preprint at <https://doi.org/10.4045/tidsskr.19.0322> (2020). Last accessed: 01/25/24 02:24.
23. Bommakanti, K. K. *et al.* Requiring smartphone ownership for mHealth interventions: who could be left out? *BMC Public Health* **20**, (2020). Last accessed: 01/25/24 [02:26pm](#).
24. Buchman, T. & Cabello, C. A New Method to Directly Observe Tuberculosis Treatment: Skype Observed Therapy, a Patient-Centered Approach. *Journal of Public Health Management and Practice* **23**, 175–177 (2017). Last Accessed: 01/25/24 [02:27pm](#).
25. Burzynski, J. *et al.* In-Person vs Electronic Directly Observed Therapy for Tuberculosis Treatment Adherence: A Randomized Noninferiority Trial. *JAMA network open* vol. 5 e2144210 Preprint at <https://doi.org/10.1001/jamanetworkopen.2021.44210> (2022). Last accessed: 01/24/25 02:30pm.
26. Casalme, D. J. O. *et al.* Feasibility and acceptability of asynchronous VOT among patients with MDR-TB. *INTERNATIONAL JOURNAL OF TUBERCULOSIS AND LUNG DISEASE* **26**, [760–765 \(2022\)](#). Last accessed: [01/24/25 02:31pm](#).
27. Chen, S.-H. *et al.* Advantage in privacy protection by using synchronous video observed treatment enhances treatment adherence among patients with latent tuberculosis infection. *Journal of infection and public health* vol. 13 1354–1359 Preprint at <https://doi.org/10.1016/j.jiph.2020.03.013> (2020). Last accessed: 01/25/24 02:32.

28. Chuck, C., Robinson, E., Macaraig, M., Alexander, M. & Burzynski, J. Enhancing management of tuberculosis treatment with video directly observed therapy in New York City. *INTERNATIONAL JOURNAL OF TUBERCULOSIS AND LUNG DISEASE* **20**, [588–593 \(2016\)](#). Last accessed: 01/24/25 03:21pm.
29. Do, D., Garfein, R. S., Cuevas-Mota, J., Collins, K. & Liu, L. Change in Patient Comfort Using Mobile Phones Following the Use of an App to Monitor Tuberculosis Treatment Adherence: Longitudinal Study. *JMIR Mhealth Uhealth* **7**, [e11638 \(2019\)](#). Last accessed: 01/24/25 03:22pm.
30. Doltu, S. *et al.* Short and long-term outcomes of video observed treatment in tuberculosis patients, the Republic of Moldova. *Journal of infection in developing countries* vol. 15 17S-24S Preprint at <https://doi.org/10.3855/jidc.14601> (2021). Last accessed: 01/24/25 03:24pm.
31. Donahue, M. L., Eberly, M. D. & Rajnik, M. Tele-TB: Using TeleMedicine to Increase Access to Directly Observed Therapy for Latent Tuberculosis Infection. *Mil Med* **186**, [25–31 \(2021\)](#). Last accessed: 01/24/25 03:25pm.
32. Garfein, R. S. *et al.* Feasibility of tuberculosis treatment monitoring by video directly observed therapy: a binational pilot study. *INTERNATIONAL JOURNAL OF TUBERCULOSIS AND LUNG DISEASE* **19**, [1057–1064 \(2015\)](#). Last accessed: 01/24/24 03:27pm.
33. Garfein, R. S. *et al.* Tuberculosis Treatment Monitoring by Video Directly Observed Therapy in 5 Health Districts, California, USA. *Emerg Infect Dis* **24**, [1806–1815 \(2018\)](#). Last accessed: 01/24/25 03:28pm.
34. Garfein, R. S. *et al.* Evaluation of recorded video-observed therapy for anti-tuberculosis treatment. *The International Journal of Tuberculosis and Lung Disease* **24**, [520–525 \(2020\)](#). Last accessed: 01/24/25 03:29pm.
35. Gassanov, M. A. *et al.* The Use of Videophone for Directly Observed Therapy for the Treatment of Tuberculosis. *Canadian Journal of Public Health* **104**, [e272–e272 \(2013\)](#). Last accessed: 01/24/25 03:31pm.
36. Guo, P., Qiao, W., Sun, Y., Liu, F. L. & Wang, C. T. Telemedicine Technologies and Tuberculosis Management: A Randomized Controlled Trial. *TELEMEDICINE AND E-HEALTH* **26**, [1150–1156 \(2020\)](#).
37. Guo, X. *et al.* A Comprehensive App That Improves Tuberculosis Treatment Management Through Video-Observed Therapy: Usability Study. *JMIR mHealth and uHealth* vol. 8 e17658 Preprint at <https://doi.org/10.2196/17658> (2020). Last accessed: 01/24/25 9 03:34.
38. Anjum Hira, F., Khalid, H., Shashikala, A. & Md Moshiul, A. Factors of Video Directly Observed Therapy Adoption in Malaysia. *International Journal of Academic Research in Business and Social Sciences* **12**, [\(2022\)](#). Last accessed: 01/24/25 03:35pm.
39. Hoffman, J. A. *et al.* Mobile direct observation treatment for tuberculosis patients: a technical feasibility pilot using mobile phones in Nairobi, Kenya. *American journal of preventive medicine* vol. 39 78–80 Preprint at <https://doi.org/10.1016/j.amepre.2010.02.018> (2010). Last accessed: 01/24/25 03:37pm.
40. Holzman, S. B., Zenilman, A. & Shah, M. Advancing Patient-Centered Care in Tuberculosis Management: A Mixed-Methods Appraisal of Video Directly Observed Therapy. *Open Forum Infect Dis* **5**, [\(2018\)](#). Last accessed: 01/24/25 03:38pm.
41. Holzman, S. B. *et al.* Use of Smartphone-Based Video Directly Observed Therapy (vDOT) in Tuberculosis Care: Single-Arm, Prospective Feasibility

- Study. *JMIR formative research* vol. 3 e13411 Preprint at <https://doi.org/10.2196/13411> (2019). Last accessed: 03:39pm.
42. Kumar, A. A. *et al.* Mobile Health for Tuberculosis Management in South India: Is Video-Based Directly Observed Treatment an Acceptable Alternative? *JMIR Mhealth Uhealth* **7**, [e11687](https://doi.org/10.2196/13411) (2019). Last accessed: 01/24/25 02:41.
 43. Krueger, K. *et al.* Videophone utilization as an alternative to directly observed therapy for tuberculosis. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease* vol. 14 779–781 Preprint at <https://discovery.ebsco.com/linkprocessor/plink?id=0c508780-6d21-3346-ad93-76c45d7bd5b1> (2010). Last accessed: 01/24/25 03:41pm.
 44. Lam, C. K. *et al.* Using Video Technology to Increase Treatment Completion for Patients With Latent Tuberculosis Infection on 3-Month Isoniazid and Rifapentine: An Implementation Study. *J Med Internet Res* **20**, [e11687](https://doi.org/10.2196/13411) (2018). Last accessed: 01/25/24 03:43pm.
 45. Lam, C. K., Fluegge, K., Macaraig, M. & Burzynski, J. Cost savings associated with video directly observed therapy for treatment of tuberculosis. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease* vol. 23 1149–1154 Preprint at <https://doi.org/10.5588/ijtld.18.0625> (2019). Last accessed: 01/25/24 03:43.
 46. Lippincott, C., Perry, A., Munk, E., Maltas, G. & Shah, M. Tuberculosis treatment adherence in the era of COVID-19. *BMC Infect Dis* **22**, [e11687](https://doi.org/10.1186/s12875-022-11111-1) (2022). Last accessed: 01/24/25 03:44pm
 47. Macaraig, M., Lobato, M. N., Pilote, K. M. & Wegener, D. A National Survey on the Use of Electronic Directly Observed Therapy for Treatment of Tuberculosis. *JOURNAL OF PUBLIC HEALTH MANAGEMENT AND PRACTICE* **24**, [567–570](https://doi.org/10.1177/1078143518781435) (2018). Last accessed: 01/25/24 03:46pm.
 48. Mirsaiedi, M. *et al.* Video directly observed therapy for treatment of tuberculosis is patient-oriented and cost-effective. *European Respiratory Journal* **46**, [871–874](https://doi.org/10.1183/13996622.015000002015) (2015). Last accessed: 01/25/24 03:48pm.
 49. Molton, J. S. *et al.* Prospective single-arm interventional pilot study to assess a smartphone-based system for measuring and supporting adherence to medication. *BMJ open* vol. 6 e014194 Preprint at <https://doi.org/10.1136/bmjopen-2016-014194> (2016). Last accessed: 01/25/24 03:48pm.
 50. Nguyen, T. A. *et al.* Video Directly Observed Therapy to support adherence with treatment for tuberculosis in Vietnam: A prospective cohort study. *INTERNATIONAL JOURNAL OF INFECTIOUS DISEASES* **65**, [85–89](https://doi.org/10.1016/j.ijid.2017.05.001) (2017). Last accessed: 01/25/24 03:52pm.
 51. Nsengiyumva, N. P. *et al.* Evaluating the potential costs and impact of digital health technologies for tuberculosis treatment support. *EUROPEAN RESPIRATORY JOURNAL* **52**, [e18001](https://doi.org/10.1183/13996622.015000002018) (2018). Last accessed: 01/25/24 03:53pm.
 52. Perry, A. *et al.* Real-world implementation of video-observed therapy in an urban TB program in the United States. *INTERNATIONAL JOURNAL OF TUBERCULOSIS AND LUNG DISEASE* **25**, [655–661](https://doi.org/10.1183/13996622.015000002021) (2021). Last accessed: 01/25/24 03:54pm.
 53. Rabinovich, L. *et al.* Perceptions and Acceptability of Digital Interventions Among Tuberculosis Patients in Cambodia: Qualitative Study of Video-Based Directly Observed Therapy. *Journal of medical Internet research* vol. 22 e20212 (2021). Last accessed: 01/25/24 03:54pm.

- e16856 Preprint at <https://doi.org/10.2196/16856> (2020). Last accessed: 01/25/24 03:55pm.
54. Rao, J. S. *et al.* Acceptability of video observed treatment vs. directly observed treatment for tuberculosis: a comparative analysis between South and Central India. *Wellcome Open Res* **7**, [152 \(2022\)](#). Last accessed: 01/25/24 03:55pm.
 55. Ravenscroft, L. *et al.* Video-observed therapy and medication adherence for tuberculosis patients: randomised controlled trial in Moldova. *EUROPEAN RESPIRATORY JOURNAL* **56**, [\(2020\)](#). Last accessed: 01/25/24 03:56pm.
 56. Ritter, L. A., Kwong, M. W. & Nasser, L. M. California Public Health Departments Remotely Treat Tuberculosis: Outcomes & Opportunities. *Calif J Health Promot* **15**, [37–45 \(2017\)](#). Last accessed: 01/25/24 03:57pm.
 57. Amoo Olufemi Samuel *et al.* Impact of a Digital Health Platform (NimCure) on Adherence Enhancement in Tuberculosis Therapy. *African Scientist* **22**, [49–57 \(2021\)](#). Last accessed: 01/25/24 03:59pm.
 58. Sekandi, J. N. *et al.* Video directly observed therapy for supporting and monitoring adherence to tuberculosis treatment in Uganda: a pilot cohort study. *ERJ Open Res* **6**, [\(2020\)](#). Last accessed: 01/25/24 03:59pm.
 59. Sekandi, J. N. *et al.* Stakeholders' Perceptions of Benefits of and Barriers to Using Video-Observed Treatment for Monitoring Patients With Tuberculosis in Uganda: Exploratory Qualitative Study. *JMIR Mhealth Uhealth* **9**, [\(2021\)](#).
 60. Sinkou, H. *et al.* Video-observed treatment for tuberculosis patients in Belarus: findings from the first programmatic experience. *European Respiratory Journal* **49**, [1602049 \(2017\)](#). Last accessed: 01/24/25 04:02pm.
 61. Wade, V. A., Karnon, J., Elliott, J. A. & Hiller, J. E. Home Videophones Improve Direct Observation in Tuberculosis Treatment: A Mixed Methods Evaluation. *PLoS One* **7**, [e50155 \(2012\)](#). Last accessed: 01/25/24 04:03pm.
 62. Zúñiga, M. L. *et al.* A Qualitative Study Exploring Stakeholder Perceptions of Video Directly Observed Therapy for Monitoring Tuberculosis Treatment in the US-Mexico Border Region. *J Mob Technol Med* **5**, [12–23 \(2016\)](#). Last accessed: 01/25/24 04:04pm.
 63. Abbara, A. *et al.* The challenges of tuberculosis control in protracted conflict: The case of Syria. *International Journal of Infectious Diseases* **90**, [53–59 \(2020\)](#). Last accessed: 01/24/25 04:05pm.
 64. Brett, K. & Severn, M. *Direct Observational Therapy for the Treatment of Tuberculosis: A Review of Clinical Evidence and Guidelines*. [\(Canadian Agency for Drugs and Technologies in Health, 2020\)](#). Last accessed: 01/25/24 04:06pm.
 65. Chakaya, J. *et al.* Global Tuberculosis Report 2020 - Reflections on the Global TB burden, treatment and prevention efforts. *Int J Infect Dis* **113 Suppl 1**, [S7–S12 \(2021\)](#). Last accessed: 01/25/24 04:12pm.
 66. Keutzer, L., Wicha, S. G. & Simonsson, U. S. Mobile Health Apps for Improvement of Tuberculosis Treatment: Descriptive Review. *JMIR mHealth and uHealth* vol. 8 e17246 Preprint at <https://doi.org/10.2196/17246> (2020). Last accessed: 01/25/24 04:13pm.
 67. Lee, Y., Raviglione, M. C. & Flahault, A. Use of Digital Technology to Enhance Tuberculosis Control: Scoping Review. *J Med Internet Res* **22**, [\(2020\)](#). Last accessed: 01/25/24 04:14.
 68. Digital, C. G. I. *et al.* The impact of digital health technologies on tuberculosis treatment: a systematic review. *EUROPEAN RESPIRATORY JOURNAL* **51**, [\(2018\)](#). Last accessed: 01/25/24 04:15pm.

69. Ridho, A. *et al.* Digital Health Technologies to Improve Medication Adherence and Treatment Outcomes in Patients With Tuberculosis: Systematic Review of Randomized Controlled Trials. *J Med Internet Res* **24**, (2022). Last accessed: 01/25/24 04:15pm.
70. Subbaraman, R. *et al.* Digital adherence technologies for the management of tuberculosis therapy: mapping the landscape and research priorities. *BMJ Glob Health* **3**, e001018 (2018). Last accessed: 01/25/24 04:17pm.
71. Truong, C. B., Tanni, K. A. & Qian, J. Video-Observed Therapy Versus Directly Observed Therapy in Patients With Tuberculosis. *American journal of preventive medicine* vol. 62 450–458 Preprint at <https://doi.org/10.1016/j.amepre.2021.10.013> (2022). Last accessed: 01/25/24 04:17pm.
72. Wong, Y. J., Ng, K. Y. & Lee, S. W. H. Digital health use in latent tuberculosis infection care: A systematic review. *Int J Med Inform* **159**, 104687 (2022). Last accessed: 01/25/24 04:18pm.
73. Implementing an Electronic Directly Observed Therapy (eDOT) Program: A Toolkit for Tuberculosis (TB) Programs. Last accessed: 01/25/24 04:19pm
74. European Centre for Disease Prevention and Control. & Sandgren, A. Guidance on tuberculosis control in vulnerable and hard-to-reach populations. *ECDC* (2016) doi:10.2900/72431. Last accessed: 01/24/25 04:20pm.
75. World Health Organization & Farr. Handbook for the use of digital technologies to support tuberculosis medication adherence. *WHO, Geneva* <https://iris.who.int/bitstream/handle/10665/259832/9789241513456-eng.pdf;jsessionid=A32F9A9DE6CE8248EF661C71B9C80283?sequence=1> (2017). Last accessed: 01/25/24 04:21pm.
76. World Health Organization. *Guidelines for treatment of drug-susceptible tuberculosis and patient care, 2017 update*. <https://iris.who.int/bitstream/handle/10665/255052/9789241550000-eng.pdf> (2017). Last accessed: 01/25/24 04:21pm.
77. Centre for Disease Control and Prevention. Tuberculosis: Data and Statistics. <https://www.cdc.gov/tb/statistics/default.htm> (2023). Last accessed: 01/25/24 04:21pm.
78. WHO Nigeria. Gearing towards a TB free Nigeria- WHO and partners scale up action. *World Health Organisation* <https://www.afro.who.int/news/gearing-towards-tb-free-nigeria-who-and-partners-scale-action> (2021). Last accessed: 01/25/24 04:22pm.
79. Iribarren, S. J., et al. (2016) 'Smartphone Applications to Support Tuberculosis Prevention and Treatment: Review and Evaluation', *JMIR Mhealth Uhealth*, 4(2). Available at: <https://mhealth.jmir.org/2016/2/e25/> (Accessed: August 2023). Last accessed: 01/25/24 04:23pm.
80. Saunders, M.J., Wingfield, T., Tovar, M.A., Herlihy, N., Rocha, C., Zevallos, K., Montoya, R., Ramos, E., Datta, S. and Evans, C.A. (2018) 'Mobile phone interventions for tuberculosis should ensure access to mobile phones to enhance equity – a prospective, observational cohort study in Peruvian shantytowns', *Trop Med Int Health*, 23(8), pp. 850-859. Available at: <https://doi.org/10.1111/tmi.13087> (Accessed: August 2023). Last accessed: 01/25/24 04:24pm.
81. Zamudio, C., et al. (2015) 'Public transportation and tuberculosis transmission in a high incidence setting', *PLoS One*. Available at: doi:10.1371/journal.pone.0115230 Last accessed: 01/25/24 04:24pm.

82. [Kosha C. \(n.d.\). 50TH World Conference on Lung Health of The International Union Against Tuberculosis and Lung Disease \(The Union\) The Official Journal of the International Union Against Tuberculosis and Lung Disease. Retrieved January 8, 2024, from <https://jata.or.jp/english/> Last accessed: 01/25/24 04:25pm.](#)

Ethical Considerations

There were no ethical issues or risk of harm to patients identified. Ethical approval exemption was sought and granted by LSTM. All the resources used were based on published research with previously anonymised participant demographics and outcomes. This study was nested within the larger “SAFEST-1 MDR-TB” programme of research, which includes an implementation study of VOT for people with MDR-TB in Mozambique. SAFEST-1 MDR-TB was approved by Research Ethics Committees at the Liverpool School of Tropical Medicine, UK (23-008) and the Comité Institucional de Ética do Instituto Nacional de Saude, Mozambique (148/CIE-INS/2023, 15th of November, 2023).

Conflicts of Interest & Funding Declaration


There were no conflicts of interest declared. ADT and CK and TW are affiliated with The Liverpool School of Tropical Medicine (LSTM), UK. KP and CK are affiliated with Instituto Nacional de Saúde (INS), Mozambique. CK is affiliated to Faculdade de Medicina, Eduardo Mondlane University, Mozambique and EM is affiliated with The Royal Liverpool University Hospital, UK. This scoping review has not sought or received any specific funding. However, it does form an output of a part of a larger study for the “User and Provider Acceptability and Feasibility of Video Observed Therapy for Multi-Drug Resistant Tuberculosis in Mozambique (SAFEST-1 MDR-TB)” for which funding of £30,000 was granted and the grant number is MRF-131-0006-RG-KHOS-C0942 (Medical Research Foundation, 2023).

TW is supported by grants from the Wellcome Trust, UK (209075/Z/17/Z), the Department of Health and Social Care (DHSC), the Foreign, Commonwealth & Development Office (FCDO), the Medical Research Council (MRC) and Wellcome, UK. Medical Research Council, Department for International Development, and Wellcome Trust (Joint Global Health Trials, MR/V004832/1), Medical Research Council (Public Health Intervention Development “PHIND” Award, MR/Y503216/1), and a Medical Research Foundation Meade Collaboration Grant in Epidemiology.

Availability of data and materials

The data are available from the corresponding author (email address) upon reasonable request.

Figure 1: PRISMA-ScR flow diagram



*Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/registers).

**If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools.

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>

Table 1: Primary Research Studies (n=47)

Author & Year	Title	Self-reported Study type and design / Country / WHO Region* / Income group	Total participants (n)	CCAT score (%) and grading
Aisyah et al. 2020 ¹⁹	Knowledge, Attitudes, and Behaviors on Utilizing Mobile Health Technology for TB in Indonesia: A Qualitative Pilot Study	Pilot study (qualitative) Indonesia SEAR LMIC	46	33 (82.5%) High quality
Asay et al. 2020 ²⁰	Cost of Tuberculosis Therapy Directly Observed on Video for Health Departments and Patients in New York City; San Francisco, California; and Rhode Island (2017-2018)	Survey USA AMR High Income	87	33 (82.5%) High quality
Bachina et al. 2022 ²¹	Programmatic Adoption and Implementation of Video-Observed Therapy in Minnesota: Prospective Observational Cohort Study	Observational USA AMR High Income	49	32 (80%) High quality
Bendiksen et al. 2020 ²²	Use of video directly observed treatment for tuberculosis in Northern Norway	Observation Norway EUR High Income	17	30 (75%) High Quality
Bommakantiet al. 2020 ²³	Requiring smartphone ownership for mHealth interventions: who could be left out?	Secondary analysis of data from single-arm trial USA AMR High Income	151	33 (82.5%) High quality
Buchman et al. 2017 ²⁴	A New Method to Directly Observe Tuberculosis Treatment: Skype Observed Therapy, a Patient-Centered Approach	Retrospective USA AMR High Income	24	29 (73%) Moderate quality
Burzynski et al. 2022 ²⁵	In-Person vs Electronic Directly Observed Therapy for Tuberculosis Treatment Adherence: A Randomized Noninferiority Trial	Observational (with cross-over and initial randomisation) USA	216	36 (90%) High quality

		AMR High Income		
Casalme et al. 2022 ²⁶	Feasibility and acceptability of asynchronous VOT among patients with MDR-TB	Observational The Philippines WPR LMIC	308	29 (72.5%) Moderate quality
Chen et al. 2020 ²⁷	Advantage in privacy protection by using synchronous video observed treatment enhances treatment adherence among patients with latent tuberculosis infection	Survey Taiwan WPR High Income	240	31 (77.5%) High quality
Chuck et al. 2016 ²⁸	Enhancing management of tuberculosis treatment with video directly observed therapy in New York City	Prospective USA AMR High Income	329	31 (77.5%) High quality
DeMaio et al. 2001 ¹⁷	The Application of Telemedicine Technology to a Directly Observed Therapy Program for Tuberculosis: A Pilot Project	Pilot single-arm USA AMR High Income	6	29 (72.5%) Moderate quality
Do et al. 2019 ²⁹	Change in Patient Comfort Using Mobile Phones Following the Use of an App to Monitor Tuberculosis Treatment Adherence: Longitudinal Study	Prospective USA AMR High Income	120	33 (82.5%) High quality
Doltu et al. 2021 ³⁰	Short and long-term outcomes of video observed treatment in tuberculosis patients, the Republic of Moldova	Observational Moldova EUR LMIC	647	33 (82.5%) High quality
Donahue et al. 2021 ³¹	Tele-TB: Using TeleMedicine to Increase Access to Directly Observed Therapy for Latent Tuberculosis Infection	Prospective USA AMR High Income	20	30 (75%) High quality

Garfein et al. 2015 ³²	Feasibility of tuberculosis treatment monitoring by video directly observed therapy: a binational pilot study	Pilot single-arm USA & Mexico AMR High Income & Upper Middle Income	52 (USA 43 + Mexico 9)	35 (87.5%) High quality
Garfein et al. 2018 ³³	Tuberculosis Treatment Monitoring by Video Directly Observed Therapy in 5 Health Districts, California, USA	Prospective USA AMR High Income	274	33 (82.5%) High quality
Garfein et al. 2020 ³⁴	Evaluation of recorded video-observed therapy for anti-tuberculosis treatment	Pilot single-arm USA AMR High Income	149	31 (77.5%) High quality
Gassanov et al. 2013 ³⁵	The use of videophone for directly observed therapy for the treatment of tuberculosis	Letter to the Editor (with original data) Canada AMR High Income	13	21 (52.5%) Moderate quality
Guo P. et al. 2020 ³⁶	Telemedicine Technologies and Tuberculosis Management: A Randomized Controlled Trial	RCT China WPR Upper Middle Income	405	32 (80%) High quality
Guo X. et al. 2020 ³⁷	A Comprehensive App That Improves Tuberculosis Treatment Management Through Video-Observed Therapy: Usability Study	Retrospective China WPR Upper Middle Income	290	29 (72.5%) Moderate quality
Hira et al. 2022 ³⁸	Factors of Video Directly Observed Therapy Adoption in Malaysia	Survey (cross-sectional) Malaysia WPR Upper Middle Income	150	28 (70%) Moderate quality

Hoffman et al. 2010 ³⁹	Mobile direct observation treatment for tuberculosis patients: a technical feasibility pilot using mobile phones in Nairobi, Kenya	Pilot study Kenya AFR LMIC	13	29 (72.5%) Moderate quality
Holzman et al. 2018 ⁴⁰	Advancing Patient-Centered Care in Tuberculosis Management: A Mixed-Methods Appraisal of Video Directly Observed Therapy	Prospective USA AMR High Income	28	34 (85%) High Quality
Holzman et al. 2019 ⁴¹	Use of Smartphone-Based Video Directly Observed Therapy (vDOT) in Tuberculosis Care: Single-Arm, Prospective Feasibility Study	Prospective India SEAR LMIC	25	34 (85%) High quality
Kumar et al. 2019 ⁴²	Mobile Health for Tuberculosis Management in South India: Is Video-Based Directly Observed Treatment an Acceptable Alternative?	Exploratory (acceptability study) India SEAR LMIC	185	34 (85%) High quality
Krueger et al. 2010 ⁴³	Videophone utilization as an alternative to directly observed therapy for tuberculosis	Retrospective USA AMR High Income	57	28 (70%) Moderate quality
Lam et al. 2018 ⁴⁴	Using Video Technology to Increase Treatment Completion for Patients With Latent Tuberculosis Infection on 3-Month Isoniazid and Rifapentine: An Implementation Study	Prospective USA AMR High Income	71	33 (82.5%) High quality
Lam et al. 2019 ⁴⁵	Cost savings associated with video directly observed therapy for treatment of tuberculosis	Retrospective USA AMR High Income	n/a - VOT sessions reported	32 (80%) High quality
Lippincott et al. 2022 ⁴⁶	Tuberculosis treatment adherence in the era of COVID-19	Retrospective USA AMR	52	34 (85%) High quality

		High Income		
Macaraig et al. 2018 ⁴⁷	A National Survey on the Use of Electronic Directly Observed Therapy for Treatment of Tuberculosis	Survey USA AMR High Income	47	34 (85%) High quality
Mirsaeidi et al. 2015 ⁴⁸	Video directly observed therapy for treatment of tuberculosis is patient-oriented and cost-effective	Letter to the Editor (with original data) USA AMR High Income	11	26 (65%) Moderate quality
Molton et al. 2016 ⁴⁹	Prospective single-arm interventional pilot study to assess a smartphone-based system for measuring and supporting adherence to medication	Prospective Singapore WPR High Income	42	36 (90%) High quality
Nguyen et al. 2017 ⁵⁰	Video Directly Observed Therapy to support adherence with treatment for tuberculosis in Vietnam: A prospective cohort study	Prospective Vietnam WPR LMIC	40	32 (80%) High quality
Nsengiyumva et al. 2018 ⁵¹	Evaluating the potential costs and impact of digital health technologies for tuberculosis treatment support	Prospective Brazil AMR Upper Middle Income	N/A Simulated cost analysis	30 (75%) High quality
Perry et al. 2021 ⁵²	Real-world implementation of video-observed therapy in an urban TB program in the United States	Prospective USA AMR High Income	163	30 (75%) High quality
Rabinovich et al. 2020 ⁵³	Perceptions and Acceptability of Digital Interventions Among Tuberculosis Patients in Cambodia: Qualitative Study of Video-Based Directly Observed Therapy	Cross-sectional (qualitative) Cambodia WPR LMIC	64	33 (82.5%) High quality

Rajalahti et al. 2022 ¹⁸	Asynchronous video supported treatment of tuberculosis is well adopted in a real-world setting – an observational study comparing two distinct applications	Observational Finland EUR High Income	30	32 (80%) High quality
Rao et al. 2022 ⁵⁴	Acceptability of video observed treatment vs. directly observed treatment for tuberculosis: a comparative analysis between South and Central India	Exploratory (cross-sectional) India SEAR LMIC	351	31 (77.5%) High quality
Ravenscroft et al. 2020 ⁵⁵	Video-observed therapy and medication adherence for tuberculosis patients: randomised controlled trial in Moldova	RCT Moldova EUR LMIC	178	34 (85%) High quality
Ritter et al. 2017 ⁵⁶	California Public Health Departments Remotely Treat Tuberculosis: Outcomes & Opportunities	Survey USA AMR High Income	56	33 (83%) High quality
Samuel et al. 2021 ⁵⁷	Impact of a Digital Health Platform (NimCure) on Adherence Enhancement in Tuberculosis Therapy	RCT Nigeria AFR LMIC	85	29 (72.5%) Moderate quality
Sekandi et al. 2020 ⁵⁸	Video directly observed therapy for supporting and monitoring adherence to tuberculosis treatment in Uganda: a pilot cohort study	Pilot cohort study Uganda AFR Low Income	50	34 (85%) High quality
Sekandi et al. 2021 ⁵⁹	Stakeholders' Perceptions of Benefits of and Barriers to Using Video-Observed Treatment for Monitoring Patients With Tuberculosis in Uganda: Exploratory Qualitative Study	Exploratory (Qualitative) Uganda AFR Low Income	30	35 (87.5%) High quality
Sinkou et al. 2017 ⁶⁰	Video-observed treatment for tuberculosis patients in Belarus: findings from the first programmatic experience	Letter to the Editor (with original data)	10	32 (90%)

		Belarus EUR Upper Middle Income		High quality
Story et al. 2019 ⁹	Smartphone-enabled video-observed versus directly observed treatment for tuberculosis: a multicentre, analyst-blinded, randomised, controlled superiority trial	RCT UK EUR High Income	226	36 (90%) High quality
Wade et al. 2012 ⁶¹	Home videophones improve direct observation in tuberculosis treatment: a mixed methods evaluation	Retrospective Australia WPR High Income	128	38 (95%) High quality
Zúñiga et al. 2016 ⁶²	A Qualitative Study Exploring Stakeholder Perceptions of Video Directly Observed Therapy for Monitoring Tuberculosis Treatment in the US-Mexico Border Region	Qualitative (FGDs) USA & Mexico AMR High Income & Upper Middle Income	22 clients + 47 providers	32 (80%) High quality

*African Region (AFR) / Region of the Americas (AMR) / South-East Asian Region (SEAR) / European Region (EUR) / Eastern Mediterranean Region (EMR) / Western Pacific Region (WPR)

Table 2: Review-type Studies (n=13)

Author & Year	Title	Self-reported Study type and Design / Country of 1 st author / WHO Region*	Studies included (n)	PRISMA or similar reporting guidelines followed	CCAT score (%) and grading
Abbara et al. 2020 ⁶³	The challenges of tuberculosis control in protracted conflict: The case of Syria	Narrative review UK EUR	Not reported	No	33 (82.5%) High quality
Alipanah et al. 2018 ⁴	Adherence interventions and outcomes of tuberculosis treatment: A systematic review and meta-analysis of trials and observational studies	Systematic review with Meta-analysis USA AMR	129	Yes	37 (92.5%) High quality
Brett et al. 2020 ⁶⁴	Direct Observational Therapy for the Treatment of Tuberculosis: A Review of Clinical Evidence and Guidelines	Quasi-systematic review Canada AMR	3x systematic reviews, 6x RCTs, 6x evidence-based guidelines	Yes	35 (87.5%) High quality
Chakaya et al. 2021 ⁶⁵	Global Tuberculosis Report 2020 – Reflections on the Global TB burden, treatment and prevention efforts	Narrative review Kenya AFR	Not reported	No	36 (90%) High quality
Garfein et al. 2019 ⁵	Synchronous and asynchronous video observed therapy (VOT) for tuberculosis treatment adherence monitoring and support	Literature review (non-systematic) USA AMR	Not reported	No	28 (70%) Moderate quality
Keutzer et al. 2020 ⁶⁶	Mobile Health Apps for Improvement of Tuberculosis Treatment: Descriptive Review	Descriptive review Sweden EUR	55	Yes	29 (72.5%) Moderate quality
Lee et al. 2020 ⁶⁷	Use of Digital Technology to Enhance Tuberculosis Control: Scoping Review	Scoping review Switzerland EUR	145	Yes	33 (82.5%) High quality

Ngwatu et al. 2018 ⁶⁸	The impact of digital health technologies on tuberculosis treatment: a systematic review	Systematic review Switzerland EUR	7	Yes	36 (90%) High quality
Ridho et al. 2022 ⁶⁹	Digital Health Technologies to Improve Medication Adherence and Treatment Outcomes in Patients With Tuberculosis: Systematic Review of Randomized Controlled Trials	Systematic review of RCTs Indonesia SEAR	16	Yes	37 (92.5%) High quality
Stoner et al. 2022 ⁸	Digital directly observed therapy to monitor adherence to medications: a scoping review	Scoping review USA AMR	28	Yes	35 (85%) High quality
Subbaraman et al. 2018 ⁷⁰	Digital adherence technologies for the management of tuberculosis therapy: mapping the landscape and research priorities	Narrative review USA AMR	Not reported	No	33 (82.5%) High quality
Truong et al. 2022 ⁷¹	Video-Observed Therapy Versus Directly Observed Therapy in Patients With Tuberculosis	Systematic review and Meta-analysis USA AMR	9	Yes	35 (87.5%) High quality
Wong et al. 2022 ⁷²	Digital health use in latent tuberculosis infection care: A systematic review	Systematic review Malaysia WPR	15	Yes	36 (90%) High quality

Table 3 : Grey Literature documents (n=6)

Organisation (author) & Year	Title	Type of document	Country / WHO Region	Current TB burden, cases per 100,000*
CDC 2017 ⁷³	Implementing an Electronic Directly Observed Therapy (eDOT) Program: A Toolkit for Tuberculosis (TB) Programs	Toolkit	USA AMR	2.5
ECDC (Sandgren et al.) 2016 ⁷⁴	Guidance on tuberculosis control in vulnerable and hard-to-reach populations	Guideline	Europe EUR	10
WHO (Falzon et al.) 2015 ¹⁰	Digital health for the end TB strategy: An agenda for action	Strategy	Worldwide	127
WHO (Farr et al.) 2017 ⁷⁵	Handbook for the use of digital technologies to support tuberculosis medication adherence	Handbook	Worldwide	127
WHO 2017 ⁷⁶	Guidelines for treatment of drug susceptible tuberculosis and patient care	Guideline	Worldwide	127
WHO Europe 2020 ⁷	Quick guide to video-supported treatment of tuberculosis	Quick guide	Europe EUR	10

*Sources: WHO, 2021; ECDC, 2022; CDC, 2023

Table 4: Reported Successes and Challenges in High-income to Middle-income Countries

High-income to Middle-income			
Successes	Challenges	Potential solutions	Key studies
<ul style="list-style-type: none"> • Reported high adherence rates with VOT • VOT was feasible for people living with TB during the COVID-19 pandemic • VOT was feasible for people with (M)DR-TB • VOT was feasible for TB preventive therapy monitoring in a paediatric population • VOT uptake in individuals older than 65 was 10% in an urban setting in the USA • Ability to learn VOT was not associated with ownership of a smartphone • When scaled-up in urban settings of the USA, VOT was cost saving compared with DOT • HCWs preferred VOT over community DOT home visits • VOT was implemented more widely in London, UK based on outcomes from the trial conducted by Story et al. 2019 	<ul style="list-style-type: none"> • Health insurance policies and frameworks must recognise VOT as a treatment in order that patients have their costs reimbursed • Poor video quality • Network connectivity problems • Difficulty navigating VOT software / uploading videos • Personal smartphone not compatible for VOT software • Lack of interpersonal interaction reported by both clients and HCW • Misuse of loaned phone for non-VOT activities • Side-effects may go unreported with VOT usage • Concerns regarding privacy and data security • Older people, males, people with fewer years of formal education, and people with lower annual income were less likely to own smartphones posing problems for equity of access and use 	<ul style="list-style-type: none"> • Advocate for recognition of VOT as an established adherence monitoring modality, including both to National TB Programs but also to social protection and health insurance schemes and policy makers • Training of HCW and clients • Loaning smartphones compatible with VOT software • Encourage ad hoc communication and reporting of side-effects via VOT and/or integration of adverse effect monitoring or interactive platforms within VOT applications • Reassure clients regarding privacy and data security 	<p>Gassanov, 2013³⁵</p> <p>Chuck, 2016²⁸</p> <p>Ritter, 2017⁵⁶</p> <p>Garfein, 2018³³</p> <p>Lam, 2018⁴⁴</p> <p>Macaraig, 2018⁴⁷</p> <p>Story, 2019⁹</p> <p>Asay, 2020²⁰</p> <p>Bendiksen, 2020²²</p> <p>Bommakanti, 2020²³</p> <p>Garfein, 2020³⁴</p> <p>Donahue, 2021³¹</p> <p>Perry, 2021⁵²</p> <p>Bachina, 2022²¹</p> <p>Burzynski, 2022²⁵</p>
Low-income to lower-middle income			
Successes	Challenges	Potential solutions	Key studies
<ul style="list-style-type: none"> • VOT was feasible in LMIC and LIC contexts 	<ul style="list-style-type: none"> • Personal preference for DOT • Persistent non-adherence with VOT requiring switch back to DOT 	<ul style="list-style-type: none"> • Improve the evidence base on implementation strategies, scalability, effectiveness (vs DOT for adherence 	<p>Hoffman, 2010³⁹</p> <p>Nguyen, 2017⁵⁰</p>

<ul style="list-style-type: none"> • VOT was feasible for people living with MDR-TB in an LMIC context (The Phillipines) • Positive feedback from clients and HCWs for VOT • Clients could feedback treatment-related problems out of hours using asynchronous VOT • Asynchronous VOT had no sessional time limits • Clients felt empowered by ad hoc communication offered by asynchronous VOT • Opportunity for HCWs to engage in Continued Professional Development in learning requisite IT skills for VOT 	<ul style="list-style-type: none"> • Client unease with recording themselves taking their medications • Concerns regarding privacy following uploading of video-recording • Lack of private place to record VOT videos • Perceived lack of contact with HCW expressed in FGD about VOT in Cambodia • Slow uploading of asynchronous VOT videos due to issues with • Poor network connection, especially in rural areas • Competing for internet bandwidth with the host hospital to deliver VOT • Study smartphone not returned despite expectations it would be so at end of study implementation • Broken or uncharged smartphone 	<p>and treatment success), and cost effectiveness of VOT and disseminate to clients, potential clients, public, and HCWs</p> <ul style="list-style-type: none"> • Reassure clients regarding privacy and data security, consider encryption similar to other platforms including pins, signatures, and digital prints to access • Encourage ad hoc communication and reporting of side-effects via VOT and/or integration of adverse effect monitoring or interactive platforms within VOT applications • Help transition VOT clients between TB centres (if feasible) • Lobby stakeholders for wider mobile internet network reach into rural communities 	<p>Holzman, 2019⁴¹ Rabinovich, 2020⁵³ Sekandi, 2020⁵⁸ Samuel, 2021⁵⁷ Sekandi, 2021⁵⁹ Casalme, 2022²⁶ Rao, 2022⁵⁴</p>
--	--	--	---

Appendix/Supplemental Materials:

An example of the MeSH (Medical Subject Headings) search terms used were: (TI (VOT OR (video AND observed AND therapy) OR VST OR (video AND support AND treatment) OR vDOT OR eDOT)) AND (TI (TB OR tuberculosis)) OR (AB (VOT OR (video AND observed AND therapy) OR VST OR (video AND support AND treatment) OR vDOT OR eDOT)) AND (AB (TB OR tuberculosis)). These MeSH terms were piloted and refined between team members ADT, EM, and KP.

Abbreviations and Glossary

ADT – An Du Tinh

AMR – Antimicrobial resistance

CCAT – Crowe Critical Appraisal Tool

CDC – Centers for Disease Control and Prevention; American public health agency

CI – Confidence Interval

CK – Celso Khosa

Client – Person living with TB; a less stigmatising term than “TB patient”

CXR – Chest x-ray

DOT – Direct Observed Therapy; commonly associated with in-person adherence monitoring for TB treatment

DS-TB – drug-sensitive TB

ECDC – European Centre for Disease Prevention and Control - the European public health agency

eDOT - electronic-Direct Observed Therapy; VOT in a different name

EM – Eleanor Morgan

FGD – Focus Group Discussions – structured group interviews to collect qualitative data

GRADE - Grading of Recommendations, Assessment, Development, and Evaluations framework for evaluating research evidence summaries

HCW – Healthcare Worker

IGRA – Interferon-gamma release assay

INS - Instituto Nacional de Saúde

JBI - Joanna Briggs Institute; consensus framework for conducting Scoping Reviews

KP – Kassia Pereira

LMIC – Lower Middle Income Country as determined by The World Bank classification

LSTM – Liverpool School of Tropical Medicine

LTBI – Latent Tuberculosis Infection

MDR-TB – Multidrug Resistant Tuberculosis

MTB – Mycobacterium tuberculosis

NHS – National Health Service

NTP - National Tuberculosis Programme

PRISMA - Preferred Reporting Items for Systematic Reviews and Meta-Analyses

RCT – Randomised Control Trial

RR – Relative Risk

SAT – Self-Administered Therapy; TB treatments taken unobserved or monitored by healthcare professionals

ScR – Scoping Review; as in PRISMA-ScR

SDG – Sustainable Development Goals as set by the WHO/UN

SMS – Short Message/Messaging Service

TB – Tuberculosis

TST – Tuberculin skin test

TW – Tom Wingfield

vDOT - virtual/video-Directed Observed Therapy; VOT in a different name

VOT – Video Observed Therapy; also known as Video-Supported Treatment, electronic-Directly Observed Therapy and virtual/video-Directly Observed Therapy

VST - Video-Supported Treatment; VOT in a different name

WHO – World Health Organisation

UKHSA – United Kingdom Health Security Agency; agency formerly known as Public Health England (PHE)

Supplementary Table 1: Summary of VOT Applications

Name	Website & App stores	Country of Origin / WHO Region	Cost	Cited or used by
Emocha Health, now renamed Scene Health	www.scene.health or via Apple App Store and Google Play Store	USA AMR	Free to download the app, but the cost of the account subscription cost to on Scene Health not reported	Bachina, 2022 ²¹ ; Perry, 2021 ⁵² ; Lippincott, 2022 ⁴⁶ ; Holzman, 2019 ⁴¹ ; Holzman 2018 ⁴⁰ ; Keutzer 2020 ⁶⁶ ; Asay, 2020 ²⁰ ; CDC, 2017 ⁷³ ; Farr, 2017 ⁷⁵
SureAdhere	www.sureadhere.com owned by Dimagi; Apple App Store and Google Play Store	USA AMR	Starting from \$120/month	Garfein, 2019 ⁵ ; Casalme 2022 ²⁶ ; Keutzer, 2020 ⁶⁶ ; Asay, 2020 ²⁰ ; Nguyen, 2017 ⁵⁰ ; CDC, 2017 ⁷³ ; Farr, 2017 ⁷⁵ ; Subbaraman, 2018 ⁷⁰
Skype	www.skype.com owned by Microsoft; Apple App Store and Google Play Store	USA AMR	Free for internet calls only, variable costs otherwise dependent on the country	Asay, 2020 ²⁰ ; Mirsaeidi, 2015 ⁴⁸ ; Macaraig, 2018 ⁴⁷ ; CDC, 2017 ⁷³ ; Buchman, 2017 ²⁴
FaceTime	Pre-installed on Apple mobile devices; Apple App Store only, owned by Apple	USA AMR	Free for calls between Apple devices	Macaraig, 2018 ⁴⁷ ; CDC, 2017 ⁷³
AiCure	www.aicure.com Apple App store and Google Play store	USA AMR	Starting from \$750 per clinic per month for up to 50 patients (Salcedo et al., 2021)	Keutzer, 2020 ⁶⁶ ; CDC, 2017 ⁷³
Acano	Via Apple App Store only, owned by Cisco Systems	USA AMR	Free	Bendiksen, 2020 ²²
Fuze	www.fuze.com owned by 8x8 Company; Apple App Store and Google Play Store	USA AMR	Starting from \$15 per user per month	CDC, 2017 ⁷³
HipaaBridge	www.everbridge.com and Google Play Store	USA AMR	Free to download on Android, unclear from Ever Bridge website if there are	CDC, 2017 ⁷³

			other costs from website applicable following download	
ooVoo	www.oovoo.com but discontinued November 2017	USA AMR	Not reported	CDC, 2017 ⁷³
Tango	www.tango.me Apple App Store and Google Play Store	USA AMR	Free to download, in-app purchases for extra features	CDC, 2017 ⁷³
Adobe Connect	www.adobe.com or via Apple App Store and Google Play Store	USA AMR	Free	Donahue, 2021 ³¹
ViaTV units *earliest example of VOT amongst included studies	Not commercially available; developed by the research team to prove earliest VOT concept	USA AMR	Not reported	DeMaio, 2001 ¹⁷
Youping Technology Co.	Not commercially available, developed for study trial	China WPR	Not reported	Guo, 2020 ³⁶
MIST (Mobile Interactive Supervised Therapy)	Not commercially available, developed for study trial	Singapore WPR	Not reported	Molton, 2016 ⁴⁹
I Like VST by TBVOT.MD	www.tbvot.md and Google Play Store	Moldova EUR	Free	WHO, 2020 ⁷
Lekar	Via Apple App Store and Google Play, owned by ABM Cloud	Ukraine EUR	Free	WHO, 2020 ⁷

HealthFox	www.healthfox.fi or via Apple App Store and Google Play Store	Finland EUR	Free	Rajalahti, 2022 ¹⁸
AppVST	Not commercially available, developed by TUH technology department	Finland EUR	Not reported	Rajalahti, 2022 ¹⁸
NIMCURE	Not commercially available; developed by Co-Creation Hub and Nigerian Institute of Medical Research (NIMR)	Nigeria AFR	Not reported	Samuel, 2021 ⁵⁷
Adhere 2TX-TB	Not reported	Not reported	Not reported	WHO, 2020 ⁷

*Applications developed or evaluated prior to the year 2000 that did not meet the study eligibility criteria for inclusion may exist. All websites accessed January 2024.

Supplementary Table 2: Crow Critical Appraisal Tool (CCAT)

Breakdown of Scores - Primary & Review Studies (n = 60)

Scoring rubric: 0 – 49% = low quality; 50-74% = moderate quality; 75-100% = high quality Adapted from Nuttall et al. 2022⁶

Study ref.	Preliminaries	Intro	Design	Sampling	Data collection	Ethical matters	Results	Discussion	Total CCAT Score (n/40) / Grading
Abbara et al. 2020 ⁶³	4	5	4	4	3	5	4	4	33 (82.5%) High Quality
Aisyah et al. 2020 ¹⁹	4	5	4	3	4	5	4	4	33 (82.5%) High Quality
Alipanah et al. 2018 ⁴	4	4	5	5	5	5	5	4	37 (92.5%) High Quality
Asay et al. 2020 ¹¹	4	5	4	4	3	4	4	5	33 (82.5%) High Quality
Bachina et al. 2022 ²¹	4	5	4	4	3	4	4	4	32 (80%) High Quality
Bendiksen et al. 2020 ²²	4	4	4	3	4	3	4	4	30 (75%) High Quality
Bommakan ti et al. 2020 ²³	4	4	4	3	5	5	4	4	33 (82.5%) High Quality
Brett et al. 2020 ⁶⁴	4	5	5	4	3	4	5	5	35 (87.5%) High Quality
Buchman et al. 2017 ²⁴	4	5	3	5	3	2	3	4	29 (73%) Moderate Quality
Burzynski et al. 2022 ²⁵	4	5	5	5	3	5	4	5	36 (90%) High Quality
Casalme et al. 2022 ²⁶	4	4	4	3	1	5	4	4	29 (72.5%) Moderate Quality
Chakaya et al. 2021 ⁶⁵	4	4	4	5	5	5	4	5	36 (90%) High Quality
Chen et al. 2020 ²⁷	4	4	4	3	3	5	4	4	31 (77.5%) High Quality

Chuck et al. 2016 ²⁸	4	5	4	3	3	4	4	4	31 (77.5%) High Quality
DeMaio et al. 2001 ¹⁷	4	4	4	3	3	3	4	4	29 (72.5%) Moderate Quality
Do et al. 2019 ²⁹	4	5	3	4	4	5	4	4	33 (82.5%) High Quality
Doltu et al. 2021 ³⁰	4	5	4	3	4	5	4	4	33 (82.5%) High Quality
Donahue et al. 2021 ³¹	4	4	4	3	4	3	4	4	30 (75%) High Quality
Gassanov et al. 2013 ³⁵	3	3	3	3	3	0	3	3	21 (52.5%) Moderate Quality
Garfein et al. 2015 ³²	4	4	5	4	4	5	4	5	35 (87.5%) High Quality
Garfein et al. 2018 ³³	4	4	5	3	5	4	4	4	33 (82.5%) High Quality
Garfein et al. 2019 ⁵	4	4	3	0	4	5	4	4	28 (70%) Moderate Quality
Garfein et al. 2020 ³⁴	4	5	4	3	3	5	3	4	31 (77.5%) High Quality
Guo et al. 2020a ³⁶	4	4	4	3	4	5	4	4	32 (80%) High Quality
Guo et al. 2020b ³⁷	4	4	3	3	4	3	4	4	29 (72.5%) Moderate Quality
Hira et al. 2022 ³⁸	4	5	4	4	4	0	4	3	28 (70%) Moderate Quality
Hoffman et al. 2010 ³⁹	4	5	4	3	3	4	3	3	29 (72.5%) Moderate Quality
Holzman et al. 2018 ⁴⁰	4	5	4	5	3	5	4	4	34 (85%) High Quality
Holzman et al. 2019 ⁴¹	4	5	4	4	3	4	5	5	34 (85%) High Quality
Keutzer et al. 2020 ⁶⁶	4	4	3	4	3	3	4	4	29 (72.5%) Moderate Quality
Krueger et al. 2010 ⁴³	4	5	4	3	3	3	3	3	28 (70%) Moderate Quality

Kumar et al. 2019 ⁴²	4	5	4	4	4	5	4	4	34 (85%) High Quality
Lam et al. 2018 ⁴⁴	4	5	4	3	5	4	4	4	33 (82.5%) High Quality
Lam et al. 2019 ⁴⁵	4	5	4	3	3	5	4	4	32 (80%) High Quality
Lee et al. 2020 ⁶⁷	4	5	4	5	3	3	4	5	33 (82.5%) High Quality
Lippincott et al. 2022 ⁴⁶	4	5	4	4	3	5	4	5	34 (85%) High Quality
Macaraig et al. 2018 ⁴⁷	4	5	4	5	4	4	4	4	34 (85%) High Quality
Mirsaeidi et al. 2015 ⁴⁸	3	4	3	3	3	4	3	3	26 (65%) Moderate Quality
Molton et al. 2016 ⁴⁹	4	5	4	5	4	5	4	5	36 (90%) High Quality
Nguyen et al. 2017 ⁵⁰	4	5	4	5	3	4	4	3	32 (80%) High Quality
Ngwatu et al. 2018 ⁶⁸	4	5	5	5	5	4	4	4	36 (90%) High Quality
Nsengiyumva et al. 2018 ⁵¹	4	4	4	3	3	4	4	4	30 (75%) High Quality
Perry et al. 2021 ⁵²	4	5	3	3	3	4	4	4	30 (75%) High Quality
Rabinovich et al. 2020 ⁵³	4	5	4	3	5	4	4	4	33 (82.5%) High Quality
Rajalahti et al. 2022 ¹⁸	4	4	4	3	4	5	4	4	32 (80%) High Quality
Rao et al. 2022 ⁵⁴	4	5	4	3	3	4	4	4	31 (77.5%) High Quality
Ravenscroft et al. 2020 ⁵⁵	4	5	4	5	3	5	4	4	34 (85%) High Quality
Ridho et al. 2022 ⁶⁹	5	4	4	5	5	4	5	5	37 (92.5%) High Quality


Ritter et al. 2017 ⁵⁶	3	5	4	5	5	2	4	5	33 (83%) High Quality
Samuel et al. 2021 ⁵⁷	4	4	4	3	3	4	4	3	29 (72.5%) Moderate Quality
Sekandi et al. 2020 ⁵⁸	4	5	4	3	4	5	4	5	34 (85%) High Quality
Sekandi et al. 2021 ⁵⁹	4	5	4	3	5	5	4	5	35 (87.5%) High Quality
Sinkou et al. 2017 ⁶⁰	4	5	4	3	3	5	4	4	32 (90%) High Quality
Stoner et al. 2022 ⁸	5	5	4	5	5	5	5	5	39 (97.5 %) High Quality
Story et al. 2019 ⁹	4	5	4	5	4	4	5	5	36 (90%) High Quality
Subbaraman et al. 2018 ⁷⁰	4	5	4	4	4	4	4	4	33 (82.5%) High Quality
Truong et al. 2022 ⁷¹	5	5	5	5	4	3	4	4	35 (87.5%) High Quality
Wade et al. 2012 ⁶¹	5	5	5	4	5	5	4	5	38 (95%) High Quality
Zúñiga et al. 2016 ⁶²	4	4	4	3	5	4	4	4	32 (80%) High Quality

Supplementary Figure 1: Geographical locations of study sites of VOT studies (n=55)

Global Representation of VOT Study Locality

Adapted from Table 3

< 2 2-4 4-6 6-8 8-10 10-12 12-14 14-16 16-18 18-20 20-22 22-24 ≥ 24



Map: ADT & EM • Source: ADT, EM, KP • Created with Datawrapper

**Legend: Due to country size or geopolitical recognition, Moldova (n=3, lower-middle income), Singapore (n=1, high income) and Taiwan (n=1, high income), are not visualised in the above map by the software used (Datawrapper, 2023). None of these three countries are high TB, HIV/TB, or MDR-TB burden countries.*