

1 Supplementary tables

2 **Table S1. Overview of the MRI scanners and parameters employed to acquire the**
 3 **T1-weighted images analyzed in the study.**

FS (T)	Manuf.	Model	TR (sec)	TE (sec)	TI (sec)	FA (°)	Native voxel vol. (mm ³)	OVERALL (N=20845) N (%)	ADNI (N=10977) N (%)	NACC (N=7819) N (%)	OASIS (N=2049) N (%)
1.5	GE	GENESIS SIGNA	0.009 to 0.035	0.002 to 0.008	0.45 to 1	8 to 60	1.2	1641 (7.9)	271 (2.5)	1370 (17.5)	0
1.5	GE	SIGNA	0.018	0.002 to 0.003	NA	25	1.8	50 (0.2)	0	50 (0.6)	0
1.5	GE	SIGNA EXCITE	0.008 to 0.029	0.002 to 0.004	0.5 to 1	8 to 25	1.1	999 (4.8)	909 (8.3)	90 (1.2)	0
1.5	GE	SIGNA HDx	0.009 to 0.011	0.004	0.5 to 1	8 to 10	1.1	587 (2.8)	425 (3.9)	162 (2.1)	0
1.5	GE	Signa HDxt	0.009 to 0.02	0.004 to 0.007	0.45 to 1	8 to 30	1.1	572 (2.7)	303 (2.8)	269 (3.4)	0
1.5	Hitachi	OASIS	NA	NA	NA	NA	0.2	1 (0.0)	0	1 (0.0)	0
1.5	Philips	Achieva	0.009	0.004	1	8	1.1	70 (0.3)	70 (0.6)	0	0
1.5	Philips	Eclipse 1.5T	0.009	0.002	NA	15	1.5	74 (0.4)	0	74 (0.9)	0
1.5	Philips	Gyrosan Intera	0.008 to 0.009	0.004	0.761 to 1	8	1.1	13 (0.1)	12 (0.1)	1 (0.0)	0
1.5	Philips	Gyrosan NT	0.009	0.004	1	8	1	19 (0.1)	19 (0.2)	0	0
1.5	Philips	Intera	0.008 to 0.01	0.003 to 0.004	1 to 1.076	8	1.1	359 (1.7)	352 (3.2)	7 (0.1)	0
1.5	Philips	Intera Achieva	0.009	0.004	1	8	1.1	6 (0.0)	6 (0.1)	0	0
1.5	Siemens	Aera	NA	NA	0.8	NA	1	7 (0.0)	0	7 (0.1)	0
1.5	Siemens	Avanto	1.7 to 2.4	0.002 to 0.004	1 to 1.1	8 to 15	1.9	452 (2.2)	443 (4.0)	7 (0.1)	2 (0.1)
1.5	Siemens	Espreo	0.011 to 2.4	0.003 to 0.006	1 to 1.1	8 to 25	1.8	36 (0.2)	24 (0.2)	12 (0.2)	0
1.5	Siemens	Magnetom VISION	NA	NA	0.3	NA	1.8	2 (0.0)	0	2 (0.0)	0
1.5	Siemens	Magnetom ESSENZA	1.85	0.003	1.1	9	1	1 (0.0)	0	1 (0.0)	0
1.5	Siemens	NUMARIS/4	2.4	0.004	1	8	1.9	2 (0.0)	2 (0.0)	0	0
1.5	Siemens	Sonata	1.9 to 3	0.003 to 0.005	0.93 to 1.1	8 to 20	1.8	498 (2.4)	376 (3.4)	88 (1.1)	34 (1.7)
1.5	Siemens	Sonata Vision	2.4 to 3	0.004	1	8	1.9	40 (0.2)	40 (0.4)	0	0
1.5	Siemens	Symphony	2 to 3	0.003 to 0.004	1 to 1.1	8 to 15	1.9	681 (3.3)	594 (5.4)	87 (1.1)	0
1.5	Siemens	Symphony Tim	2.4 to 3	0.002 to 0.004	1	8	1.9	126 (0.6)	124 (1.1)	2 (0.0)	0
1.5	Siemens	Vision	9.7	4	NA	10	1.3	172 (0.8)	0	0	172 (8.4)
3	GE	Discovery MR750	0.006 to 0.009	0.002 to 0.003	0.4 to 1.06	8 to 12	1.1	2990 (14.3)	958 (8.7)	2032 (26.0)	0
3	GE	Discovery MR750w	0.008	0.003	0.4	11	1.1	132 (0.6)	132 (1.2)	0	0
3	GE	GENESIS SIGNA	0.007	0.003	0.9	8	1.2	18 (0.1)	18 (0.2)	0	0
3	GE	SIGNA EXCITE	0.007 to 0.008	0.003	0.45 to 0.9	8 to 15	1.2	50 (0.2)	44 (0.4)	6 (0.1)	0
3	GE	SIGNA HDx	0.007	0.003	0.4 to 0.9	8 to 11	1.3	53 (0.3)	53 (0.5)	0	0

3	GE	Signa HDxt	0.005 to 0.02	0.002 to 0.004	0.4 to 0.9	8 to 27	1.2	658 (3.2)	427 (3.9)	231 (3.0)	0
3	GE	Signa MR360	2.1	0.003	0.9	8	1	1 (0.0)	0	1 (0.0)	0
3	GE	SIGNA PET/MR	0.008	0.003	0.4	11	1.3	77 (0.4)	0	77 (1.0)	0
3	GE	SIGNA Premier	0.007	0.003	0.4 to 0.45	11 to 12	1	89 (0.4)	56 (0.5)	33 (0.4)	0
3	GE	SIGNA UHP	0.007 to 2.077	0.003	0.4 to 0.9	8 to 11	1	10 (0.0)	7 (0.1)	3 (0.0)	0
3	Philips	Achieva	0.006 to 0.007	0.003	0.805 to 0.9	8 to 9	1.2	889 (4.3)	597 (5.4)	292 (3.7)	0
3	Philips	Achieva dStream	0.006 to 0.007	0.003	NA	8 to 9	1	186 (0.9)	138 (1.3)	48 (0.6)	0
3	Philips	GEMINI	0.007	0.003	NA	9	1.2	44 (0.2)	35 (0.3)	9 (0.1)	0
3	Philips	Ingenia	0.007	0.003	0.9	9	1.1	209 (1.0)	209 (1.9)	0	0
3	Philips	Ingenia Elition X	0.007	0.003	NA	9	1	14 (0.1)	14 (0.1)	0	0
3	Philips	Ingenuity	0.007	0.003	NA	9	1.3	19 (0.1)	16 (0.1)	3 (0.0)	0
3	Philips	Intera	0.007	0.003	0.845 to 0.869	8 to 9	1.2	273 (1.3)	273 (2.5)	0	0
3	Philips	Intera Achieva	0.007	0.003	0.855	8	1.2	1 (0.0)	1 (0.0)	0	0
3	Siemens	Allegra	2.3 to 2.5	0.003 to 0.004	0.9 to 1.1	8 to 9	1.1	136 (0.7)	108 (1.0)	28 (0.4)	0
3	Siemens	Biograph mMR	2.3 to 2.4	0.002 to 0.003	0.9 to 1	8 to 9	1.3	584 (2.8)	26 (0.2)	5 (0.1)	553 (27.0)
3	Siemens	Magnetom Prisma Fit	2.3	0.003	0.9	9	1	12 (0.1)	12 (0.1)	0	0
3	Siemens	Magnetom Vida	2.3 to 2.4	0.003	0.9	8 to 9	1.2	219 (1.1)	3 (0.0)	1 (0.0)	215 (10.5)
3	Siemens	Prisma	2.3 to 2.4	0.002 to 0.003	0.9 to 1.06	8 to 9	1	957 (4.6)	255 (2.3)	702 (9.0)	0
3	Siemens	Prisma fit	1.8 to 2.4	0.002 to 0.005	0.9 to 1	8 to 10	1	810 (3.9)	663 (6.0)	146 (1.9)	1 (0.0)
3	Siemens	Skyra	1.8 to 2.3	0.002 to 0.003	0.649 to 0.962	8 to 10	1.1	1696 (8.1)	489 (4.5)	1207 (15.4)	0
3	Siemens	Skyra fit	2.3	0.003	0.9	9	1	26 (0.1)	26 (0.2)	0	0
3	Siemens	Trio	1.62 to 2.5	0.002 to 0.004	0.9 to 1.1	7 to 15	1.2	144 (0.7)	121 (1.1)	23 (0.3)	0
3	Siemens	TrioTim	1.31 to 3.2	0.002 to 0.455	0.9 to 1.2	7 to 120	1.1	3098 (14.9)	1547 (14.1)	479 (6.1)	1072 (52.3)
3	Siemens	Verio	1.7 to 2.3	0.002 to 0.004	0.9	8 to 9	1.2	805 (3.9)	767 (7.0)	38 (0.5)	0
Not available / Unknown			0.007 to 2.3	0.002 to 0.005	0.4 to 0.9	8 to 60	1	237 (1.1)	12 (0.1)	225 (2.9)	0

5 **Table S2. Multi-variate associations of PVS and WML markers with the**
6 **demographic and clinical covariates included in all the models of this study.**
7

Variable	Estimate	Standard error	T value	Adjusted P value
WM-PVS count				
Age (years)	-4.31	0.14	-30.43	<0.001
Female sex	-24.35	3.12	-7.80	<0.001
Education (years)	-1.50	0.44	-3.40	0.001
Race (compared to White)				
Black	27.18	4.13	6.58	<0.001
Asian	28.27	14.07	2.01	0.089
American Indians	9.36	9.53	0.98	0.652
More than one reported	1.03	20.64	0.05	1
CDR global score	-62.10	3.20	-19.42	<0.001
Body Mass Index (kg/m ²)	1.97	0.26	7.63	<0.001
Dyslipidemia	2.02	2.64	0.77	0.887
Hypertension	-6.43	2.78	-2.32	0.041
Diabetes	-11.14	4.04	-2.76	0.012
History of cardio-/cerebro-vascular disease	-1.68	2.88	-0.58	1
Family history of dementia	-4.13	2.58	-1.60	0.219
BG-PVS count				
Age (years)	-1.43	0.05	-30.68	<0.001
Female sex	-5.89	1.03	-5.72	<0.001
Education (years)	0.43	0.15	2.94	0.006
Race (compared to White)				
Black	11.96	1.36	8.77	<0.001
Asian	11.30	4.64	2.43	0.03
American Indians	7.15	3.15	2.27	0.046
More than one reported	15.82	6.81	2.32	0.04
CDR global score	-16.96	1.06	-16.07	<0.001
Body Mass Index (kg/m ²)	0.91	0.09	10.68	<0.001
Dyslipidemia	1.14	0.87	1.31	0.379
Hypertension	-4.57	0.92	-4.99	<0.001
Diabetes	-3.37	1.33	-2.53	0.023
History of cardio-/cerebro-vascular disease	-1.89	0.95	-1.99	0.093
Family history of dementia	0.54	0.85	0.63	1
WM-PVS diameter				
Age (years)	0.03	0.00	17.81	<0.001
Female sex	0.18	0.03	5.39	<0.001
Education (years)	0.00	0.00	-0.31	1
Race (compared to White)				
Black	-0.15	0.04	-3.28	0.002

Asian	-0.09	0.15	-0.56	1
American Indians	-0.42	0.10	-4.14	<0.001
More than one reported	-0.61	0.22	-2.72	0.013
CDR global score	0.19	0.03	5.44	<0.001
Body Mass Index (kg/m ²)	-0.02	0.00	-5.63	<0.001
Dyslipidemia	0.03	0.03	0.98	0.658
Hypertension	0.21	0.03	6.86	<0.001
Diabetes	-0.16	0.04	-3.76	<0.001
History of cardio-/cerebro-vascular disease	-0.04	0.03	-1.23	0.436
Family history of dementia	0.01	0.03	0.25	1
BG-PVS diameter				
Age (years)	0.02	0.00	19.30	<0.001
Female sex	0.09	0.02	4.13	<0.001
Education (years)	-0.01	0.00	-2.82	0.01
Race (compared to White)				
Black	-0.19	0.03	-6.56	<0.001
Asian	-0.20	0.10	-2.03	0.085
American Indians	-0.15	0.07	-2.26	0.047
More than one reported	-0.38	0.14	-2.64	0.017
CDR global score	0.18	0.02	8.03	<0.001
Body Mass Index (kg/m ²)	-0.02	0.00	-9.19	<0.001
Dyslipidemia	-0.03	0.02	-1.92	0.111
Hypertension	0.12	0.02	6.39	<0.001
Diabetes	-0.05	0.03	-1.77	0.152
History of cardio-/cerebro-vascular disease	0.05	0.02	2.34	0.039
Family history of dementia	-0.02	0.02	-1.16	0.494
P-WML log volume				
Age (years)	0.15	0.00	47.25	<0.001
Female sex	0.64	0.07	9.32	<0.001
Education (years)	-0.01	0.01	-1.50	0.267
Race (compared to White)				
Black	0.03	0.09	0.31	1
Asian	-0.56	0.31	-1.82	0.138
American Indians	0.15	0.21	0.72	0.943
More than one reported	-0.25	0.45	-0.55	1
CDR global score	1.33	0.07	19.06	<0.001
Body Mass Index (kg/m ²)	-0.03	0.01	-5.98	<0.001
Dyslipidemia	0.00	0.06	0.05	1
Hypertension	0.45	0.06	7.36	<0.001
Diabetes	0.13	0.09	1.44	0.298
History of cardio-/cerebro-vascular disease	0.08	0.06	1.24	0.433
Family history of dementia	-0.10	0.06	-1.72	0.17

D-WML log volume				
Age (years)	0.02	0.00	10.38	<0.001
Female sex	0.28	0.05	5.51	<0.001
Education (years)	-0.01	0.01	-1.18	0.479
Race (compared to White)				
Black	0.32	0.07	4.83	<0.001
Asian	-0.21	0.23	-0.94	0.698
American Indians	-0.18	0.15	-1.15	0.504
More than one reported	-0.66	0.33	-1.97	0.097
CDR global score	-0.25	0.05	-4.83	<0.001
Body Mass Index (kg/m ²)	-0.01	0.00	-2.83	0.009
Dyslipidemia	-0.03	0.04	-0.65	1
Hypertension	0.21	0.04	4.73	<0.001
Diabetes	-0.09	0.07	-1.42	0.31
History of cardio-/cerebro-vascular disease	-0.03	0.05	-0.58	1
Family history of dementia	0.00	0.04	-0.09	1

8
9 This analysis was run in the entire study population (N=10,004), including non-
10 demented and demented subjects.

11 **Table S3. Multi-variate associations of PVS and WML markers with the clinical**
 12 **covariates assessed in the sensitivity analyses.**
 13

Variable	Total number of subjects	Estimate	Standard error	T value	Adjusted P value
WM-PVS count					
APOE (ref. ε3ε3)	8599				
ε2ε2		-40.02	23.01	-1.74	0.164
ε2ε3		1.48	4.79	0.31	1
ε2ε4		3.61	8.66	0.42	1
ε3ε4		0.55	3.07	0.18	1
ε4ε4		-8.51	5.42	-1.57	0.233
Baseline MMSE	7036	6.43	0.46	14.11	<0.001
Tobacco smoking	9267	-5.29	2.74	-1.93	0.106
A-beta positivity	3906	-8.77	4.06	-2.16	0.062
Tau positivity	2466	8.72	4.92	1.77	0.153
BG-PVS count					
APOE (ref. ε3ε3)	8599				
ε2ε2		-7.57	7.42	-1.02	0.616
ε2ε3		1.65	1.55	1.07	0.573
ε2ε4		0.49	2.79	0.18	1
ε3ε4		-1.19	0.99	-1.20	0.461
ε4ε4		-5.07	1.75	-2.90	0.007
Baseline MMSE	7036	1.56	0.14	11.16	<0.001
Tobacco smoking	9267	-7.24	0.90	-8.03	<0.001
A-beta positivity	3906	-3.92	1.45	-2.71	0.014
Tau positivity	2466	-1.62	1.69	-0.96	0.679
WM-PVS diameter					
APOE (ref. ε3ε3)	8599				
ε2ε2		0.21	0.25	0.83	0.81
ε2ε3		-0.09	0.05	-1.72	0.17
ε2ε4		0.04	0.09	0.44	1
ε3ε4		0.02	0.03	0.61	1
ε4ε4		0.16	0.06	2.73	0.013
Baseline MMSE	7036	-0.02	0.01	-2.94	0.007
Tobacco smoking	9267	0.16	0.03	5.59	<0.001
A-beta positivity	3906	0.15	0.05	2.95	0.006
Tau positivity	2466	0.17	0.06	2.80	0.01
BG-PVS diameter					
APOE (ref. ε3ε3)	8599				
ε2ε2		0.05	0.16	0.33	1
ε2ε3		-0.02	0.03	-0.72	0.943
ε2ε4		-0.04	0.06	-0.73	0.934
ε3ε4		0.02	0.02	0.79	0.856
ε4ε4		0.09	0.04	2.31	0.041

Baseline MMSE	7036	-0.01	0.00	-3.63	<0.001
Tobacco smoking	9267	0.12	0.02	6.08	<0.001
A-beta positivity	3906	0.07	0.03	2.05	0.081
Tau positivity	2466	0.10	0.04	2.39	0.034
P-WML log volume					
APOE (ref. ε3ε3)	8599				
ε2ε2		0.51	0.50	1.03	0.609
ε2ε3		-0.01	0.10	-0.14	1
ε2ε4		0.19	0.19	1.01	0.624
ε3ε4		-0.02	0.07	-0.30	1
ε4ε4		0.26	0.12	2.22	0.053
Baseline MMSE	7036	-0.14	0.01	-14.05	<0.001
Tobacco smoking	9267	0.19	0.06	3.14	0.003
A-beta positivity	3906	0.40	0.09	4.26	<0.001
Tau positivity	2466	-0.22	0.11	-1.91	0.113
D-WML log volume					
APOE (ref. ε3ε3)	8599				
ε2ε2		0.07	0.37	0.19	1
ε2ε3		-0.09	0.08	-1.11	0.536
ε2ε4		0.09	0.14	0.62	1
ε3ε4		-0.06	0.05	-1.21	0.449
ε4ε4		0.24	0.09	2.70	0.014
Baseline MMSE	7036	0.01	0.01	1.77	0.153
Tobacco smoking	9267	-0.01	0.04	-0.16	1
A-beta positivity	3906	0.17	0.07	2.44	0.029
Tau positivity	2466	0.10	0.09	1.22	0.444

14
15 These general linear models included the same covariates as in the primary analysis
16 (Table S2), and the indicated variable was included as additional covariate. For
17 “MMSE”, the variable “MMSE score at the baseline” was used in place of the CDR
18 score.

19
20

21 **Table S4. Estimated marginal means for PVS and WML markers.**
 22

Marker	Estimated marginal mean \pm standard error		Adj. P value
	Non-demented	Demented	
WM-PVS count	445 \pm 6	371 \pm 7	<0.001
BG-PVS count	144 \pm 2	123 \pm 2	<0.001
WM-PVS diameter (mm)	1.96 \pm 0.01	1.99 \pm 0.01	<0.001
BG-PVS diameter (mm)	1.57 \pm 0	1.60 \pm 0	<0.001
P-WML log volume	4.48 \pm 0.12	5.89 \pm 0.15	<0.001
D-WML log volume	2.56 \pm 0.09	2.41 \pm 0.11	0.051

23
 24 Marginal means were estimated controlling for the same covariates as in the primary
 25 analysis (Table S2).
 26

27 **Table S5. Sensitivity analysis on the ANCOVA models.**

Covariate	N. of demented patients / total	Marker	Estimated marginal mean ± standard error		Adj. P value
			Non-demented	Demented	
Artifacts	1123 / 9955	WM-PVS count	446 ± 6	372 ± 7	<0.001
		BG-PVS count	144 ± 2	123 ± 2	<0.001
		WM-PVS diameter	1.96 ± 0.01	1.99 ± 0.01	<0.001
		BG-PVS diameter	1.57 ± 0	1.6 ± 0	<0.001
		P-WML log volume	4.47 ± 0.12	5.89 ± 0.15	<0.001
		D-WML log volume	2.56 ± 0.09	2.43 ± 0.11	0.1
APOE	974 / 8599	WM-PVS count	437 ± 7	362 ± 8	<0.001
		BG-PVS count	139 ± 2	119 ± 3	<0.001
		WM-PVS diameter	1.97 ± 0.01	2 ± 0.01	<0.001
		BG-PVS diameter	1.58 ± 0.01	1.6 ± 0.01	<0.001
		P-WML log volume	4.67 ± 0.16	6.05 ± 0.19	<0.001
		D-WML log volume	2.64 ± 0.12	2.49 ± 0.14	0.08
Baseline MMSE	1255 / 7036	WM-PVS count	450 ± 7	403 ± 8	<0.001
		BG-PVS count	143 ± 2	131 ± 2	<0.001
		WM-PVS diameter	1.95 ± 0.01	1.96 ± 0.01	0.026
		BG-PVS diameter	1.57 ± 0	1.58 ± 0.01	0.004
		P-WML log volume	4.47 ± 0.14	5.37 ± 0.16	<0.001
		D-WML log volume	2.52 ± 0.11	2.41 ± 0.12	0.205
History of tobacco smoking	1053 / 9267	WM-PVS count	447 ± 6	372 ± 7	<0.001
		BG-PVS count	1.96 ± 0.01	1.99 ± 0.01	<0.001
		WM-PVS diameter	143 ± 2	122 ± 2	<0.001
		BG-PVS diameter	1.57 ± 0	1.6 ± 0	<0.001
		P-WML log volume	4.48 ± 0.13	5.89 ± 0.15	<0.001
		D-WML log volume	2.55 ± 0.09	2.39 ± 0.11	0.037
A-beta positivity	327 / 3906	WM-PVS count	445 ± 10	391 ± 12	<0.001
		BG-PVS count	142 ± 4	128 ± 4	<0.001
		WM-PVS diameter	1.97 ± 0.01	1.98 ± 0.01	0.306
		BG-PVS diameter	1.58 ± 0.01	1.58 ± 0.01	0.553
		P-WML log volume	4.94 ± 0.23	6.11 ± 0.27	<0.001
		D-WML log volume	2.55 ± 0.17	2.41 ± 0.2	0.43
Tau positivity	217 / 2466	WM-PVS count	453 ± 12	380 ± 14	<0.001
		BG-PVS count	137 ± 4	121 ± 5	<0.001
		WM-PVS diameter	1.98 ± 0.01	2.01 ± 0.02	0.01
		BG-PVS diameter	1.59 ± 0.01	1.6 ± 0.01	0.169
		P-WML log volume	4.84 ± 0.27	6.3 ± 0.32	<0.001
		D-WML log volume	2.63 ± 0.2	2.46 ± 0.24	0.487

28

29 These models were adjusted for the same covariates as in the primary analysis (Table

30 S2). For “Artifacts”, cases where the estimate of the vascular marker may have been

31 affected by MRI artifacts or the MRI data were acquired after the injection of a contrast
32 agent were excluded from the model. For “MMSE”, the definition of dementia was based
33 on the MMSE score rather than CDR. For all the other models, the indicated variable
34 was included as additional covariate in the model.
35

36
37

Table S6. Baseline characteristics of non-demented individuals included in the analysis of the hazard ratio for dementia.

Characteristic	Overall (N=7518)	ADNI (N=1931)	NACC (N=4433)	OASIS (N=1154)
Age at baseline – yr	71.0 ± 9.5	73.1 ± 7.1	70.3 ± 10.3	69.9 ± 9.1
Female – %	56.4	46.6	61	55.5
Education – yr	15.9 ± 2.9	16.1 ± 2.7	15.8 ± 3.0	15.9 ± 2.7
Race – %				
White	86.1	91.6	84.1	84.6
Black	11.1	5	12.9	14.7
Asian	0.8	0.2	1.2	0
American Indians	1.7	2	1.8	0.4
More than one reported	0.4	1.2	0	0.3
Body Mass Index – kg/m ²	27.2 ± 5.2	27.1 ± 4.9	27.0 ± 5.2	27.9 ± 5.4
Dyslipidemia – %	55.4	53.6	58.1	48.1
Hypertension – %	50.2	48.8	51.2	48.6
Diabetes – %	12.5	10.3	14	10.8
History of cardio- /cerebro-vascular disease – %	28.1	36.5	25.2	24.9
Family history of dementia – %	53.6	53.6	52.2	59.5
History of tobacco smoking* – %	26.7	50.5	44.8	37.1
APOE* – %				
ε2ε2	0.3	0.1	0.4	0.6
ε2ε3	9.6	7.8	10.3	10.6
ε2ε4	2.5	2	2.5	3.7
ε3ε3	50.4	48	52.1	48.7
ε3ε4	30.8	33.1	29.6	31
ε4ε4	6.3	9	5.3	5.3
A-beta positivity* – %	49.5	63.9	40.5	34
Tau positivity* – %	54.9	65.6	39.2	43
Median follow-up – yr	4.1 (2.3 to 7.1)	4.0 (2.1 to 6.8)	4.1 (2.3 to 6.6)	6.1 (3.2 to 10.1)
Number of conversions to dementia	1493	469	803	221
WM-PVS count	435 (346 to 537)	382 (308 to 480)	459 (369 to 563)	434 (350 to 525)

BG-PVS count	135 (107 to 167)	105 (88 to 132)	147 (122 to 177)	133 (113 to 158)
WM-PVS diameter – mm	1.96 (1.88 to 2.07)	2.07 (1.96 to 2.18)	1.93 (1.86 to 2.02)	1.96 (1.88 to 2.05)
BG-PVS diameter – mm	1.57 (1.52 to 1.64)	1.65 (1.57 to 1.71)	1.54 (1.50 to 1.59)	1.57 (1.53 to 1.62)
P-WML volume – mm ³	254.50 (0 to 1452)	581 (46 to 1920.50)	143 (0 to 1189)	223 (0 to 1521)
D-WML volume – mm ³	17 (1 to 74)	37 (7 to 107)	8 (0 to 55)	25 (4 to 94.75)

38 Plus-minus values are means \pm standard deviations, entries with parentheses are
39 medians (interquartile range). Data from three studies — the Alzheimer’s Disease
40 Neuroimaging Initiative (ADNI), the National Alzheimer’s Coordinating Center (NACC),
41 and the Open Access Series of Imaging Studies (OASIS) — are shown. *Total number
42 of subjects with available history of tobacco smoking, Apolipoprotein E (APOE)
43 genotype, amyloid-beta status and tau status were 3918, 6968, 3774, and 2434,
44 respectively.
45

46 **Table S7. Sensitivity analysis on the HR for dementia**

Characteristic and overall prevalence (%) or mean \pm standard deviation		N. of conversions to dementia / total	HR [95% CI]				
			WM-PVS count	BG-PVS count	WM-PVS diameter	BG-PVS diameter	P-WML
Artifacts – %	0.4	1487 / 7490	0.80 [0.76 0.84]	0.91 [0.89 0.93]	1.10 [1.05 1.15]	1.21 [1.13 1.30]	1.14 [1.11 1.16]
APOE – %		1418 / 6914	0.79 [0.75 0.83]	0.91 [0.89 0.93]	1.09 [1.04 1.14]	1.19 [1.11 1.27]	1.14 [1.11 1.17]
ε2ε2	0.3						
ε2ε3	9.6						
ε2ε4	2.5						
ε3ε3	50.4						
ε3ε4	30.8						
ε4ε4	6.3						
Baseline MMSE	28.5 \pm 1.5	1259 / 5311	0.86 [0.81 0.91]	0.93 [0.91 0.95]	1.09 [1.04 1.14]	1.13 [1.05 1.22]	1.10 [1.07 1.14]
History of tobacco smoking – %	37.1	1056 / 3918	0.80 [0.76 0.85]	0.92 [0.9 0.94]	1.09 [1.04 1.14]	1.17 [1.09 1.26]	1.12 [1.09 1.15]
A-beta positivity – %	49.7	697 / 3774	0.80 [0.74 0.86]	0.91 [0.89 0.94]	1.05 [0.99 1.12]	1.15 [1.05 1.27]	1.12 [1.08 1.16]
Tau positivity – %	54.9	465 / 2434	0.77 [0.70 0.85]	0.91 [0.88 0.94]	1.08 [1.01 1.17]	1.18 [1.06 1.32]	1.13 [1.08 1.18]

47 These models were adjusted for the same covariates as in the primary analysis (Fig. 1).
 48 For “Artifacts”, cases where the estimate of the vascular marker may have been
 49 affected by MRI artifacts or the MRI data were acquired after the injection of a contrast
 50 agent were excluded from the model. For “MMSE”, the definition of dementia was based
 51 on the MMSE score rather than CDR, and the corresponding MMSE at the baseline was
 52 used as covariate in place of CDR score at the baseline. For all the other models, the
 53 indicated variable was included as additional covariate in the model. Hazard ratios (HR)
 54 for WM-PVS and BG-PVS count are per 100 and 10 units increase, respectively, for
 55 WM-PVS and BG-PVS diameter are per 0.1 mm units increase, and for P-WML volume
 56 are per 1 logarithmic unit increase. See also Fig. S7 for spline plots.
 57

58
59
60

Table S8. Baseline characteristics of non-demented individuals included in the analysis of the linear mixed-effects model for brain atrophy and for the longitudinal trajectories of PVS and WML markers.

Characteristic	Overall (N=3389)	ADNI (N=1833)	NACC (N=1176)	OASIS (N=380)
Age at baseline – yr	70.7 ± 9.8	73.3 ± 7.0	67.8 ± 12.1	67.5 ± 9.6
Female – %	53.3	46.0	63.4	56.8
Education – yr	15.9 ± 2.9	16.1 ± 2.7	15.7 ± 3.1	15.9 ± 2.6
Race – %				
White	89.2	92.4	84.6	87.8
Black	7.9	4.4	12.4	11.4
Asian	0.5	0.3	0.9	0
American Indians	1.7	1.7	2	0.5
More than one reported	0.7	1.2	0	0.3
Body Mass Index – kg/m ²	27.0 ± 5.0	27.0 ± 4.8	27.3 ± 5.2	27.9 ± 5.0
Dyslipidemia – %	51.5	53.4	50.8	44.3
Hypertension – %	46.4	48.9	42.6	45.9
Diabetes – %	10.9	10	12.8	9.5
History of cardio- /cerebro-vascular disease – %	30.3	36.8	23.7	19.6
Family history of dementia – %	55.5	53.8	55.2	64
History of tobacco smoking* – %	33.7	27.3	40.3	45.4
APOE* – %				
ε2ε2	0.1	0.1	0.2	0.3
ε2ε3	9.7	7.9	11.5	13
ε2ε4	2.6	2.1	3.3	2.9
ε3ε3	49.2	47.5	51.2	51.3
ε3ε4	31.6	33.4	29.9	27.8
ε4ε4	6.7	8.9	3.9	4.8
A-beta positivity* – %	53.3	62.2	24.2	25.8
Tau positivity* – %	58.6	62.9	34.1	42
Median follow-up – yr	3.1 (2.0 to 5.6)	3.1 (2.0 to 5.7)	3.1 (2.0 to 4.8)	4.0 (1.7 to 7.8)
WM-PVS count	422 (333 to 526)	379 (306 to 474)	481 (388 to 586)	453 (368 to 539)
BG-PVS count	122	104	148	139

	(97 to 156)	(87 to 128)	(123 to 173)	(117.75 to 169)
WM-PVS diameter – mm	2.00 (1.90 to 2.12)	2.08 (1.97 to 2.18)	1.92 (1.85 to 2.01)	1.94 (1.86 to 2.04)
BG-PVS diameter – mm	1.60 (1.53 to 1.68)	1.65 (1.58 to 1.71)	1.55 (1.51 to 1.60)	1.56 (1.52 to 1.62)
P-WML volume – mm ³	286 (0 to 1535)	592 (49 to 1923)	56 (0 to 842)	101.50 (0 to 829.25)
D-WML volume – mm ³	25 (3 to 90)	39 (7 to 110)	12 (1 to 57.25)	20.50 (4 to 73.25)

61 Plus-minus values are means ± standard deviations, entries with parentheses are
62 medians (interquartile range). Data from three studies — the Alzheimer’s Disease
63 Neuroimaging Initiative (ADNI), the National Alzheimer’s Coordinating Center (NACC),
64 and the Open Access Series of Imaging Studies (OASIS) — are shown. *Total number
65 of subjects with available history of tobacco smoking, Apolipoprotein E (APOE)
66 genotype, amyloid-beta status and tau status were 3297, 3353, 2118, and 1625,
67 respectively.

68
69
70
71

72 **Table S9. Estimates and adjusted significance of the interaction term “marker” by**
 73 **“time” in linear mixed-effects models with the grey matter volume, cortical**
 74 **thickness, or white matter volume as dependent variables.**

Marker	Grey matter volume (mm ³ / year)	Cortical thickness (µm / year)	White matter volume (mm ³ / year)
WM-PVS count (per 100-unit)	709±100, P<0.001	4±1, P<0.001	591±84, P<0.001
BG-PVS count (per 10-unit)	187±33, P<0.001	1±0, P<0.001	142±26, P<0.001
WM-PVS diameter (per 0.1 mm)	-382±84, P<0.001	-3±1, P<0.001	-64±67, P=0.675
BG-PVS diameter (per 0.1 mm)	-576±137, P<0.001	-5±1, P<0.001	-283±108, P=0.018
P-WML log volume	-288±46, P<0.001	-2±0, P<0.001	-73±35, P=0.08
D-WML log volume	-189±62, P=0.004	-1±0, P=0.01	-2±48, P=1

75
 76 Estimates indicate the value of the dependent variable preserved (positive values) or
 77 lost (negative values) per year for each unit increase in the marker. All models were
 78 adjusted for age, sex, race, educational level, body mass index, CDR global score at
 79 the baseline, history of diabetes, cardio-/cerebro-vascular disease, hypertension,
 80 dyslipidemia, family history of dementia, intracranial volume, the baseline value of the
 81 dependent variable (grey matter volume, cortical thickness, or white matter volume),
 82 field strength, manufacturer, and intra-individual consistency of the protocol used for the
 83 longitudinal MRI acquisitions (consistent versus non-consistent protocol). “P” indicates
 84 adjusted P-values.

85
 86

87
88
89

Table S10. Regional estimates (mm³/year) and adjusted significance of the interaction term “marker” by “time” in linear mixed-effects models with the volume of the corresponding region as dependent variable.

Region	WM-PVS count	BG-PVS count	WM-PVS diameter	BG-PVS diameter	P-WML log volume	D-WML log volume
lh_bankssts	3.3 ± 0.63, P<0.001	0.49 ± 0.21, P=1	-1.52 ± 0.53, P=0.344	-1.55 ± 0.86, P=1	-0.87 ± 0.29, P=0.213	-0.56 ± 0.39, P=1
lh_caudalanteriorcingulate	0.43 ± 0.48, P=1	-0.05 ± 0.16, P=1	0.9 ± 0.4, P=1	1.2 ± 0.66, P=1	-0.14 ± 0.22, P=1	-0.21 ± 0.3, P=1
lh_caudalmiddlefrontal	4.83 ± 1.31, P=0.019	0.93 ± 0.44, P=1	-0.34 ± 1.09, P=1	-0.69 ± 1.78, P=1	-2.19 ± 0.58, P=0.016	-0.49 ± 0.79, P=1
lh_cuneus	3.19 ± 0.85, P=0.015	0.24 ± 0.28, P=1	0.15 ± 0.71, P=1	-0.3 ± 1.16, P=1	-0.53 ± 0.38, P=1	-1.15 ± 0.52, P=1
lh_entorhinal	6.49 ± 0.94, P<0.001	2.47 ± 0.31, P<0.001	-3.83 ± 0.79, P<0.001	-6.76 ± 1.28, P<0.001	-1.97 ± 0.43, P<0.001	-0.04 ± 0.58, P=1
lh_fusiform	21.26 ± 2.56, P<0.001	7.13 ± 0.84, P<0.001	-13.66 ± 2.15, P<0.001	-21.65 ± 3.49, P<0.001	-6.01 ± 1.18, P<0.001	-3.74 ± 1.59, P=1
lh_inferiorparietal	19.68 ± 2.93, P<0.001	5.85 ± 0.97, P<0.001	-9.22 ± 2.47, P<0.016	-17.55 ± 4, P=0.001	-6.59 ± 1.34, P<0.001	-4.41 ± 1.81, P=1
lh_inferiortemporal	24.5 ± 3.08, P<0.001	8.04 ± 1.01, P<0.001	-13.43 ± 2.59, P<0.001	-24.06 ± 4.2, P<0.001	-7.63 ± 1.41, P<0.001	-4.67 ± 1.91, P=1
lh_isthmuscingulate	3.78 ± 0.59, P<0.001	0.78 ± 0.2, P=0.006	-0.59 ± 0.5, P=1	-1.77 ± 0.81, P=1	-1.14 ± 0.27, P=0.002	-0.08 ± 0.36, P=1
lh_lateraloccipital	20.63 ± 3.42, P<0.001	7.35 ± 1.13, P<0.001	-12.97 ± 2.86, P=0.001	-25.33 ± 4.65, P<0.001	-8.26 ± 1.55, P<0.001	-6.05 ± 2.1, P=0.34
lh_lateralorbitofrontal	7.41 ± 1.45, P<0.001	2.54 ± 0.48, P<0.001	-1.34 ± 1.21, P=1	-4.28 ± 1.97, P=1	-2.59 ± 0.65, P=0.006	-1.02 ± 0.88, P=1
lh_lingual	8.04 ± 1.76, P<0.001	1.89 ± 0.59, P<0.113	-2.69 ± 1.48, P=1	-5.02 ± 2.41, P=1	-1.6 ± 0.8, P=1	-0.99 ± 1.08, P=1
lh_medialorbitofrontal	6.28 ± 1.49, P=0.002	1.56 ± 0.49, P=0.137	0.01 ± 1.25, P=1	-0.58 ± 2.04, P=1	-2.22 ± 0.68, P=0.096	-0.66 ± 0.92, P=1
lh_middletemporal	24.18 ± 3.01, P<0.001	8.47 ± 0.98, P<0.001	-14.21 ± 2.53, P<0.001	-26.78 ± 4.09, P<0.001	-6.71 ± 1.38, P<0.001	-3.15 ± 1.87, P=1
lh parahippocampal	5.58 ± 0.7, P<0.001	1.67 ± 0.23, P<0.001	-2.83 ± 0.59, P<0.001	-4.63 ± 0.96, P<0.001	-1.32 ± 0.32, P=0.004	-0.4 ± 0.43, P=1
lh_paracentral	0.79 ± 0.7, P=1	-0.13 ± 0.24, P=1	0.62 ± 0.59, P=1	1.25 ± 0.96, P=1	-0.96 ± 0.31, P=0.184	-0.71 ± 0.42, P=1
lh_parsopercularis	5.58 ± 0.97, P<0.001	1.29 ± 0.32, P=0.006	-2.1 ± 0.81, P=0.858	-2.92 ± 1.33, P=1	-1.65 ± 0.44, P=0.015	-0.77 ± 0.59, P=1
lh_parsorbitalis	3 ± 0.69, P=0.001	1.14 ± 0.23, P<0.001	-1.54 ± 0.58, P=0.661	-3.02 ± 0.94, P=0.116	-0.66 ± 0.31, P=1	-0.23 ± 0.42, P=1
lh_parstriangularis	3.32 ± 0.82, P=0.005	0.76 ± 0.27, P=0.5	-0.78 ± 0.69, P=1	-1.73 ± 1.12, P=1	-1.03 ± 0.37, P=0.456	-0.64 ± 0.5, P=1
lh_pericalcarine	0.03 ± 0.73, P=1	-1.01 ± 0.24, P=0.003	2.01 ± 0.6, P=0.076	3.25 ± 0.99, P=0.092	0.53 ± 0.33, P=1	-0.03 ± 0.44, P=1
lh_postcentral	5.3 ± 2, P=0.689	0.56 ± 0.68, P=1	0.34 ± 1.67, P=1	-1.76 ± 2.73, P=1	-2.55 ± 0.89, P=0.352	-1.21 ± 1.2, P=1
lh_posteriorcingulate	2.75 ± 0.72, P=0.013	0.55 ± 0.24, P=1	-0.39 ± 0.61, P=1	0.32 ± 0.99, P=1	-0.67 ± 0.33, P=1	-0.37 ± 0.44, P=1
lh_precentral	10.19 ± 2.54, P=0.005	1.1 ± 0.87, P=1	-4.1 ± 2.12, P=1	-6.74 ± 3.48, P=1	-4.67 ± 1.13, P=0.003	-2.17 ± 1.53, P=1
lh_precuneus	9.64 ± 1.93, P<0.001	2.46 ± 0.65, P=0.013	-2.24 ± 1.62, P=1	-3.73 ± 2.64, P=1	-3.59 ± 0.87, P=0.003	-2.64 ± 1.18, P=1
lh_rostral anteriorcingulate	1.39 ± 0.59, P=1	0.7 ± 0.2, P=0.032	-0.59 ± 0.5, P=1	-1.48 ± 0.81, P=1	-0.3 ± 0.27, P=1	-0.68 ± 0.36, P=1
lh_rostralmiddlefrontal	10.25 ± 3.04, P=0.065	1.23 ± 1.03, P=1	-0.89 ± 2.54, P=1	1.21 ± 4.15, P=1	-4.65 ± 1.36, P=0.053	-2.48 ± 1.84, P=1
lh_superiorfrontal	12.01 ± 4.28, P=0.437	3.73 ± 1.44, P=0.831	1.03 ± 3.58, P=1	-2.23 ± 5.84, P=1	-5.35 ± 1.92, P=0.466	-3.49 ± 2.6, P=1
lh_superiorparietal	10.82 ± 2.99, P=0.026	3.08 ± 1, P=0.19	-3.55 ± 2.5, P=1	-6.17 ± 4.08, P=1	-6 ± 1.34, P<0.001	-4.91 ± 1.82, P=0.595
lh_superiortemporal	24.98 ± 2.96, P<0.001	7.78 ± 0.98, P<0.001	-13.58 ± 2.49, P<0.001	-23.93 ± 4.04, P<0.001	-7.22 ± 1.35, P<0.001	-2.82 ± 1.84, P=1
lh_supramarginal	17.13 ± 2.57, P<0.001	3.48 ± 0.86, P=0.005	-4.76 ± 2.16, P=1	-8.85 ± 3.52, P=1	-6.68 ± 1.16, P<0.001	-2.12 ± 1.58, P=1

lh_frontalpole	2.07 ± 0.47, P=0.001	0.76 ± 0.16, P<0.001	-1.01 ± 0.4, P=0.937	-1.56 ± 0.64, P=1	-0.22 ± 0.22, P=1	-0.46 ± 0.29, P=1
lh_temporalpole	7.68 ± 1.22, P<0.001	2.19 ± 0.4, P<0.001	-3.33 ± 1.03, P=0.102	-6.46 ± 1.66, P=0.009	-1.85 ± 0.56, P=0.083	-0.3 ± 0.76, P=1
lh_transversetemporal	1.71 ± 0.33, P<0.001	0.34 ± 0.11, P=0.157	-1.1 ± 0.27, P=0.006	-1.47 ± 0.45, P=0.088	-0.49 ± 0.15, P=0.076	-0.87 ± 0.2, P=0.001
lh_insula	8.98 ± 1.56, P<0.001	2.11 ± 0.51, P=0.004	-2.72 ± 1.31, P=1	-3.09 ± 2.13, P=1	-3.15 ± 0.71, P=0.001	-1.13 ± 0.96, P=1
rh_bankssts	2.9 ± 0.51, P<0.001	0.85 ± 0.17, P<0.001	-2 ± 0.42, P<0.001	-2.91 ± 0.69, P=0.002	-1.09 ± 0.23, P<0.001	-0.98 ± 0.31, P=0.131
rh_caudalanteriorcingulate	1.05 ± 0.64, P=1	-0.02 ± 0.21, P=1	0 ± 0.54, P=1	1 ± 0.87, P=1	-0.21 ± 0.29, P=1	-0.09 ± 0.39, P=1
rh_caudalmiddlefrontal	4.31 ± 1.24, P=0.046	0.81 ± 0.42, P=1	-0.9 ± 1.04, P=1	-1.34 ± 1.7, P=1	-2.69 ± 0.56, P<0.001	-1.24 ± 0.75, P=1
rh_cuneus	3.39 ± 0.94, P=0.026	1.13 ± 0.31, P=0.027	-1.03 ± 0.78, P=1	-2.8 ± 1.27, P=1	-1.36 ± 0.42, P=0.109	-1.82 ± 0.57, P=0.123
rh_entorhinal	5.68 ± 0.9, P<0.001	2.42 ± 0.29, P<0.001	-3.46 ± 0.76, P<0.001	-7.42 ± 1.22, P<0.001	-1.57 ± 0.41, P=0.012	-1.28 ± 0.56, P=1
rh_fusiform	18.26 ± 2.39, P<0.001	7.02 ± 0.78, P<0.001	-11.15 ± 2.01, P<0.001	-20.61 ± 3.25, P<0.001	-7.09 ± 1.09, P=0.001	-4.93 ± 1.48, P=0.073
rh_inferiorparietal	17.91 ± 3.32, P<0.001	5.6 ± 1.11, P<0.001	-7.45 ± 2.79, P=0.652	-17.12 ± 4.52, P=0.014	-7.36 ± 1.5, P<0.001	-4.76 ± 2.03, P=1
rh_inferiortemporal	22.17 ± 2.92, P<0.001	7.42 ± 0.96, P<0.001	-12.12 ± 2.46, P<0.001	-21.49 ± 3.99, P<0.001	-8.36 ± 1.33, P<0.001	-4.53 ± 1.81, P=1
rh_isthmuscingulate	2.82 ± 0.65, P=0.001	0.9 ± 0.22, P=0.003	-0.42 ± 0.54, P=1	-1.4 ± 0.89, P=1	-0.92 ± 0.29, P=0.154	-0.45 ± 0.4, P=1
rh_lateraloccipital	18.44 ± 3.44, P<0.001	4.96 ± 1.15, P<0.001	-9.21 ± 2.89, P=0.125	-16.17 ± 4.69, P=0.05	-9.18 ± 1.56, P<0.001	-6.2 ± 2.1, P=0.281
rh_lateralorbitofrontal	10.67 ± 1.66, P<0.001	3.03 ± 0.55, P<0.001	-3.22 ± 1.4, P=1	-7.24 ± 2.27, P=0.123	-3.5 ± 0.75, P<0.001	-0.4 ± 1.02, P=1
rh_lingual	8.67 ± 1.8, P<0.001	3.26 ± 0.6, P<0.001	-3.26 ± 1.51, P=1	-6.57 ± 2.45, P=0.645	-3.54 ± 0.81, P=0.001	-2.14 ± 1.1, P=1
rh_medialorbitofrontal	6.02 ± 1.52, P=0.007	2.37 ± 0.5, P<0.001	-2.23 ± 1.27, P=1	-4.19 ± 2.07, P=1	-2.68 ± 0.69, P=0.009	-1.79 ± 0.93, P=1
rh_middletemporal	26.71 ± 3.3, P<0.001	9.94 ± 1.08, P<0.001	-18.26 ± 2.77, P<0.001	-31.5 ± 4.48, P<0.001	-8.87 ± 1.51, P<0.001	-4.03 ± 2.04, P=1
rh parahippocampal	4.59 ± 0.68, P<0.001	1.84 ± 0.22, P<0.001	-2.63 ± 0.57, P<0.001	-5.29 ± 0.92, P<0.001	-1.46 ± 0.31, P<0.001	-0.79 ± 0.42, P=1
rh_paracentral	0.42 ± 0.86, P=1	-0.51 ± 0.29, P=1	0.41 ± 0.72, P=1	2.35 ± 1.17, P=1	-0.64 ± 0.38, P=1	-0.82 ± 0.52, P=1
rh_parsopercularis	5.6 ± 0.92, P<0.001	1.89 ± 0.3, P<0.001	-3.04 ± 0.77, P=0.007	-5.7 ± 1.25, P<0.001	-2.2 ± 0.42, P<0.001	-0.6 ± 0.56, P=1
rh_parsorbitalis	4.27 ± 0.89, P<0.001	2.17 ± 0.29, P<0.001	-3.37 ± 0.74, P=0.001	-6.75 ± 1.2, P<0.001	-1.44 ± 0.4, P=0.032	-1.03 ± 0.55, P=1
rh_parstriangularis	4.35 ± 1, P=0.001	1.99 ± 0.33, P<0.001	-3.44 ± 0.83, P=0.003	-6.29 ± 1.35, P<0.001	-2.09 ± 0.45, P<0.001	-1.04 ± 0.61, P=1
rh_pericalcarine	0.93 ± 0.83, P=1	-1.07 ± 0.28, P=0.01	1.85 ± 0.69, P=0.647	3.44 ± 1.13, P=0.2	-0.15 ± 0.37, P=1	0 ± 0.5, P=1
rh_postcentral	2.4 ± 2.03, P=1	0.43 ± 0.69, P=1	0.97 ± 1.69, P=1	-0.06 ± 2.77, P=1	-1.58 ± 0.91, P=1	-1.65 ± 1.23, P=1
rh_posteriorcingulate	3.02 ± 0.77, P=0.007	1.09 ± 0.25, P=0.002	-0.66 ± 0.64, P=1	-1.21 ± 1.05, P=1	-1.05 ± 0.35, P=0.228	-0.5 ± 0.47, P=1
rh_precentral	8.97 ± 2.56, P=0.04	2.14 ± 0.88, P=1	-5.21 ± 2.13, P=1	-9.12 ± 3.5, P=0.795	-4.91 ± 1.13, P=0.001	-3.43 ± 1.54, P=1
rh_precuneus	8.95 ± 1.97, P=0.001	2.55 ± 0.66, P=0.01	-3.99 ± 1.65, P=1	-5.67 ± 2.69, P=1	-4.38 ± 0.89, P<0.001	-3.95 ± 1.2, P=0.09
rh_rostralanteriorcingulate	2.25 ± 0.51, P=0.001	0.36 ± 0.17, P=1	-0.35 ± 0.43, P=1	-0.22 ± 0.7, P=1	-0.57 ± 0.23, P=1	-0.19 ± 0.31, P=1
rh_rostralmiddlefrontal	12.57 ± 3.19, P=0.007	5.41 ± 1.07, P<0.001	-6.83 ± 2.67, P=0.921	-13.38 ± 4.35, P=0.184	-7.04 ± 1.43, P<0.001	-4.87 ± 1.94, P=1
rh_superiorfrontal	11.89 ± 4.45, P=0.654	3.35 ± 1.49, P=1	1.65 ± 3.72, P=1	-0.25 ± 6.07, P=1	-7.76 ± 2, P=0.009	-3.85 ± 2.7, P=1
rh_superiorparietal	9.03 ± 2.95, P=0.195	2.26 ± 0.99, P=1	-0.74 ± 2.47, P=1	-3.58 ± 4.02, P=1	-5.43 ± 1.32, P=0.004	-5.08 ± 1.79, P=0.395
rh_superiortemporal	24.55 ± 2.96, P<0.001	9.99 ± 0.97, P<0.001	-18.43 ± 2.48, P<0.001	-31.12 ± 4.02, P<0.001	-8.92 ± 1.35, P<0.001	-5.29 ± 1.83, P=0.34
rh_supramarginal	13.95 ± 2.2, P<0.001	4.96 ± 0.73, P<0.001	-8.98 ± 1.85, P<0.001	-16.7 ± 3, P<0.001	-5.36 ± 1, P<0.001	-2.46 ± 1.35, P=1

rh_frontalpole	3.01 ± 0.62, P<0.001	1.22 ± 0.2, P<0.001	-2.15 ± 0.52, P=0.003	-3.34 ± 0.85, P=0.007	-1.3 ± 0.28, P<0.001	-1.44 ± 0.38, P=0.014
rh_temporalpole	6.78 ± 1.27, P<0.001	2.7 ± 0.41, P<0.001	-4.87 ± 1.06, P<0.001	-8.03 ± 1.73, P<0.001	-2.87 ± 0.58, P<0.001	-2.44 ± 0.78, P=0.161
rh_transversetemporal	1.67 ± 0.28, P<0.001	0.58 ± 0.09, P<0.001	-0.84 ± 0.23, P=0.028	-1.94 ± 0.38, P<0.001	-0.45 ± 0.13, P=0.032	-0.54 ± 0.17, P=0.122
rh_insula	9.11 ± 1.55, P<0.001	2.83 ± 0.51, P<0.001	-3.75 ± 1.3, P=0.342	-6.26 ± 2.11, P=0.266	-3.52 ± 0.71, P<0.001	-0.9 ± 0.96, P=1
Left-Thalamus-Proper	-1.49 ± 1.27, P=1	-0.49 ± 0.42, P=1	0.29 ± 1.06, P=1	2.46 ± 1.73, P=1	-1.01 ± 0.57, P=1	-1.13 ± 0.77, P=1
Right-Thalamus-Proper	1.04 ± 1.29, P=1	-0.06 ± 0.43, P=1	-0.97 ± 1.08, P=1	1.25 ± 1.76, P=1	-1.36 ± 0.59, P=1	-1.18 ± 0.79, P=1
Right-Lateral-Ventricle	-117.67 ± 13.2, P<0.001	-34.94 ± 4.29, P<0.001	55.56 ± 11.12, P<0.001	92.96 ± 18.04, P<0.001	47.6 ± 6.1, P<0.001	9.66 ± 8.23, P=1
Left-Hippocampus	2.49 ± 0.88, P=0.43	1.88 ± 0.29, P<0.001	-4.12 ± 0.74, P<0.001	-6.63 ± 1.19, P<0.001	-0.55 ± 0.4, P=1	-1.81 ± 0.54, P=0.074
Left-Lateral-Ventricle	-112.15 ± 15.48, P<0.001	-27.37 ± 5.03, P<0.001	37.26 ± 13.03, P=0.367	63.74 ± 21.14, P=0.223	44.61 ± 7.15, P<0.001	-2.75 ± 9.62, P=1
Right-Putamen	2.98 ± 1.2, P=1	0.36 ± 0.4, P=1	-2.2 ± 1, P=1	-0.15 ± 1.64, P=1	-0.76 ± 0.55, P=1	-0.38 ± 0.74, P=1
Right-Amygdala	1.01 ± 0.42, P=1	0.33 ± 0.14, P=1	-1.81 ± 0.35, P=0.001	-1.62 ± 0.58, P=0.438	-0.33 ± 0.19, P=1	-1.02 ± 0.26, P=0.007
Left-Putamen	1.1 ± 1.2, P=1	-0.4 ± 0.4, P=1	-1.69 ± 1.01, P=1	3.47 ± 1.64, P=1	0.04 ± 0.55, P=1	-0.61 ± 0.74, P=1
Right-VentralDC	-1.25 ± 0.57, P=1	0.29 ± 0.2, P=1	-0.47 ± 0.47, P=1	-0.45 ± 0.78, P=1	0.08 ± 0.25, P=1	-0.38 ± 0.33, P=1
Left-VentralDC	-2.39 ± 0.56, P=0.002	0.28 ± 0.2, P=1	0.04 ± 0.46, P=1	-0.38 ± 0.77, P=1	0.24 ± 0.24, P=1	-0.48 ± 0.32, P=1
Right-Hippocampus	3.17 ± 0.91, P=0.046	2.16 ± 0.3, P<0.001	-3.95 ± 0.76, P<0.001	-7.34 ± 1.24, P<0.001	-0.69 ± 0.42, P=1	-1.88 ± 0.56, P=0.071
Left-Amygdala	0.63 ± 0.39, P=1	0.18 ± 0.13, P=1	-1.13 ± 0.33, P=0.048	-1.57 ± 0.53, P=0.282	-0.2 ± 0.18, P=1	-0.52 ± 0.24, P=1
Left-Pallidum	-0.84 ± 0.45, P=1	0.25 ± 0.15, P=1	-0.21 ± 0.38, P=1	-1.2 ± 0.62, P=1	0.27 ± 0.2, P=1	0.1 ± 0.27, P=1
Right-Pallidum	-1.3 ± 0.45, P=0.296	0.33 ± 0.15, P=1	-0.29 ± 0.37, P=1	-1.1 ± 0.61, P=1	0.61 ± 0.2, P=0.215	0.13 ± 0.27, P=1
Left-Caudate	1.48 ± 0.96, P=1	-0.55 ± 0.32, P=1	-0.01 ± 0.8, P=1	5.32 ± 1.3, P=0.004	-0.11 ± 0.43, P=1	-0.07 ± 0.58, P=1
Right-Caudate	-0.33 ± 1.02, P=1	-0.52 ± 0.34, P=1	0.48 ± 0.85, P=1	5.27 ± 1.38, P=0.012	0.49 ± 0.46, P=1	-0.27 ± 0.62, P=1
3rd-ventricle	0.1 ± 0.79, P=1	-1.28 ± 0.26, P<0.001	2.04 ± 0.66, P=0.163	3.54 ± 1.07, P=0.079	0.7 ± 0.36, P=1	0.93 ± 0.48, P=1
4th-ventricle	-0.37 ± 0.87, P=1	-1.18 ± 0.29, P=0.004	0.99 ± 0.72, P=1	1.84 ± 1.18, P=1	1.46 ± 0.39, P=0.016	0.04 ± 0.53, P=1

90
91
92
93
94
95
96
97
98
99
100

The estimates indicate the grey matter or ventricular volume for that region preserved (positive values) or lost (negative values) per year for each unit increase in the marker assessed at the baseline. Units were 100 for WM-PVS count, 10 for BG-PVS count, 0.1 mm for WM- and BG-PVS diameter, and 1 log for P-WML and D-WML volume. See also Figure 2A. Parcellations performed by Freesurfer according to the Desikan-Killany atlas. Bankssts: Banks of the Superior Temporal Sulcus; DC: diencephalon. Lh: left hemisphere; Rh: right hemisphere. Models were fully adjusted for all the same covariates included in the main linear mixed-effect model. P values are corrected for 86 comparisons with Holm-Bonferroni procedure.

101
102
103

Table S11. Regional estimates ($\mu\text{m}/\text{year}$) and adjusted significance of the interaction term “marker” by “time” in linear mixed-effects models with the thickness of the corresponding region as dependent variable.

Region	WM-PVS count	BG-PVS count	WM-PVS diameter	BG-PVS diameter	P-WML log volume	D-WML log volume
lh_bankssts	2.89 ± 0.49, P<0.001	0.98 ± 0.17, P<0.001	-2.22 ± 0.42, P<0.001	-3.44 ± 0.69, P<0.001	-0.69 ± 0.23, P=0.148	-0.56 ± 0.31, P=1
lh_caudalanteriorcingulate	2.05 ± 0.54, P=0.012	0.64 ± 0.18, P=0.032	-0.7 ± 0.47, P=1	-0.79 ± 0.76, P=1	-0.75 ± 0.25, P=0.175	-0.07 ± 0.34, P=1
lh_caudalmiddlefrontal	1.61 ± 0.41, P=0.007	0.66 ± 0.14, P<0.001	-0.81 ± 0.35, P=1	-1.69 ± 0.58, P=0.247	-0.77 ± 0.18, P=0.002	-0.23 ± 0.25, P=1
lh_cuneus	0.49 ± 0.31, P=1	-0.05 ± 0.11, P=1	0.22 ± 0.26, P=1	0 ± 0.43, P=1	-0.2 ± 0.14, P=1	-0.45 ± 0.19, P=1
lh_entorhinal	5.79 ± 0.95, P<0.001	2.97 ± 0.31, P<0.001	-4.7 ± 0.81, P<0.001	-8.9 ± 1.32, P<0.001	-1.92 ± 0.44, P<0.001	-1.16 ± 0.6, P=1
lh_frontalpole	1.3 ± 0.64, P=1	1.12 ± 0.22, P<0.001	-0.72 ± 0.54, P=1	-3 ± 0.89, P=0.052	-0.12 ± 0.29, P=1	-0.17 ± 0.39, P=1
lh_fusiform	3.79 ± 0.52, P<0.001	1.64 ± 0.17, P<0.001	-3 ± 0.45, P<0.001	-5.42 ± 0.73, P<0.001	-1.24 ± 0.24, P<0.001	-0.65 ± 0.33, P=1
lh_inferiorparietal	2.34 ± 0.43, P<0.001	0.8 ± 0.14, P<0.001	-1.56 ± 0.37, P=0.002	-2.69 ± 0.6, P<0.001	-0.92 ± 0.19, P<0.001	-0.64 ± 0.27, P=1
lh_inferiortemporal	3.29 ± 0.52, P<0.001	1.54 ± 0.17, P<0.001	-2.25 ± 0.45, P<0.001	-4.57 ± 0.73, P<0.001	-0.95 ± 0.24, P=0.005	-0.57 ± 0.33, P=1
lh_insula	2.26 ± 0.56, P=0.004	0.89 ± 0.19, P<0.001	-1.03 ± 0.48, P=1	-2.04 ± 0.78, P=0.645	-0.59 ± 0.26, P=1	-0.26 ± 0.35, P=1
lh_isthmuscingulate	1.74 ± 0.38, P<0.001	0.64 ± 0.13, P<0.001	-0.73 ± 0.32, P=1	-1.87 ± 0.53, P=0.029	-0.52 ± 0.17, P=0.164	-0.17 ± 0.24, P=1
lh_lateraloccipital	1.41 ± 0.36, P=0.007	0.59 ± 0.12, P<0.001	-0.85 ± 0.31, P=0.414	-2.04 ± 0.5, P=0.004	-0.66 ± 0.16, P=0.003	-0.44 ± 0.22, P=1
lh_lateralorbitofrontal	1.66 ± 0.42, P=0.006	0.93 ± 0.14, P<0.001	-0.74 ± 0.36, P=1	-1.8 ± 0.58, P=0.143	-0.63 ± 0.19, P=0.06	-0.15 ± 0.26, P=1
lh_lingual	0.64 ± 0.34, P=1	0.22 ± 0.12, P=1	-0.17 ± 0.29, P=1	-0.68 ± 0.47, P=1	-0.21 ± 0.15, P=1	-0.24 ± 0.21, P=1
lh_medialorbitofrontal	1.53 ± 0.43, P=0.03	0.74 ± 0.15, P<0.001	-0.49 ± 0.37, P=1	-0.97 ± 0.6, P=1	-0.58 ± 0.2, P=0.239	-0.29 ± 0.27, P=1
lh_middletemporal	3.73 ± 0.53, P<0.001	1.43 ± 0.18, P<0.001	-2.27 ± 0.46, P<0.001	-4.34 ± 0.74, P<0.001	-1.05 ± 0.24, P=0.001	-0.49 ± 0.34, P=1
lh_paracentral	0.63 ± 0.42, P=1	0.37 ± 0.15, P=0.872	-0.68 ± 0.36, P=1	-1.09 ± 0.59, P=1	-0.52 ± 0.19, P=0.383	-0.55 ± 0.26, P=1
lh parahippocampal	4.77 ± 0.71, P<0.001	1.7 ± 0.24, P<0.001	-2.72 ± 0.61, P=0.001	-4.98 ± 0.99, P<0.001	-1.55 ± 0.32, P<0.001	-0.86 ± 0.44, P=1
lh_parsopercularis	1.91 ± 0.39, P<0.001	0.68 ± 0.13, P<0.001	-1.02 ± 0.33, P=0.138	-1.78 ± 0.54, P=0.069	-0.76 ± 0.17, P=0.001	-0.35 ± 0.24, P=1
lh_parsorbitalis	0.81 ± 0.49, P=1	0.71 ± 0.17, P=0.002	-0.49 ± 0.42, P=1	-1.47 ± 0.69, P=1	-0.22 ± 0.22, P=1	0.17 ± 0.3, P=1
lh_parstriangularis	1.01 ± 0.36, P=0.338	0.44 ± 0.12, P=0.023	-0.29 ± 0.3, P=1	-0.87 ± 0.5, P=1	-0.39 ± 0.16, P=1	-0.39 ± 0.22, P=1
lh_pericalcarine	-0.43 ± 0.34, P=1	-0.25 ± 0.12, P=1	0.79 ± 0.29, P=0.493	0.64 ± 0.48, P=1	0.07 ± 0.15, P=1	-0.14 ± 0.21, P=1
lh_postcentral	0.93 ± 0.33, P=0.323	0.33 ± 0.11, P=0.239	-0.39 ± 0.28, P=1	-1.42 ± 0.46, P=0.138	-0.41 ± 0.15, P=0.391	-0.44 ± 0.2, P=1
lh_posteriorcingulate	1.76 ± 0.38, P<0.001	0.64 ± 0.13, P<0.001	-0.78 ± 0.32, P=1	-1.22 ± 0.52, P=1	-0.77 ± 0.17, P<0.001	-0.17 ± 0.23, P=1
lh_precentral	2.07 ± 0.42, P<0.001	0.79 ± 0.15, P<0.001	-1.49 ± 0.36, P=0.002	-3 ± 0.59, P<0.001	-0.82 ± 0.19, P=0.001	-0.58 ± 0.26, P=1
lh_precuneus	1.49 ± 0.37, P=0.004	0.67 ± 0.12, P<0.001	-1.14 ± 0.31, P=0.02	-1.96 ± 0.51, P=0.009	-0.66 ± 0.17, P=0.005	-0.63 ± 0.23, P=0.383
lh_rostralanteriorcingulate	0.63 ± 0.46, P=1	0.62 ± 0.16, P=0.005	-0.65 ± 0.4, P=1	-1.72 ± 0.64, P=0.538	-0.35 ± 0.21, P=1	-0.05 ± 0.29, P=1
lh_rostralmiddlefrontal	0.59 ± 0.34, P=1	0.15 ± 0.12, P=1	0 ± 0.29, P=1	0.11 ± 0.48, P=1	-0.38 ± 0.15, P=0.9	-0.24 ± 0.21, P=1
lh_superiorfrontal	0.83 ± 0.39, P=1	0.61 ± 0.13, P<0.001	-0.39 ± 0.33, P=1	-1.19 ± 0.54, P=1	-0.53 ± 0.17, P=0.161	-0.25 ± 0.24, P=1
lh_superiorparietal	1.37 ± 0.35, P=0.008	0.51 ± 0.12, P=0.002	-0.93 ± 0.3, P=0.135	-1.68 ± 0.49, P=0.046	-0.73 ± 0.16, P<0.001	-0.63 ± 0.22, P=0.265

lh_superiortemporal	3.5 ± 0.49, P<0.001	1.43 ± 0.16, P<0.001	-2.72 ± 0.42, P<0.001	-4.95 ± 0.68, P<0.001	-1.14 ± 0.23, P<0.001	-0.58 ± 0.31, P=1
lh_supramarginal	2.13 ± 0.39, P<0.001	0.71 ± 0.13, P<0.001	-1.29 ± 0.33, P=0.006	-2.57 ± 0.54, P<0.001	-0.96 ± 0.17, P<0.001	-0.41 ± 0.24, P=1
lh_temporalpole	3.86 ± 0.92, P=0.002	2.14 ± 0.31, P<0.001	-2.82 ± 0.79, P=0.026	-6.39 ± 1.28, P<0.001	-1.61 ± 0.42, P=0.009	-0.55 ± 0.58, P=1
lh_transversetemporal	3.43 ± 0.51, P<0.001	1.01 ± 0.18, P<0.001	-2.57 ± 0.44, P<0.001	-4.55 ± 0.71, P<0.001	-1.13 ± 0.23, P<0.001	-1.38 ± 0.32, P=0.001
rh_bankssts	2.7 ± 0.5, P<0.001	1.32 ± 0.17, P<0.001	-2.56 ± 0.43, P<0.001	-4.42 ± 0.7, P<0.001	-0.92 ± 0.23, P=0.004	-0.86 ± 0.31, P=0.433
rh_caudalanteriorcingulate	0.44 ± 0.58, P=1	0.15 ± 0.19, P=1	-0.07 ± 0.49, P=1	0.02 ± 0.8, P=1	-0.38 ± 0.26, P=1	-0.09 ± 0.36, P=1
rh_caudalmiddlefrontal	1.75 ± 0.42, P=0.002	0.67 ± 0.15, P<0.001	-1.37 ± 0.36, P=0.01	-2.27 ± 0.59, P<0.009	-0.97 ± 0.19, P<0.001	-0.68 ± 0.26, P=0.617
rh_cuneus	0.38 ± 0.34, P=1	0.24 ± 0.12, P=1	-0.3 ± 0.29, P=1	-0.94 ± 0.48, P=1	-0.29 ± 0.15, P=1	-0.73 ± 0.21, P=0.035
rh_entorhinal	5.76 ± 1.01, P<0.001	2.81 ± 0.34, P<0.001	-4.15 ± 0.87, P<0.001	-9.28 ± 1.41, P<0.001	-1.97 ± 0.46, P=0.002	-1.56 ± 0.64, P=1
rh_frontalpole	0.97 ± 0.69, P=1	0.97 ± 0.23, P=0.003	-0.52 ± 0.59, P=1	-3.24 ± 0.95, P=0.049	-0.53 ± 0.31, P=1	-0.52 ± 0.42, P=1
rh_fusiform	3.38 ± 0.54, P<0.001	1.77 ± 0.18, P<0.001	-2.8 ± 0.46, P<0.001	-5.67 ± 0.75, P<0.001	-1.31 ± 0.25, P<0.001	-0.98 ± 0.34, P=0.272
rh_inferiorparietal	1.65 ± 0.41, P=0.004	0.83 ± 0.14, P<0.001	-0.85 ± 0.35, P=1	-2.41 ± 0.57, P=0.002	-0.87 ± 0.18, P<0.001	-0.71 ± 0.25, P=0.339
rh_inferiortemporal	3.09 ± 0.51, P<0.001	1.53 ± 0.17, P<0.001	-2.27 ± 0.44, P<0.001	-4.81 ± 0.71, P<0.001	-1.29 ± 0.23, P<0.001	-0.9 ± 0.32, P=0.347
rh_insula	2 ± 0.58, P=0.043	1.28 ± 0.19, P<0.001	-1.8 ± 0.5, P=0.022	-3.53 ± 0.81, P<0.001	-0.89 ± 0.27, P=0.063	-0.48 ± 0.37, P=1
rh_isthmuscingulate	1.86 ± 0.42, P=0.001	0.75 ± 0.14, P<0.001	-1.01 ± 0.36, P=0.392	-2.3 ± 0.59, P=0.007	-0.59 ± 0.19, P=0.169	-0.19 ± 0.26, P=1
rh_lateraloccipital	0.84 ± 0.37, P=1	0.44 ± 0.13, P=0.03	-0.44 ± 0.31, P=1	-1.63 ± 0.51, P=0.109	-0.56 ± 0.17, P=0.056	-0.76 ± 0.23, P=0.06
rh_lateralorbitofrontal	2.17 ± 0.46, P<0.001	1.14 ± 0.15, P<0.001	-1.71 ± 0.4, P=0.001	-3.61 ± 0.64, P<0.001	-0.89 ± 0.21, P=0.002	-0.31 ± 0.29, P=1
rh_lingual	0.71 ± 0.36, P=1	0.51 ± 0.12, P=0.002	-0.79 ± 0.31, P=0.751	-1.63 ± 0.5, P=0.083	-0.32 ± 0.16, P=1	-0.41 ± 0.22, P=1
rh_medialorbitofrontal	1.19 ± 0.47, P=0.807	0.65 ± 0.16, P=0.003	-0.57 ± 0.4, P=1	-1.37 ± 0.65, P=1	-0.66 ± 0.21, P=0.141	-0.58 ± 0.29, P=1
rh_middletemporal	3.14 ± 0.52, P<0.001	1.55 ± 0.17, P<0.001	-2.46 ± 0.44, P<0.001	-4.97 ± 0.71, P<0.001	-1.04 ± 0.24, P=0.001	-0.75 ± 0.32, P=1
rh_paracentral	0.82 ± 0.44, P=1	0.32 ± 0.15, P=1	-0.7 ± 0.37, P=1	-0.86 ± 0.61, P=1	-0.49 ± 0.19, P=0.808	-0.52 ± 0.27, P=1
rh parahippocampal	4.24 ± 0.71, P<0.001	1.93 ± 0.24, P<0.001	-2.94 ± 0.61, P<0.001	-5.91 ± 0.99, P<0.001	-1.66 ± 0.33, P<0.001	-1.17 ± 0.45, P=0.654
rh_parsopercularis	2.07 ± 0.4, P<0.001	0.87 ± 0.14, P<0.001	-1.59 ± 0.34, P<0.001	-2.93 ± 0.56, P<0.001	-0.87 ± 0.18, P<0.001	-0.35 ± 0.25, P=1
rh_parsorbitalis	0.94 ± 0.49, P=1	1.09 ± 0.17, P<0.001	-1.14 ± 0.42, P=0.485	-3.47 ± 0.68, P<0.001	-0.57 ± 0.22, P=0.784	-0.46 ± 0.31, P=1
rh_parstriangularis	1.23 ± 0.37, P=0.057	0.62 ± 0.13, P<0.001	-1.23 ± 0.31, P=0.006	-2.1 ± 0.51, P=0.003	-0.59 ± 0.16, P=0.025	-0.53 ± 0.23, P=1
rh_pericalcarine	-0.39 ± 0.35, P=1	-0.23 ± 0.12, P=1	0.23 ± 0.3, P=1	0.21 ± 0.49, P=1	0 ± 0.16, P=1	-0.49 ± 0.21, P=1
rh_postcentral	0.56 ± 0.34, P=1	0.38 ± 0.12, P=0.09	-0.3 ± 0.29, P=1	-1.31 ± 0.47, P=0.395	-0.36 ± 0.15, P=1	-0.44 ± 0.21, P=1
rh_posteriorcingulate	2.18 ± 0.43, P<0.001	0.79 ± 0.14, P<0.001	-0.88 ± 0.37, P=1	-1.66 ± 0.59, P=0.364	-0.83 ± 0.19, P=0.001	-0.01 ± 0.27, P=1
rh_precentral	1.97 ± 0.43, P<0.001	0.87 ± 0.15, P<0.001	-1.63 ± 0.36, P=0.001	-3.18 ± 0.6, P<0.001	-0.89 ± 0.19, P<0.001	-0.62 ± 0.26, P=1
rh_precuneus	1.65 ± 0.36, P<0.001	0.66 ± 0.12, P<0.001	-1.23 ± 0.31, P=0.004	-2.19 ± 0.5, P<0.001	-0.73 ± 0.16, P<0.001	-0.81 ± 0.22, P=0.019
rh_rostralanteriorcingulate	2.07 ± 0.52, P=0.006	0.52 ± 0.18, P=0.215	-0.42 ± 0.45, P=1	-0.73 ± 0.73, P=1	-0.5 ± 0.24, P=1	0.06 ± 0.33, P=1
rh_rostralmiddlefrontal	0.54 ± 0.36, P=1	0.4 ± 0.12, P=0.067	-0.4 ± 0.3, P=1	-0.83 ± 0.5, P=1	-0.48 ± 0.16, P=0.193	-0.46 ± 0.22, P=1
rh_superiorfrontal	0.53 ± 0.41, P=1	0.5 ± 0.14, P=0.026	-0.16 ± 0.35, P=1	-1.09 ± 0.57, P=1	-0.64 ± 0.18, P=0.039	-0.29 ± 0.25, P=1
rh_superiorparietal	1.03 ± 0.36, P=0.293	0.4 ± 0.12, P=0.09	-0.4 ± 0.31, P=1	-0.98 ± 0.5, P=1	-0.67 ± 0.16, P=0.002	-0.79 ± 0.22, P=0.024

rh_superiortemporal	2.91 ± 0.51, P<0.001	1.66 ± 0.17, P<0.001	-3.05 ± 0.44, P<0.001	-5.6 ± 0.71, P<0.001	-1.19 ± 0.23, P<0.001	-0.91 ± 0.32, P=0.308
rh_supramarginal	1.93 ± 0.38, P<0.001	1.03 ± 0.13, P<0.001	-1.73 ± 0.33, P<0.001	-3.57 ± 0.53, P<0.001	-0.9 ± 0.17, P<0.001	-0.65 ± 0.24, P=0.413
rh_temporalpole	4.67 ± 1, P<0.001	2.66 ± 0.33, P<0.001	-4.91 ± 0.85, P<0.001	-8.98 ± 1.38, P<0.001	-1.86 ± 0.46, P=0.003	-1.82 ± 0.62, P=0.247
rh_transversetemporal	3.08 ± 0.57, P<0.001	1.39 ± 0.2, P<0.001	-2.36 ± 0.49, P<0.001	-5.33 ± 0.79, P<0.001	-0.79 ± 0.26, P=0.153	-0.86 ± 0.35, P=1
lh_MeanThickness	1.93 ± 0.33, P<0.001	0.7 ± 0.11, P<0.001	-1.22 ± 0.28, P=0.001	-2.17 ± 0.45, P<0.001	-0.72 ± 0.15, P<0.001	-0.47 ± 0.2, P=1
rh_MeanThickness	1.65 ± 0.33, P<0.001	0.78 ± 0.11, P<0.001	-1.22 ± 0.28, P=0.001	-2.51 ± 0.46, P<0.001	-0.76 ± 0.15, P<0.001	-0.64 ± 0.21, P=0.138

104
105 The estimates indicate the cortical thickness for that region preserved (positive values)
106 or lost (negative values) per year for each unit increase in the marker assessed at the
107 baseline. Units were 100 for WM-PVS count, 10 for BG-PVS count, 0.1 mm for WM-
108 and BG-PVS diameter, and 1 log for P-WML and D-WML volume. See also Figure 2B.
109 Parcellations performed by Freesurfer according to the Desikan-Killany atlas. Bankssts:
110 Banks of the Superior Temporal Sulcus; Lh: left hemisphere; Rh: right hemisphere.
111 Models were fully adjusted for all the same covariates included in the main linear mixed-
112 effect model. P values are corrected for 70 comparisons with Holm-Bonferroni
113 procedure.

114
115

116
117
118

Table S12. Regional estimates (mm³/year) and adjusted significance of the interaction term “marker” by “time” in linear mixed-effects models with the white matter volume of the corresponding region as dependent variable.

Region	WM-PVS count	BG-PVS count	WM-PVS diameter	BG-PVS diameter	P-WML log volume	D-WML log volume
lh_bankssts	3.41 ± 0.55, P<0.001	0.67 ± 0.17, P=0.005	-0.44 ± 0.44, P=1	-1.47 ± 0.71, P=1	-0.82 ± 0.23, P=0.028	-0.15 ± 0.32, P=1
lh_caudalanteriorcingulate	4.32 ± 0.85, P<0.001	1.24 ± 0.26, P<0.001	-0.35 ± 0.67, P=1	-2.62 ± 1.09, P=1	-0.94 ± 0.36, P=0.614	-0.99 ± 0.49, P=1
lh_caudalmiddlefrontal	7.82 ± 1.14, P<0.001	1.47 ± 0.36, P=0.003	1.15 ± 0.91, P=1	-0.77 ± 1.48, P=1	-1.06 ± 0.48, P=1	0.25 ± 0.66, P=1
lh_cuneus	6.73 ± 0.86, P<0.001	2.1 ± 0.27, P<0.001	-2.39 ± 0.68, P=0.031	-5.81 ± 1.11, P<0.001	-1.32 ± 0.36, P=0.017	-0.11 ± 0.49, P=1
lh_entorhinal	2.4 ± 0.37, P<0.001	0.57 ± 0.11, P<0.001	-0.87 ± 0.29, P=0.18	-1.94 ± 0.47, P=0.003	-0.49 ± 0.15, P=0.107	0.17 ± 0.21, P=1
lh_frontalpole	0.95 ± 0.21, P=0.001	0.21 ± 0.07, P=0.151	-0.43 ± 0.17, P=0.703	-0.4 ± 0.28, P=1	-0.11 ± 0.09, P=1	-0.15 ± 0.12, P=1
lh_fusiform	14.5 ± 1.5, P<0.001	3.25 ± 0.48, P<0.001	-2.96 ± 1.21, P=1	-8.63 ± 1.97, P=0.001	-2.47 ± 0.64, P=0.009	-0.28 ± 0.88, P=1
lh_inferiorparietal	14.3 ± 2.12, P<0.001	3.27 ± 0.67, P<0.001	-2.69 ± 1.69, P=1	-9.9 ± 2.74, P=0.021	-1.31 ± 0.9, P=1	-0.46 ± 1.22, P=1
lh_inferiortemporal	16.75 ± 1.65, P<0.001	3.75 ± 0.52, P<0.001	-5.95 ± 1.34, P=0.001	-12.98 ± 2.17, P<0.001	-2.48 ± 0.71, P=0.036	-0.75 ± 0.97, P=1
lh_insula	7.6 ± 1.39, P<0.001	2.08 ± 0.43, P<0.001	-2.42 ± 1.09, P=1	-4.98 ± 1.77, P=0.345	-3.05 ± 0.58, P<0.001	-1.51 ± 0.79, P=1
lh_isthmuscingulate	6.37 ± 0.71, P<0.001	1.59 ± 0.22, P<0.001	-0.79 ± 0.57, P=1	-4.27 ± 0.92, P<0.001	-1.37 ± 0.3, P<0.001	-0.15 ± 0.41, P=1
lh_lateraloccipital	25.09 ± 2.75, P<0.001	7.68 ± 0.87, P<0.001	-12.72 ± 2.18, P<0.001	-24.8 ± 3.57, P<0.001	-4.65 ± 1.17, P=0.005	-2.02 ± 1.59, P=1
lh_lateralorbitofrontal	11.53 ± 1.31, P<0.001	2.65 ± 0.41, P<0.001	-1.7 ± 1.05, P=1	-6.45 ± 1.71, P=0.011	-1.77 ± 0.56, P=0.107	-0.38 ± 0.76, P=1
lh_lingual	12.16 ± 1.53, P<0.001	3.51 ± 0.48, P<0.001	-3.92 ± 1.21, P=0.08	-9.81 ± 1.97, P<0.001	-2.31 ± 0.64, P=0.021	-0.01 ± 0.87, P=1
lh_medialorbitofrontal	6.51 ± 0.92, P<0.001	1.29 ± 0.29, P=0.001	1.52 ± 0.73, P=1	-1.57 ± 1.2, P=1	-0.74 ± 0.39, P=1	0.99 ± 0.53, P=1
lh_middletemporal	12.6 ± 1.5, P<0.001	3.33 ± 0.47, P<0.001	-4.78 ± 1.21, P=0.005	-10.63 ± 1.96, P<0.001	-1.28 ± 0.65, P=1	-0.37 ± 0.88, P=1
lh_paracentral	5.64 ± 0.98, P<0.001	0.72 ± 0.31, P=1	1.77 ± 0.77, P=1	0.61 ± 1.26, P=1	-0.14 ± 0.41, P=1	0.83 ± 0.56, P=1
lh parahippocampal	2.2 ± 0.42, P<0.001	0.67 ± 0.13, P<0.001	-1.2 ± 0.33, P=0.017	-1.94 ± 0.53, P=0.019	-0.33 ± 0.17, P=1	0.19 ± 0.24, P=1
lh_parsopercularis	4.69 ± 0.92, P<0.001	0.97 ± 0.29, P=0.05	0.52 ± 0.73, P=1	-1.1 ± 1.18, P=1	0.25 ± 0.39, P=1	0.57 ± 0.53, P=1
lh_parsorbitalis	2.12 ± 0.4, P<0.001	0.52 ± 0.13, P=0.003	-0.76 ± 0.31, P=1	-1.47 ± 0.51, P=0.292	-0.12 ± 0.17, P=1	0.07 ± 0.23, P=1
lh_parstriangularis	5.06 ± 0.77, P<0.001	0.91 ± 0.24, P=0.011	-1.25 ± 0.61, P=1	-2.51 ± 0.99, P=0.801	-0.49 ± 0.33, P=1	0.38 ± 0.44, P=1
lh_pericalcarine	5.53 ± 0.88, P<0.001	1.19 ± 0.28, P=0.001	-1.27 ± 0.69, P=1	-3.5 ± 1.13, P=0.134	-0.67 ± 0.37, P=1	0.45 ± 0.5, P=1
lh_postcentral	11.37 ± 1.95, P<0.001	1.84 ± 0.63, P=0.237	0.86 ± 1.54, P=1	-0.98 ± 2.53, P=1	0.05 ± 0.81, P=1	2.05 ± 1.1, P=1
lh_posteriorcingulate	8.58 ± 1.17, P<0.001	2.44 ± 0.35, P<0.001	-1.93 ± 0.93, P=1	-5.41 ± 1.51, P=0.023	-2.23 ± 0.5, P=0.001	-1.29 ± 0.68, P=1
lh_precentral	20.53 ± 3.05, P<0.001	3.42 ± 0.97, P=0.03	4.2 ± 2.43, P=1	0.56 ± 3.98, P=1	-2.78 ± 1.28, P=1	3.01 ± 1.75, P=1
lh_precuneus	12.86 ± 1.84, P<0.001	2.31 ± 0.58, P=0.005	1.41 ± 1.47, P=1	-4.67 ± 2.39, P=1	-1.9 ± 0.78, P=0.998	0.75 ± 1.06, P=1
lh_rostralanteriorcingulate	3.37 ± 0.52, P<0.001	0.82 ± 0.16, P<0.001	-0.04 ± 0.41, P=1	-1.84 ± 0.67, P=0.406	-0.45 ± 0.22, P=1	-0.47 ± 0.3, P=1
lh_rostralmiddlefrontal	16.78 ± 2.88, P<0.001	3.93 ± 0.9, P<0.001	-0.61 ± 2.29, P=1	-3.79 ± 3.74, P=1	-1.41 ± 1.22, P=1	0.62 ± 1.66, P=1
lh_superiorfrontal	23.88 ± 4.15, P<0.001	3.41 ± 1.31, P=0.619	6.11 ± 3.31, P=1	3.53 ± 5.4, P=1	-0.78 ± 1.76, P=1	2.37 ± 2.39, P=1
lh_superiorparietal	19.46 ± 2.67, P<0.001	4.75 ± 0.84, P<0.001	-1.73 ± 2.14, P=1	-9.56 ± 3.48, P=0.413	-3.14 ± 1.13, P=0.382	0.33 ± 1.55, P=1

lh_superiortemporal	15.38 ± 1.64, P<0.001	3.93 ± 0.52, P<0.001	-2.75 ± 1.32, P=1	-9.05 ± 2.15, P=0.002	-1.54 ± 0.7, P=1	0.23 ± 0.96, P=1
lh_supramarginal	13.78 ± 1.85, P<0.001	2.77 ± 0.59, P<0.001	-1.18 ± 1.49, P=1	-6.09 ± 2.43, P=0.836	-1.47 ± 0.79, P=1	0.72 ± 1.08, P=1
lh_temporalpole	2.75 ± 0.4, P<0.001	0.66 ± 0.12, P<0.001	-1.5 ± 0.31, P<0.001	-2.68 ± 0.51, P<0.001	-0.48 ± 0.17, P=0.274	-0.09 ± 0.23, P=1
lh_transversetemporal	0.42 ± 0.3, P=1	-0.06 ± 0.09, P=1	0.56 ± 0.23, P=1	0.54 ± 0.38, P=1	0.34 ± 0.12, P=0.361	0.19 ± 0.17, P=1
rh_bankssts	4.59 ± 0.57, P<0.001	1.05 ± 0.17, P<0.001	-1.08 ± 0.45, P=1	-1.93 ± 0.73, P=0.577	-0.88 ± 0.24, P=0.018	-0.34 ± 0.33, P=1
rh_caudalanteriorcingulate	4.03 ± 0.85, P<0.001	0.45 ± 0.26, P=1	-0.48 ± 0.67, P=1	0.03 ± 1.08, P=1	-0.49 ± 0.36, P=1	-0.61 ± 0.49, P=1
rh_caudalmiddlefrontal	6.12 ± 1.1, P<0.001	0.92 ± 0.35, P=0.546	1.04 ± 0.87, P=1	1.25 ± 1.43, P=1	-1.07 ± 0.46, P=1	0.26 ± 0.63, P=1
rh_cuneus	6.17 ± 0.83, P<0.001	1.8 ± 0.26, P<0.001	-0.64 ± 0.66, P=1	-3.39 ± 1.07, P=0.111	-1.34 ± 0.35, P=0.008	0.15 ± 0.47, P=1
rh_entorhinal	2.31 ± 0.34, P<0.001	0.53 ± 0.11, P<0.001	-0.75 ± 0.27, P=0.327	-1.42 ± 0.44, P=0.075	-0.38 ± 0.14, P=0.509	-0.22 ± 0.19, P=1
rh_frontalpole	1.53 ± 0.27, P<0.001	0.46 ± 0.09, P<0.001	-1.03 ± 0.21, P<0.001	-1.22 ± 0.35, P=0.04	-0.35 ± 0.11, P=0.173	-0.32 ± 0.16, P=1
rh_fusiform	14.68 ± 1.42, P<0.001	3.1 ± 0.45, P<0.001	-2.98 ± 1.16, P=0.702	-6.34 ± 1.89, P=0.055	-3.36 ± 0.61, P<0.001	-0.89 ± 0.84, P=1
rh_inferiorparietal	23.49 ± 2.55, P<0.001	5.05 ± 0.82, P<0.001	-7.48 ± 2.07, P=0.022	-14.86 ± 3.36, P=0.001	-3.08 ± 1.1, P=0.363	-0.63 ± 1.51, P=1
rh_inferiortemporal	15.48 ± 1.56, P<0.001	3.12 ± 0.5, P<0.001	-5.1 ± 1.28, P=0.005	-8.77 ± 2.07, P=0.002	-2.74 ± 0.68, P=0.004	-0.89 ± 0.93, P=1
rh_insula	4.55 ± 1.26, P=0.02	1.22 ± 0.39, P=0.113	-1.76 ± 0.98, P=1	-0.93 ± 1.6, P=1	-1.88 ± 0.52, P=0.021	-1.32 ± 0.71, P=1
rh_isthmuscingulate	4.04 ± 0.77, P<0.001	0.9 ± 0.24, P=0.012	0.1 ± 0.61, P=1	-1.14 ± 1, P=1	-1 ± 0.33, P=0.158	0.23 ± 0.45, P=1
rh_lateraloccipital	29.68 ± 2.68, P<0.001	7.59 ± 0.86, P<0.001	-11.84 ± 2.17, P<0.001	-22.17 ± 3.54, P<0.001	-5.87 ± 1.15, P<0.001	-1.17 ± 1.58, P=1
rh_lateralorbitofrontal	9.87 ± 1.3, P<0.001	2.07 ± 0.41, P<0.001	-0.11 ± 1.04, P=1	-2.06 ± 1.7, P=1	-2.23 ± 0.55, P=0.004	0.21 ± 0.76, P=1
rh_lingual	12.35 ± 1.49, P<0.001	3.97 ± 0.47, P<0.001	-2.66 ± 1.19, P=1	-8.88 ± 1.94, P<0.001	-2.66 ± 0.62, P<0.001	-1.09 ± 0.85, P=1
rh_medialorbitofrontal	8.24 ± 0.97, P<0.001	1.96 ± 0.31, P<0.001	-1.47 ± 0.78, P=1	-3.36 ± 1.27, P=0.565	-1.41 ± 0.41, P=0.043	0.3 ± 0.56, P=1
rh_middletemporal	15.73 ± 1.77, P<0.001	3.75 ± 0.56, P<0.001	-7.46 ± 1.43, P<0.001	-11.3 ± 2.32, P<0.001	-2.2 ± 0.77, P=0.284	-1.28 ± 1.05, P=1
rh_paracentral	5.82 ± 1.03, P<0.001	0.63 ± 0.33, P=1	2.2 ± 0.81, P=0.481	2 ± 1.33, P=1	-0.9 ± 0.43, P=1	0.79 ± 0.59, P=1
rh parahippocampal	2.83 ± 0.39, P<0.001	0.95 ± 0.12, P<0.001	-1.48 ± 0.31, P<0.001	-2.33 ± 0.5, P<0.001	-0.77 ± 0.16, P<0.001	-0.37 ± 0.22, P=1
rh_parsopercularis	4.98 ± 0.8, P<0.001	1.16 ± 0.25, P<0.001	-0.52 ± 0.64, P=1	-2.26 ± 1.03, P=1	-0.8 ± 0.34, P=1	0.08 ± 0.46, P=1
rh_parsorbitalis	2.97 ± 0.49, P<0.001	0.97 ± 0.16, P<0.001	-1.84 ± 0.39, P<0.001	-2.97 ± 0.64, P<0.001	-0.39 ± 0.21, P=1	-0.28 ± 0.28, P=1
rh_parstriangularis	5.64 ± 0.91, P<0.001	1.39 ± 0.28, P<0.001	-1.99 ± 0.72, P=0.414	-3.62 ± 1.18, P=0.147	-0.76 ± 0.39, P=1	0.01 ± 0.53, P=1
rh_pericalcarine	4.95 ± 0.79, P<0.001	1.27 ± 0.26, P<0.001	-0.71 ± 0.62, P=1	-2.16 ± 1.03, P=1	-0.82 ± 0.33, P=0.828	0.29 ± 0.45, P=1
rh_postcentral	12.42 ± 2.06, P<0.001	1.47 ± 0.66, P=1	0.01 ± 1.63, P=1	0.13 ± 2.68, P=1	-0.8 ± 0.86, P=1	0.85 ± 1.17, P=1
rh_posteriorcingulate	7.06 ± 1.03, P<0.001	1.8 ± 0.31, P<0.001	-0.97 ± 0.82, P=1	-2.26 ± 1.33, P=1	-2.04 ± 0.44, P<0.001	-0.73 ± 0.6, P=1
rh_precentral	19.91 ± 3.01, P<0.001	2.99 ± 0.96, P=0.127	3.08 ± 2.39, P=1	0.82 ± 3.92, P=1	-2.48 ± 1.26, P=1	1.67 ± 1.72, P=1
rh_precuneus	15.5 ± 1.73, P<0.001	3.21 ± 0.55, P<0.001	0.13 ± 1.4, P=1	-3.04 ± 2.29, P=1	-2.38 ± 0.74, P=0.095	0.45 ± 1.01, P=1
rh_rostralanteriorcingulate	2.69 ± 0.45, P<0.001	0.41 ± 0.14, P=0.251	-0.5 ± 0.36, P=1	-0.46 ± 0.58, P=1	-0.48 ± 0.19, P=0.776	-0.02 ± 0.26, P=1
rh_rostralmiddlefrontal	21.01 ± 3, P<0.001	5.78 ± 0.94, P<0.001	-6.93 ± 2.4, P=0.27	-13.51 ± 3.9, P=0.037	-4.3 ± 1.28, P=0.054	-1.29 ± 1.74, P=1
rh_superiorfrontal	26.24 ± 3.99, P<0.001	4.8 ± 1.26, P=0.01	0.5 ± 3.21, P=1	-0.54 ± 5.23, P=1	-4.27 ± 1.7, P=0.836	-0.08 ± 2.32, P=1
rh_superiorparietal	22.44 ± 2.39, P<0.001	5.36 ± 0.76, P<0.001	-4.3 ± 1.93, P=1	-12.29 ± 3.14, P=0.007	-3.84 ± 1.02, P=0.012	0.42 ± 1.39, P=1

rh_superiortemporal	14.03 ± 1.76, P<0.001	3.41 ± 0.55, P<0.001	-4.63 ± 1.42, P=0.078	-7.77 ± 2.31, P=0.055	-2.28 ± 0.76, P=0.181	-0.73 ± 1.04, P=1
rh_supramarginal	12.76 ± 1.73, P<0.001	2.71 ± 0.55, P<0.001	-2.73 ± 1.4, P=1	-5.7 ± 2.28, P=0.847	-2.22 ± 0.74, P=0.189	0.33 ± 1.01, P=1
rh_temporalpole	1.86 ± 0.34, P<0.001	0.49 ± 0.11, P<0.001	-0.82 ± 0.27, P=0.171	-1.43 ± 0.44, P=0.083	-0.51 ± 0.14, P=0.023	-0.3 ± 0.19, P=1
rh_transversetemporal	0.4 ± 0.26, P=1	0.06 ± 0.08, P=1	0.4 ± 0.2, P=1	0.08 ± 0.33, P=1	0.05 ± 0.11, P=1	-0.12 ± 0.15, P=1

119
120
121
122
123
124
125
126
127
128
129
130

The estimates indicate the white matter volume for that region preserved (positive values) or lost (negative values) per year for each unit increase in the marker assessed at the baseline. Units were 100 for WM-PVS count, 10 for BG-PVS count, 0.1 mm for WM- and BG-PVS diameter, and 1 log for P-WML and D-WML volume. See also Figure 2C. Parcellations performed by Freesurfer according to the Desikan-Killany atlas. Bankssts: Banks of the Superior Temporal Sulcus; Lh: left hemisphere; Rh: right hemisphere. Models were fully adjusted for all the same covariates included in the main linear mixed-effect model. P values are corrected for 68 comparisons with Holm-Bonferroni procedure.

131
132
133

Table S13. Sensitivity analysis on the interaction term “marker” by “time” in linear mixed-effects models with the grey matter volume, cortical thickness, or white matter volume as dependent variables.

Marker	Covariate	Grey matter volume (mm ³ / year)	Cortical thickness (µm / year)	White matter volume (mm ³ / year)
WM-PVS count (per 100-unit)	Artifacts	720±100, P<0.001	4±1, P<0.001	582±84, P<0.001
	APOE	696±99, P<0.001	4±1, P<0.001	605±84, P<0.001
	MMSE	812±104, P<0.001	4±1, P<0.001	645±88, P<0.001
	Tobacco	728±101, P<0.001	4±1, P<0.001	596±85, P<0.001
	Abeta	561±113, P<0.001	3±1, P<0.001	744±84, P<0.001
	Tau	413±112, P<0.001	3±1, P=0.001	550±79, P<0.001
BG-PVS count (per 10-unit)	Artifacts	190±33, P<0.001	1±0, P<0.001	143±26, P<0.001
	APOE	178±33, P<0.001	1±0, P<0.001	140±26, P<0.001
	MMSE	255±35, P<0.001	2±0, P<0.001	170±27, P<0.001
	Tobacco	190±33, P<0.001	1±0, P<0.001	142±26, P<0.001
	Abeta	161±35, P<0.001	1±0, P<0.001	151±25, P<0.001
	Tau	88±36, P=0.031	1±0, P<0.001	189±24, P<0.001
WM-PVS diameter (per 0.1 mm)	Artifacts	-374±84, P<0.001	-2±1, P<0.001	-64±67, P=0.671
	APOE	-341±83, P<0.001	-2±1, P<0.001	-71±67, P=0.57
	MMSE	-463±87, P<0.001	-3±1, P<0.001	-93±70, P=0.37
	Tobacco	-378±85, P<0.001	-3±1, P<0.001	-58±67, P=0.775
	Abeta	-251±90, P=0.011	-2±1, P=0.001	73±66, P=0.526
	Tau	-170±91, P=0.123	-2±1, P=0.022	-15±62, P=1
BG-PVS diameter (per 0.1 mm)	Artifacts	-571±136, P<0.001	-5±1, P<0.001	-282±108, P=0.019
	APOE	-543±136, P<0.001	-4±1, P<0.001	-274±108, P=0.023
	MMSE	-726±141, P<0.001	-6±1, P<0.001	-334±114, P=0.007
	Tobacco	-577±138, P<0.001	-5±1, P<0.001	-278±109, P=0.022
	Abeta	-338±146, P=0.041	-4±1, P<0.001	-202±106, P=0.112
	Tau	-177±145, P=0.444	-2±1, P=0.042	-311±98, P=0.003
P-WML log volume	Artifacts	-277±45, P<0.001	-1±0, P<0.001	-71±36, P=0.09
	APOE	-274±45, P<0.001	-1±0, P<0.001	-71±36, P=0.089
	MMSE	-283±48, P<0.001	-1±0, P<0.001	-81±38, P=0.066
	Tobacco	-285±46, P<0.001	-2±0, P<0.001	-68±36, P=0.118
	Abeta	-208±48, P<0.001	-1±0, P<0.001	-25±34, P=0.91
	Tau	-227±47, P<0.001	-2±0, P<0.001	-61±31, P=0.093
D-WML log volume	Artifacts	-185±61, P=0.005	-1±0, P=0.01	1±48, P=1
	APOE	-165±61, P=0.014	-1±0, P=0.028	7±48, P=1
	MMSE	-144±65, P=0.053	-1±0, P=0.067	2±52, P=1
	Tobacco	-191±62, P=0.004	-1±0, P=0.013	-1±49, P=1
	Abeta	-157±65, P=0.03	-1±0, P=0.052	69±46, P=0.259
	Tau	-151±62, P=0.031	-1±0, P=0.088	57±41, P=0.326

134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144

Estimates indicate the value of the dependent variable preserved (positive values) or lost (negative values) per year for each unit increase in the marker. All models were adjusted for the same covariates as in all the other main models. “P” indicates adjusted P-values.

Table S14. Estimated marginal means for baseline PVS and WML markers in non-demented individuals who converted to dementia versus non-converters.

Marker (at baseline MRI)	Estimated marginal mean ± standard error		Adj. P value
	Non-converters	Converters	
WM-PVS count	453 ± 7	415 ± 8	<0.001
BG-PVS count	143 ± 2	127 ± 2	<0.001
WM-PVS diameter (mm)	1.96 ± 0.01	1.98 ± 0.01	<0.001
BG-PVS diameter (mm)	1.58 ± 0	1.6 ± 0.01	<0.001
P-WML log volume	4.28 ± 0.15	5.22 ± 0.17	<0.001
D-WML log volume	2.5 ± 0.11	2.51 ± 0.13	1

145
 146
 147
 148
 149
 150
 151
 152
 153

Marginal means were estimated controlling for the same covariates as in the primary analysis (Table S2).

Table S15. Estimates and adjusted significance of the logistic regression models assessing the association of medium- and high-risk tertiles with amyloid and tau positive status controlling for demographic and clinical covariates.

Tertile	Marker	Amyloid-β + status	Tau + status
Medium-risk + High-risk (compared to Low-risk)	WM-PVS count	0.12 ± 0.1, P=0.489	0.01 ± 0.12, P=1
	BG-PVS count	0.18 ± 0.1, P=0.16	0.15 ± 0.13, P=0.444
	WM-PVS diameter	0.13 ± 0.1, P=0.352	0.17 ± 0.12, P=0.318
	BG-PVS diameter	0.15 ± 0.1, P=0.236	0.13 ± 0.12, P=0.57
	P-WML volume	0.27 ± 0.1, P=0.017	-0.24 ± 0.12, P=0.096
	D-WML volume	0.16 ± 0.09, P=0.193	-0.16 ± 0.11, P=0.33
	Total grey matter volume	0.28 ± 0.1, P=0.013	0.07 ± 0.13, P=1
	Mean cortical thickness	0.49 ± 0.11, P<0.001	0.17 ± 0.11, P=0.227
High-risk only (compared to Low-risk + Medium-risk)	WM-PVS count	0.01 ± 0.09, P=1	-0.04 ± 0.11, P=1
	BG-PVS count	0.12 ± 0.09, P=0.408	0.01 ± 0.12, P=1
	WM-PVS diameter	-0.05 ± 0.09, P=1	-0.17 ± 0.1, P=0.199
	BG-PVS diameter	0.07 ± 0.09, P=0.867	0.13 ± 0.12, P=0.57
	P-WML volume	0.03 ± 0.08, P=1	-0.07 ± 0.1, P=0.965
	D-WML volume	0.05 ± 0.08, P=1	-0.32 ± 0.1, P=0.003
	Total grey matter volume	0.3 ± 0.11, P=0.01	-0.14 ± 0.13, P=0.554
	Mean cortical thickness	0.25 ± 0.09, P=0.009	0.01 ± 0.12, P=1

154

155 **Table S16. Tertile limits for PVS and WML markers.**
 156

Marker	Low-risk tertile limits	Medium-risk tertile limits	High-risk tertile limits
WM-PVS count	840 to 498	497 to 377	376 to 86
BG-PVS count	272 to 155	154 to 117	116 to 34
WM-PVS diameter (mm)	1.69 to 1.9	1.91 to 2.03	2.04 to 2.37
BG-PVS diameter (mm)	1.37 to 1.53	1.54 to 1.61	1.62 to 1.83
P-WML volume (log)	0 to 2.83	2.84 to 6.74	6.75 to 9.56
D-WML volume (log)	0 to 1.1	1.11 to 3.81	3.82 to 6.7

157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168

Table S17. Baseline characteristics of the participants whose brain MRI data have been used for the robust vessel segmentation method development and validation.

Dataset	Cohort	N. subjects	Age – yr	Female – %	Body Mass Index – kg/m ²
Test-Retest	MarkVCID	39	71.6 ± 10.1	69.2	NA
Inter-scanner	MarkVCID	19	68.5 ± 9.3	52.6	NA
Inter-field-strength	ADNI	115	74.6 ± 6.6	51.3	26.3 ± 4.5
Biological validation	HCP	2163	33.1 ± 17.8	55.4	25.4 ± 5.6*

*Body Mass index was available in 2160 subjects.

169
 170
 171
 172
 173
 174
 175
 176

177 **Table S18. Overview of the MRI scanners and T1-weighted image parameters**
 178 **employed to acquire the data used for the robust vessel segmentation method**
 179 **development and validation.**

FS (T)	Manuf.	Model	TR (sec)	TE (msec)	TI (sec)	FA (°)	Native voxel vol. (mm ³)	N. scans analyzed	Cohort
Test-retest dataset									
3	Philips	Achieva dStream	0.0095	1.8 / 3.7 / 5.5 / 7.3	NA	7	1	12	MarkVCID
3	Siemens	Prisma	2.53	1.7 / 3.6 / 5.4 / 7.3	1.1	7	1	12	MarkVCID
3	Siemens	Prisma fit	2.53	1.7 / 3.6 / 5.4 / 7.3	1.1	7	1	40	MarkVCID
3	Siemens	TrioTim	2.53	1.6 / 3.5 / 5.3 / 7.2	1.2	7	1	14	MarkVCID
Inter-scanner dataset									
3	GE	DISCOVERY MR750w	0.0095	3.7	0.6	10	0.5	19	MarkVCID
3	Philips	Achieva dStream	0.0095	1.8 / 3.7 / 5.5 / 7.3	NA	7	1	19	MarkVCID
3	Siemens	Prisma fit	2.53	1.7 / 3.6 / 5.4 / 7.3	1.1	7	1	19	MarkVCID
3	Siemens	TrioTim	2.53	1.6 / 3.5 / 5.3 / 7.2	1.2	7	1	19	MarkVCID
Inter-field-strength dataset									
1.5	GE	GENESIS SIGNA	0.01	4.1	1	8	1.1	47	ADNI
1.5	GE	SIGNA EXCITE	0.0086 to 0.0092	3.8 to 4.1	1	8	1.1	93	ADNI
1.5	GE	SIGNA HDx	0.0086 to 0.0088	3.8 to 3.9	1	8	1.1	30	ADNI
1.5	GE	Signa HDxt	0.0086	3.8	1	8	1.1	4	ADNI
1.5	Philips	Achieva	0.0086	4	NA	8	1.1	8	ADNI
1.5	Philips	Gyrosan NT	0.0086	4	1	8	1.1	5	ADNI
1.5	Philips	Intera	0.0085 to 0.0086	4	1	8	1.1	34	ADNI
1.5	Siemens	Avanto	2.4	3.5	1	8	1.9	6	ADNI
1.5	Siemens	Espreo	2.4	3.6	1	8	1.9	3	ADNI
1.5	Siemens	Sonata	3	3.5	1	8	1.9	9	ADNI
1.5	Siemens	Symphony	3	3.9	1	8	1.9	13	ADNI
1.5	Siemens	Symphony Tim	3	3.6	1	8	1.9	47	ADNI
3	GE	GENESIS SIGNA	0.0075	3.1	0.9	8	1.2	9	ADNI
3	GE	SIGNA EXCITE	0.0066 to 0.0070	2.8 to 2.9	0.9	8	1.2	15	ADNI
3	GE	SIGNA HDx	0.0066	2.8	0.9	8	1.2	20	ADNI
3	GE	Signa HDxt	0.0066	2.8	0.9	8	1.2	1	ADNI
3	Philips	Achieva	0.0066 to 0.0069	3.1 to 3.2	0.85	8	1.2	47	ADNI
3	Philips	Intera	0.0067 to 0.007	3.1 to 3.3	0.85	8	1.2	61	ADNI
3	Siemens	Allegra	2.3	2.9	0.9	9	1.2	7	ADNI
3	Siemens	Trio	2.3	2.9	0.9	9	1.2	90	ADNI
3	Siemens	TrioTim	2.3	2.9	0.9	9	1.2	49	ADNI
Biological validation dataset									
3	Siemens	Prisma	2.5	1.8 / 3.6 / 5.4 / 7.2	1	8	0.512	1096	HCPA/HCPD
3	Siemens	Connectome Skyra	2.4	2.1	1	8	0.343	1067	HCPYA