

Supporting Information for

Detection of hemagglutinin H5 influenza A virus sequence in municipal wastewater solids at wastewater treatment plants with increases in influenza A in spring, 2024

Marlene K. Wolfe¹, Dorothea Duong², Bridgette Shelden², Elana M. G. Chan³, Vikram Chan-Herur², Stephen Hilton¹, Abigail Harvey Paulos¹, Alessandro Zulli³, Bradley J. White², Alexandria B. Boehm^{3*}

1. Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University, 1518 Clifton Rd, Atlanta, GA, USA, 30322
2. Verily Life Sciences LLC, South San Francisco, CA, USA
3. Department of Civil and Environmental Engineering, Stanford University, 473 Via Ortega, Stanford, California, 94305

*Corresponding author: aboehm@stanford.edu, 650-724-9128

Additional details related to EMMI guidelines. The 78 wastewater samples from the study that were used for measuring H5 were selected at random for this analysis. The results for the samples from the prospective study are represented by results reported by Boehm et al.¹.

The average (standard deviation) number of partitions (droplets) for the across 6 replicate wells was 93902 (21281) for the multiplex reaction. The volume of the partitions, as reported by the machine vendor is 0.00085 μ L. The average (standard deviation) of copies per partition for the H5 target is 5.0×10^{-4} (1.4×10^{-3}), and the average (standard deviation) of copies per partition for the IAV M gene target is 9.7×10^{-4} (1.9×10^{-3}). An example fluorescent plot from the QX600 (6 color reader) is included in the Stanford Digital Repository with the deposited wastewater data (<https://purl.stanford.edu/tr782wk3364>).

Figure S1. Location of the primers and probe within a reference genome (NC_007362). Within the HA segment of *A/goose/Guangdong/1/1996*, (<https://www.ncbi.nlm.nih.gov/datasets/taxonomy/93838/>), the amplicon is located between bases 1086-1256. IAV_H5_P1 is the probe, IAV_H5_F1 and IAV_H5_R3 are the forward and reverse primers, respectively.




Figure S2. Concentrations of IAV M gene (units of copies per gram dry weight) measured at four plants since late August 2023. These plants were chosen because they show a clear sharp increase in IAV M gene concentrations after the typical influenza season period between November and February. Symbols represent measured concentrations of samples, and lines represent the 5-adjacent sample, centered, trimmed averages. Errors on the measurements are standard deviations. Any point at the top of a plot shown without error bars represents samples with concentrations larger than that axis limit. Vertical lines indicate key dates in the avian influenza outbreak. February 8, 2024 USDA declares ongoing HPAI poultry outbreak, March 7, 2024: Unknown dairy cattle illness first reported, March 20, 2024: Samples collected from dairy cattle in Texas, March 25, 2024: Texas confirms H5N1 in dairy cattle, March 28, 2024: Positive H5N1 specimen collected from human.




Figure S3. Histogram of IAV M gene concentrations from prospective monitoring described in the paper. The dashed lines indicate the maximum values of IAV M gene measurements made at the three WWTPs; these maximums occurred in March or April 2024.




Table S1. Primers and probes developed to target the H5 version of the HA gene in influenza. The probe contained the fluorescent molecule fluorescein amidite (FAM) as well as ZEN, a proprietary internal quencher from IDT; and IBFQ, Iowa Black FQ.

Forward primer	TATAGARGGAGGATGGCAGG
Reverse primer	ACDGCCTCAAAYTGAGTGTT
Probe	AGGGGAGTGGKTACGCTGCRGAC

Table S2. Panel of pathogens used for in vitro specificity testing. All are inactivated pathogens from Zeptomatrix (Buffalo, NY) unless “Twist” appears in front of the name in which case it is nucleic acid purchased from Twist (South San Francisco, CA).

Parainfluenza 1	Adenovirus 31	Coronavirus NL63
Parainfluenza 2	Rhinovirus Type 1A	Coronavirus OC43
Parainfluenza 3	RSV A	<i>B. parapertussis</i>
Parainfluenza 4	RSV B	<i>B. pertussis</i>
Influenza A H1N1	SARS-CoV-2	Twist Influenza B
Influenza AH1	<i>M. pneumoniae</i>	Twist Influenza A H1N1
Influenza AH3	<i>C. pneumoniae</i>	Twist Influenza A H3N2
Influenza B	Metapneumovirus 8	
Adenovirus 1	Coronavirus HKU-1	
Adenovirus 3	Coronavirus 229E	

Table S3. Sample date ranges for the three WWTPs. Dates are in month/day/year format.

	Potter/Randall County, TX (North)	Potter/Randall County, TX (North)	Dallas County, TX
Date range	2/4/24 - 4/14/24	2/4/24 - 4/14/24	2/29/24 - 4/16/24

Table S4. WWTPs with secondary IAV wastewater event onsets or increases (identified visually) in late March and early April, 2024.

WWTPs with secondary springtime onset events	WWTPs still in seasonal onset with, visually identified recent increases
<p>Bridgewater, NJ Cahaba River, Birmingham, AL Chattanooga, TN East Bank, New Orleans, LA Eastern, Orange County, FL Esparto, CA Fremont, CA Gainesville, TX Gautier, MS Gilroy, CA Kailua, Honolulu, HI Kinston, NC Lancaster, CA Las Gallinas, San Rafael, CA Little Falls Run, Stafford, VA Lompoc, CA Marina, CA Modesto, CA Muscatine, IA Newark, CA Northwest, Orange County, FL Oakhurst, NJ P20, Kansas City, KS Pascagoula Moss Point, MS Petaluma, CA Regional, Laguna Niguel, CA Riverside, CA San Mateo, CA Seaford, DE Snohomish, WA South River, Atlanta, GA South, Laredo, TX Sunnyvale, TX University Park, PA Village Creek, Birmingham, AL Wahiawa, Honolulu, HI West Bank, New Orleans, LA West Contra Costa County, CA West Railroad, San Rafael, CA Wheeling, WV Wichita Falls, TX Windsor, CA Woodland, CA Zacate Creek, Laredo, TX</p>	<p>Bessemer, AL Dallas Central, Dallas, TX Hollywood Road, Amarillo, TX Jenison, MI Louisville, KY Mill Valley, CA Montpelier, VT Sausalito, CA River Road, Amarillo, TX Salina, KS South Bend, IN Southside, Dallas, TX St. Cloud, MN White Rock Central, Dallas, TX Wolcott, Kansas City, KS</p>

References

- (1) Boehm, A. B.; Wolfe, M. K.; Wigginton, K. R.; Bidwell, A.; White, B. J.; Hughes, B.; Duong; Chan-Herur, V.; Bischel, H. N.; Naughton, C. C. Human Viral Nucleic Acids Concentrations in Wastewater Solids from Central and Coastal California USA. *Scientific Data* **2023**, *10*, 396.