Title

Clostridioides difficile transmission: a compartmental model accounting for environmental spore persistence

Authors

Jon Edman-Wallér (1); MD, Kristina Rizzardi (2); PhD, Gunnar Jacobsson (1, 3); MD, PhD, associate professor; and Philip Gerlee (1, 4, 5), PhD, associate professor

Author affiliations

1. Centre for Antibiotic Resistance Research (CARe), Department of Infectious Diseases,

Institute of Biomedicine, Sahlgrenska Academy, University of Gothenburg

- 2. Department of Microbiology, Public Health Agency of Sweden, Solna, Sweden
- 3. Department of Infectious Diseases, Skaraborg Hospital, Skövde, Sweden
- 4. Department of Mathematical Sciences, Chalmers University of Technology, Gothenburg
- 5. Department of Mathematical Sciences, University of Gothenburg, Gothenburg

Corresponding author

Jon Edman Wallér, Postmästaregatan 10 C, 431 66 Mölndal, Sweden. Telephone:

+46720044676, E-mail: jon.edman@vgregion.se

Previous presentations

Oral presentations of preliminary results have been given at two Swedish and Nordic Infection Prevention and Control conferences during 2023.

Abbreviated title

C. difficile: a compartmental model Note This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Word count

2693 words.

Keywords: Clostridioides difficile; mathematical modeling; compartmental model;

healthcare-associated infections

Abstract

Objective: To set up a compartmental model, including environmental spore levels, for *C*. *difficile* transmission dynamics in a hospital and determine the effect of preventive interventions on infection and colonization rates.

Design: Intervention study within a compartmental mathematical modeling framework.

Setting: A simulated Swedish 500-bed secondary care hospital.

Interventions modeled: Antibiotic stewardship, improved isolation of infected patients, improved general cleaning and disinfection.

Results: Antibiotic stewardship had the largest effect on infections, with a 30.6% decrease in infection prevalence. Improved general cleaning and disinfection had the largest effect on colonization (-22.5%) and environmental spore levels (-39.7%). Improved isolation of infected patients had modest effects in comparison.

Conclusions: Modeling that includes the dynamics of environmental spores can aid our understanding of *C. difficile* transmission within hospitals. Antibiotic stewardship and improved general cleaning and disinfection showed the largest potential for prevention in our modeled setting.

Introduction

Clostridioides difficile infections (CDI) are among the most common healthcare-associated infections worldwide. Forming spores resistant to ordinarily used disinfectants, the bacteria can spread from infected or asymptomatically colonized patients to the hospital environment and indirectly to new patients. Hospitals harbor both colonized or infected patients at risk of spreading *C. difficile* and patients at risk of acquiring *C. difficile* during their hospital stay. However, in the absence of epidemic strains, only a minority of infections can be traced to transmission from other symptomatic patients. ^{1, 2} Other possible sources of transmission include (but are not limited to) environmental exposition in the community, transmission via healthcare workers (HCW), and transmission from asymptomatic carriers. Interventions to reduce the rate of CDI may be costly and, in some cases, of questionable impact.

C. difficile transmission dynamics have been modeled previously, with various versions of adapted compartmental models based on the classic Susceptible, Infected, Recovered (SIR) model, ³⁻⁵ or agent-based models. ⁶⁻⁸ SIR models often assume a rather direct person-to-person transmission, which does not describe the dynamics of *C. difficile* well. Spores from infected or colonized patients are deposited in the hospital environment, where they accumulate if not removed, and can later infect a new patient. Without specific modifications, an SIR model does not model the gradual increase of infections if cleaning and disinfection routines are lacking. When transmission from the environment has been included in previous compartmental models, it has been modeled as a constant transmission source. ⁹ Some agent-based modeling studies have incorporated environmental transmission, ^{6, 8} but include many assumptions of probabilities that are mainly based on expert opinion. In this study, we present a simple, modified compartmental model encompassing a separate compartment for the environmental spore reservoir. We then determine the modeled effects of several possible

interventions for reducing hospital CDI incidence, colonization, and environmental spore contamination.

Methods

Setting

The hospital in the model was assumed to be a 500-bed secondary-care hospital in Sweden. The model used aggregated compartments with assumptions of homogeneity. Input data from sources similar to our setting were preferred. Sweden has a low epidemic ribotype 027 (RT027) prevalence and a high diversity of strains. ¹⁰ The number of hospital beds is among the lowest in the OECD. ¹¹ Antibiotic consumption is low, ¹² as is the prevalence and burden of antimicrobial resistance. ¹³ *C. difficile* outbreaks have occurred infrequently, and incidence has declined in recent years. ¹⁰ Still, CDI is a common healthcare-related infection with an incidence of 61 cases per 100,000 in 2021. ¹⁴ Routines for infection control of CDI patients vary within the country but generally include care in a single room with a private bathroom until two days after symptom resolution; hand-washing for healthcare workers followed by alcoholic hand rub after contact with the patient or their surroundings; personal protective equipment based on contact precautions; daily disinfection of patient-near surfaces with a disinfectant with or without sporicidal effect; and thorough cleaning of the room at discharge.

Model

The model was adapted from the work by Yakob et al. ³ with the following changes: Firstly, we disregarded the exposed compartment, into which unexposed patients enter upon contact with *C. difficile* spores, and which they leave for the colonized compartment after a predefined period. In Yakob et al., this compartment played an essential role since they investigated the effect of CDI screening upon hospital admission, a procedure that can only detect colonized patients. The interventions we consider do not rely on the exposed/colonized

distinction, and we could let patients flow directly from unexposed to colonized. Secondly, we did not consider separate compartments for patients who recently have taken antimicrobials and thus are vulnerable. This was motivated by the fact that while antibiotics increase the risk of CDI, antibiotic treatment is not a prerequisite for infection. Lastly, to more accurately capture the fact that transmission occurs via the spores in the environment, we included an explicit compartment for environmental spores. This environment should be interpreted broadly and includes, for instance, the hands of healthcare workers (HCWs). HCWs were not modeled as separate compartments as they are infrequently infected or colonized. ¹⁵ We thus ended up with the following compartments in our model: patients unexposed to CDI (U), patients that have become colonized with *C. difficile* spores (C), patients that are infected (I), and those that are in recovery from an infection (R). Additionally, there is an environmental compartment (E) with spores instead of patients.

We consider a hospital with a fixed number of beds (N). The flow of newly admitted patients occurs into the U, C, and I compartments, where the proportion admitted into each compartment is controlled by the parameters ε_U , ε_C , and ε_I . The outflow of patients occurs from the U, C, and R compartments, is equal for the three compartments and given by κ . Spores in the environment are produced by colonized, infected, and recovered patients at rates α_C , α_I , and α_R , respectively, which increase the spore reservoir level E. Cleaning of the hospital environment reduces the number of spores at a rate δ . The number of new colonizations per time unit depends on the presence of unexposed patients and spores in the environment. It is controlled by the parameter β and the environmental spore reservoir E. The variable E is given in a non-dimensional normalized form, where the spore level at equilibrium in the baseline scenario equals one (see below). Colonized patients become infected at a rate θ , and infected patients become recovered at a rate ρ . No recurrences occur

within the hospital, so recovery is the last possible stage for patients before they are discharged.

The following system of coupled ordinary differential equations describes the model:

$$\frac{dU}{dt} = \varepsilon_U F - \beta EU - \kappa U$$
$$\frac{dC}{dt} = \varepsilon_C F + \beta EU - \theta C - \kappa C,$$
$$\frac{dI}{dt} = \varepsilon_I F + \theta C - \rho I,$$
$$\frac{dR}{dt} = \rho I - \kappa R,$$
$$\frac{dE}{dt} = \alpha_C C + \alpha_I I + \alpha_R R - \delta E,$$

where $F = \kappa(N - I)$ is the outflow of patients, and N = U + C + I + R is the total number of patients. A schematic description is given in Figure 1, which also shows the flow of patients between the compartments and the flow of spores to and from the environmental compartment E. The model was solved numerically using the Isoda-solver as implemented in the SciPy-method odeint. The initial condition was set to (U,C,I,R,E)(t = 0) = (0.95N,0.05N,0,0,0) and, to determine the equilibrium state, the model was solved for 1000 days. We tried a range of initial conditions and found that the equilibrium was insensitive to the exact choice of initial conditions. The code is available at https://github.com/philipgerlee/CDITransmission.

Determination of parameter values

First, a literature search was performed for each model parameter to find evidence for a range of plausible values. The values of α_C , α_I and α_R , representing the contribution of spores from individual patients per day to the total environmental reservoir of spores, were difficult to

estimate based on the literature and were only estimated relative to each other at this stage. Second, after setting arbitrary values for the α parameters and mid-range values for all other parameters, the model was solved numerically, and the proportion of colonized and infected patients in equilibrium was determined. Third, these values were compared to expected colonization levels (5-13.5%, see Table 1) and CDI prevalence (0.2-0.5%, ^{16, 17}, own data) among hospitalized patients in our setting. When the values were not within these ranges, the parameters were changed manually within their respective range until the model resulted in colonization and CDI prevalence within the expected ranges. The α parameters were adjusted first, and then the other parameters. Fourth, for the model to be internally consistent at baseline, βE (in the model) should be equal to our estimation of β at baseline, i.e., E should be = 1. This is because β is estimated based on studies close to our setting, with a certain level of environmental spores. Thus, the estimation of β is actually an estimation of β E. Parameter values, especially the α values (contribution to environmental spores by patients) and the δ value (clearance of spores from the environment), were adjusted within their ranges until the model resulted in E = 1 at equilibrium as well as expected levels of colonization and infection. Parameters, parameter ranges, final baseline model parameters, and outputs in the equilibrium state are presented in Table 1.

Interventions

Having built a model representing our current setting, we simulated three possible CDIreducing interventions to estimate their expected effect on the environmental spore burden and the prevalence of *C. difficile* infection and colonization. The effects of the interventions were studied on the scale from no change to very ambitious while still within the possible.

Intervention 1: Antibiotic stewardship program

A recently updated meta-analysis ³² of the increased risk for CDI in hospitalized patients for different classes of antibiotics was used to estimate the odds ratios (ORs) for different antibiotic classes. The point estimates of ORs in the study were used except when confidence intervals overlapped 1; in that case, the OR was set to 1. Swedish data on hospital antibiotic use in 2021 ³³ in defined daily doses (DDD) for each category was multiplied by OR - 1 to estimate the increase of CDI cases attributable to the antibiotic classes combined (table 2). Estimates of the increased CDI risk attributable to different antibiotic classes combined with Swedish data on hospital consumption of antibiotics (Table 2) resulted in an estimation that antibiotics with an OR >1 increase the number of cases by 44.1%, implying that $\frac{0.441}{1 + 0.441}$ = 30.6% of cases could be prevented if low-risk antibiotics or no antibiotics replaced high-risk antibiotics. To model this intervention, we varied the θ parameter from 0.004 to as low as 0.00233 to achieve an up to 30.6% decrease in the infected population I at equilibrium.

Intervention 2: Enhanced efforts on reducing the spread from infected patients to the environment.

For this intervention, we imagined that the transmission of spores from infected patients would be virtually eliminated. For instance, infected patients could be promptly moved to a separate building not part of the main hospital after diagnosis. We modeled this as a reduction of α_I up to 90%, from 0.001 to 0.0001, with all other variables being constant.

Intervention 3: General increased removal of spores

In the baseline model, spores are removed at a rate $\delta = 0.0175$. This represents the total effect of cleaning, disinfection, hand hygiene, etc., that remove spores from the hospital environment and thus prevents further transmission. While our model cannot determine the individual contribution of each of these practices for spore removal, we can simulate an intervention aimed at various hygienic measures, where the removal of spores increases by a certain amount. We modeled this by increasing δ by up to 30%, from 0.0175 to 0.02275.

Intervention 4: Bundled intervention

To study the effect of an ambitious bundled intervention with maximal effect aimed at antibiotics and isolation of infected patients as well as general cleaning and disinfection, we combined the three interventions above at their maximal effect.

Results

Intervention 1: Antibiotic stewardship program

This intervention resulted in a maximal decrease of colonized patients from 12.1% to 11.8% (-2.5%), a maximal decrease in disease prevalence from 0.43% to 0.30% (-30.6%), and a maximal reduction of environmental spores from 0.998 to 0.936 (-6.2%) (Figure 2A).

Intervention 2: Enhanced efforts on reducing the spread from infected patients to the environment

This intervention resulted in a maximal decrease of colonized patients from 12.1% to 10.7% (-12.2%), a maximal decrease of disease prevalence from 0.43% to 0.39% (-8.6%), and a maximal reduction of environmental spores from 0.998 to 0.781 (-21.8%) (Figure 2B).

Intervention 3: General increased removal of spores

This intervention resulted in a maximal decrease of colonized patients from 12.1% to 9.4% (– 22.5%), a maximal decrease in disease prevalence from 0.43% to 0.36% (–16.0%), and a maximal reduction of environmental spores from 0.998 to 0.601 (–39.7%) (Figure 2C).

Intervention 4: Bundled intervention

A combination of all three interventions above at their maximum effect resulted in a decrease in colonized patients from 12.1% to 8.6% (-28.9%), a decrease in disease prevalence from 0.43% to 0.25% (-41.5%), and a reduction of environmental spores from 0.998 to 0.484 (-51.5%) (not included in Figure).

Discussion

In this study, we aimed to model the transmission dynamics of *C. difficile* infections in a Swedish hospital setting. We modeled three intervention strategies and found that the antibiotic stewardship program had the largest impact on the prevalence of CDI. The impact on environmental spore levels and the prevalence of colonized patients was, however, modest. Vigorous isolation of infected patients moderately decreased colonization rates and environmental spore levels, while the effect on CDI prevalence was modest. Increased reduction of spores by overall cleaning and disinfection greatly reduced colonization rates and environmental spore levels, and moderately decreased CDI prevalence.

The maximum effect of an antibiotic stewardship program, a decrease in cases by 30.6%, might seem on the low side in relation to epidemiological studies on the effects of such programs, which often have achieved CDI incidence decreases of around 50%. ³⁴ However, Sweden has a favorable antimicrobial resistance situation, and antimicrobial stewardship programs are already active on national, regional, and local levels. For instance, beta-lactamase-sensitive penicillins with low CDI risk make up almost 10% of the total, almost as much as cephalosporins. ³³ Thus, it should be expected that the maximal impact of further stewardship measures in our setting is limited. Also, an antibiotic stewardship program might affect the progression from unexposed to colonized or lead to changes in lengths of stay,

which were not modeled in this study. However, different antibiotics may increase or decrease the risk of colonization, ²⁰ and antibiotics overall did not increase the risk of *C. difficile* colonization in a recent meta-analysis. ³⁵

The maximum effect of general cleaning and disinfection was considerable in our study. This contrasts with a recent review and meta-analysis of the effect of environmental cleaning bundles ³⁶, which concluded that such interventions seem to have little or no effect on CDI incidence. However, the bundles in the included studies may have been ineffective in increasing the overall reduction rate of environmental spores. Thus, our model implies that if we could substantially increase the reduction rate of spores, this would significantly affect CDI incidence. However, still needs to be determined what interventions would be needed to increase this rate substantially.

Focusing on isolating patients had modest effects in our model, especially on disease prevalence. In this scenario, colonized patients continue contributing spores to the environmental reservoir. Asymptomatic patients as important sources of *C. difficile* infections have been discussed in the literature in recent years. ³⁷⁻³⁹ While testing for colonization and isolating these patients may be practically unfeasible and ethically questionable, our results suggest that more effective cleaning and disinfection practices for all patients may be a way to reduce the risk of transmission from colonized patients; this, however, depends on these practices to increase the riddance of spores to a meaningful degree.

A main strength of our study is that the dynamic effects of the environmental spore reservoir are modeled in a way that, as far as we know, has not been performed previously. We have also calibrated the inputs to ensure that the model produces output that is plausible for our chosen setting and, as far as possible, based our assumptions on scientific data rather than expert opinion. While this study was performed with a Swedish hospital in mind, the model could easily be adjusted to mirror other settings.

Our study has limitations. No model is better than its assumptions, and for some parameters (e.g., the relations between the different α parameters), it was difficult to find clear scientific support for a given level. The model is also a highly simplified version of a hospital. It does not consider movements of patients within the hospital building, division into different wards, or movement of HCWs. The interventions were modeled in a simplified way as well. These shortcomings could be addressed by considering a more complex, potentially agent-based model, but such an approach requires detailed knowledge about model parameters.

We conclude that, according to our model, antibiotic stewardship measures have the greatest potential to decrease CDI incidence in our setting. Cleaning and disinfection efforts, implemented in a way that substantially decreases the environmental spore reservoir, have the greatest potential to prevent in-hospital acquisition of *C. difficile*.

Acknowledgements

Financial support. None reported.

Potential conflicts of interest. All authors report no conflicts of interest relevant to this article.

References

 Eyre DW, Cule ML, Wilson DJ, Griffiths D, Vaughan A, O'Connor L, et al.
 Diverse sources of *C. difficile* infection identified on whole-genome sequencing. *N Engl J Med.* 2013;369(13):1195-205.

2. Edman-Wallér JE, Toepfer M, Karp J, Rizzardi K, Jacobsson G, Werner M. *Clostridioides difficile* outbreak detection: Evaluation by ribotyping and whole-genome sequencing of a surveillance algorithm based on ward-specific cutoffs. *Infect Control Hosp Epidemiol.* 2023:1-5.

3. Yakob L, Riley TV, Paterson DL, Clements AC. *Clostridium difficile* exposure as an insidious source of infection in healthcare settings: an epidemiological model. *BMC Infect Dis.* 2013;13:376.

4. Starr JM, Campbell A, Renshaw E, Poxton IR, Gibson GJ. Spatio-temporal stochastic modelling of *Clostridium difficile*. *J Hosp Infect*. 2009;71(1):49-56.

5. Lofgren ET, Moehring RW, Anderson DJ, Weber DJ, Fefferman NH. A mathematical model to evaluate the routine use of fecal microbiota transplantation to prevent incident and recurrent *Clostridium difficile* infection. *Infect Control Hosp Epidemiol*. 2014;35(1):18-27.

6. Rubin MA, Jones M, Leecaster M, Khader K, Ray W, Huttner A, et al. A simulation-based assessment of strategies to control *Clostridium difficile* transmission and infection. *PLoS One*. 2013;8(11):e80671.

7. Lanzas C, Dubberke ER. Effectiveness of screening hospital admissions to detect asymptomatic carriers of *Clostridium difficile*: a modeling evaluation. *Infect Control Hosp Epidemiol*. 2014;35(8):1043-50.

8. Codella J, Safdar N, Heffernan R, Alagoz O. An agent-based simulation model for *Clostridium difficile* infection control. *Med Decis Making*. 2015;35(2):211-29.

9. Gingras G, Guertin MH, Laprise JF, Drolet M, Brisson M. Mathematical Modeling of the Transmission Dynamics of *Clostridium difficile* Infection and Colonization in Healthcare Settings: A Systematic Review. *PLoS One*. 2016;11(9):e0163880.

Rizzardi K, Norén T, Aspevall O, Mäkitalo B, Toepfer M, Johansson Å, et al.
 National Surveillance for *Clostridioides difficile* Infection, Sweden, 2009-2016. *Emerg Infect Dis*. 2018;24(9):1617-25.

11. Organisation for Economic Co-operation and Development. Hospital beds. 2021[Available from: https://data.oecd.org/healtheqt/hospital-beds.htm].

12. Eureopean Centre for Disease Control. Antimicrobial consumption in the EU/EEA (ESAC-Net). 2021 [Available from:

https://www.ecdc.europa.eu/sites/default/files/documents/ESAC-Net_AER_2021_final-rev.pdf].

European Centre for Disease Control. Assessing the health burden of infections
with antibiotic-resistant bacteria in the EU/EEA, 2016-2020. 2022 [Available from:
https://www.ecdc.europa.eu/sites/default/files/documents/Health-burden-infections-antibiotic-resistant-bacteria.pdf].

Public Health Agency of Sweden. *Clostridioides difficile*-infektion –sjukdomsstatistik. 2022 [Available from:

https://www.folkhalsomyndigheten.se/folkhalsorapportering-statistik/statistik-ao/sjukdomsstatistik/clostridium-difficile-infektion/?t=county].

15. Friedman ND, Pollard J, Stupart D, Knight DR, Khajehnoori M, Davey EK, et al. Prevalence of *Clostridium difficile* colonization among healthcare workers. *BMC Infect Dis.* 2013;13:459.

16. The Swedish Association of Local Authorities and Regions. Vårdrelaterade infektioner. 2019 [Available from:

https://skr.se/download/18.5627773817e39e979ef38d91/1642167986844/7585-756-5.pdf].

European Centre for Disease Control. Point prevalence survey of healthcareassociated infections and antimicrobial use in European acute care hospitals 2011–2012. 2013
[Available from:

https://www.ecdc.europa.eu/sites/default/files/media/en/publications/Publications/healthcare-associated-infections-antimicrobial-use-PPS.pdf].

18. Aronsson B, Möllby R, Nord CE. Antimicrobial agents and *Clostridium difficile*in acute enteric disease: epidemiological data from Sweden, 1980-1982. *J Infect Dis*.
1985;151(3):476-81.

Säll O, Johansson K, Norén T. Low colonization rates of *Clostridium difficile* among patients and healthcare workers at Örebro University Hospital in Sweden. *Apmis*. 2015;123(3):240-4.

20. Dubberke ER, Reske KA, Seiler S, Hink T, Kwon JH, Burnham CA. Risk Factors for Acquisition and Loss of *Clostridium difficile* Colonization in Hospitalized Patients. *Antimicrob Agents Chemother*. 2015;59(8):4533-43.

21. Enkirch T, Mernelius S, Magnusson C, Kühlmann-Berenzon S, Bengnér M,
Åkerlund T, et al. Molecular epidemiology of community- and hospital-associated *Clostridioides difficile* infections in Jönköping, Sweden, October 2017 - March 2018. *Apmis*.
2022;130(11):661-70.

22. National Board of Health and Welfare. DRG-statistik. 2020 [Available from: https://www.socialstyrelsen.se/globalassets/sharepoint-dokument/artikelkatalog/statistik/2020-11-7042.pdf].

23. Organisation of Economic Co-operation and Development. Average length of stay in hospital. 2021 [Available from: https://www.oecd-ilibrary.org/sites/c762d8be-en/index.html?itemId=/content/component/c762d8be-en].

24. Eyre DW, Griffiths D, Vaughan A, Golubchik T, Acharya M, O'Connor L, et al.
Asymptomatic *Clostridium difficile* colonisation and onward transmission. *PLoS One*.
2013;8(11):e78445.

25. Louie TJ, Miller MA, Mullane KM, Weiss K, Lentnek A, Golan Y, et al.
Fidaxomicin versus vancomycin for *Clostridium difficile* infection. *N Engl J Med*.
2011;364(5):422-31.

26. Cornely OA, Crook DW, Esposito R, Poirier A, Somera MS, Weiss K, et al. Fidaxomicin versus vancomycin for infection with *Clostridium difficile* in Europe, Canada, and the USA: a double-blind, non-inferiority, randomised controlled trial. *Lancet Infect Dis*. 2012;12(4):281-9.

Wilcox MH, Howe R. Diarrhoea caused by *Clostridium difficile*: response time for treatment with metronidazole and vancomycin. *J Antimicrob Chemother*. 1995;36(4):673-9.

28. Jung KS, Park JJ, Chon YE, Jung ES, Lee HJ, Jang HW, et al. Risk Factors for Treatment Failure and Recurrence after Metronidazole Treatment for *Clostridium difficile*-associated Diarrhea. *Gut Liver*. 2010;4(3):332-7.

29. Fekety R, Kim KH, Brown D, Batts DH, Cudmore M, Silva J, Jr. Epidemiology of antibiotic-associated colitis; isolation of *Clostridium difficile* from the hospital environment. *Am J Med.* 1981;70(4):906-8.

30. Warren BG, Turner NA, Addison R, Nelson A, Barrett A, Addison B, et al. The Impact of Infection Versus Colonization on *Clostridioides difficile* Environmental

Contamination in Hospitalized Patients With Diarrhea. *Open Forum Infect Dis*. 2022;9(4):ofac069.

31. Sethi AK, Al-Nassir WN, Nerandzic MM, Bobulsky GS, Donskey CJ.
Persistence of skin contamination and environmental shedding of *Clostridium difficile* during and after treatment of C. difficile infection. *Infect Control Hosp Epidemiol*. 2010;31(1):21-7.

32. Slimings C, Riley TV. Antibiotics and healthcare facility-associated *Clostridioides difficile* infection: systematic review and meta-analysis 2020 update. *J Antimicrob Chemother*. 2021;76(7):1676-88.

Public Health Agency of Sweden and the National Veterinary Institute.
 SWEDRES SWARM 2021 Sales of antibiotics and occurrence of antibiotic resistance in
 Sweden. 2022 [Available from:

https://www.sva.se/media/8da965da486b11e/swedres_svarm_2021.pdf].

Feazel LM, Malhotra A, Perencevich EN, Kaboli P, Diekema DJ, Schweizer
 ML. Effect of antibiotic stewardship programmes on *Clostridium difficile* incidence: a
 systematic review and meta-analysis. *J Antimicrob Chemother*. 2014;69(7):1748-54.

35. Anjewierden S, Han Z, Brown AM, Donskey CJ, Deshpande A. Risk factors for *Clostridioides difficile* colonization among hospitalized adults: A meta-analysis and systematic review. *Infect Control Hosp Epidemiol*. 2021;42(5):565-72.

36. Chau JPC, Liu X, Lo SHS, Chien WT, Wan X. Effects of environmental cleaning bundles on reducing healthcare-associated *Clostridioides difficile* infection: a systematic review and meta-analysis. *J Hosp Infect*. 2020;106(4):734-44.

37. Sheth PM, Douchant K, Uyanwune Y, Larocque M, Anantharajah A,
Borgundvaag E, et al. Evidence of transmission of *Clostridium difficile* in asymptomatic patients following admission screening in a tertiary care hospital. *PLoS One*.
2019;14(2):e0207138.

38. Riggs MM, Sethi AK, Zabarsky TF, Eckstein EC, Jump RL, Donskey CJ.
Asymptomatic carriers are a potential source for transmission of epidemic and nonepidemic *Clostridium difficile* strains among long-term care facility residents. *Clin Infect Dis*.
2007;45(8):992-8.

39. Durovic A, Widmer AF, Tschudin-Sutter S. New insights into transmission of *Clostridium difficile* infection-narrative review. *Clin Microbiol Infect*. 2018;24(5):483-92.

Figure legends

Figure 1. Schematic description of the model. Boxes correspond to compartments and black arrows indicate patient flow between compartments. Spore flows are represented by grey arrows. The dotted line represents an increased patient flow from unexposed to colonized (C) dependent on the level of spores in the environment (E). For interpretation of symbols, see Table 1.

Figure 2. The outcome of the three interventions on the fraction of colonized and infected. A: The effect of decreased use of high-risk antibiotics; B: The effect of case isolation; C: The effect of increased general cleaning and disinfection.

Tables

Table 1. Model variables and parameter values. CDI: C. difficile infection.

Model para	Model parameters						
Parameter	Interpretation	Input value	Baseline	Comments and references			
		range	model value				
N	Number of patients in the		500	Aimed to represent a			
	hospital			secondary-care hospital			
				in Sweden			
F	Inflow of patients		κ(N-I)	Balanced against outflow			
				for a constant N			
ευ	Proportion unexposed at		0.949	$1 - (\epsilon_{\rm C} + \epsilon_{\rm I} + \epsilon_{\rm R})$			
	admission						
ε _C	Proportion colonized by	0.019-0.115	0.05	1.9%, ¹⁸ 5.2%, ¹⁹ 11.5%			
	toxigenic C. difficile at			20			
	admission						
ε _I	CDI prevalence at	0.0008-	0.001	Incidence of CDI in			
	admission	0.002		western Sweden in 2016:			
				0.8/1000 days of care 10			
				implies (0.8/к)/1000			
				admissions = 0.004, of			
				which a minority (range			
				20–50%) occur before			
				hospital admission. ²¹			

	κ	Rate of discharge from	0.17-0.26	0.2	Mean length of stay at
		hospital for U, C, R			Swedish hospitals $= 3.9$
					days, ²² 5.7 days. ²³
-	β	Rate unexposed ->	0.016-0.021	0.017	Net colonized during
		colonized			their stay: 8.5%, ²⁴ 6.4%
					20
					Divided by 1/κ to obtain
					rate per day: 2.1%, ²⁴
					1.6% ²⁰
-	θ	Rate colonized ->	0.003-0.006	0.004	Incidence of CDI in
		infected			western Sweden in 2016:
					0.8/1000 days of care, ¹⁰
					of which a majority
					(range 50–90%, 0.0004-
					0.007) occurred among
					hospitalized patients. ²¹
					As all infections occur
					among colonized
					patients, this rate is
					divided by %C (at
					equilibrium = 0.121) for a
					range of 0.003-0.006.
	ρ	Rate infected ->	0.14-0.33	0.16	1-3 days from symptoms
		recovered			to treatment, 2-4 days
1			1	1	1

				from treatment to
				symptom resolution ^{25, 26}
				for vancomycin, 4.6 days
				for Metronidazole. ²⁷
				Metronidazole is still
				widely used in Sweden,
				with a rate of treatment
				failure around 12% ²⁸ ,
				after which vancomycin
				is usually prescribed and
				clinical cure is delayed.
				This makes the lower end
				of the range most
				plausible in our setting.
δ	Clearance rate of spores	0.01-0.03	0.0175	Transmission between
	from the environment			patients via the hospital
				environment has been
				traced several months
				from donor to recipient. ¹
				Persistence of spores in
				the hospital environment
				up to five months. ²⁹
αι	Relative increase of E per	-	0.001	Value determined by
	day caused by an			feedback of model output
				(see Methods).

	individual infected			
	patient			
α _C	Relative increase of E per	0.08-0.65 *	0.00025	0.25 * α _I
	day caused by an	$\alpha_{\rm I}$		Relative frequency of
	individual colonized			excess hospital
	patient			contamination (rates
				above uncolonized
				controls) in colonized
				versus infected patients
				range from 0.08-0.65
				depending on room area.
				30
α _R	Relative increase of E per	0.02-0.2 *	0.0001	$0.1 * \alpha_{I}$
	day caused by an	α_{I}		Low frequency of <i>C</i> .
	individual patient in			<i>difficile</i> in stools during
	recovery			treatment. ³¹
Model state	variables for internal and ex	ternal consiste	rncy	1
Variable	Interpretation	Target	Baseline	Comments and references
		walua	model	

,		1	200001110	
		value	model	
			output value	
Е	Hospital spore reservoir	0.99-1.01	0.998	
	at equilibrium			
U	Proportion of unexposed	-	435 patients	
	patients		(87.1%)	

С	Proportion of colonized	5-13.5%	61 patients	$> \epsilon_{\rm C} (0.05)$ – more than
	patients (prevalence of		(12.1%)	the proportion colonized
	colonization) at			at admission
	equilibrium			$< \epsilon_{C} + \beta/\kappa (0.135) - less$
				than the proportion
				colonized at discharge
Ι	Proportion of infected	0.2-0.5%	2 patients	0.36% (own hospital
	patients (prevalence of		(0.43%)	data)
	CDI) at equilibrium			0.2%, ¹⁶ 0.5% ¹⁷
R	Proportion of recovering	-	2 patients	
	patients		(0.34%)	

Table 2. Antibiotics: consumption and increased C. difficile infection (CDI) risk. DDD:

Defined daily dose. OR: Odds Ratio.

Antibiotic class	DDD/100	Share of	OR	OR-1	Increase of
	patient-days	total			CDI cases
		DDD/100			attributable to
		patient-			the antibiotic
		days			class (DDD
					share
					* (OR - 1)
Cephalosporins	8.7	11.5%	1.8	0.8	9.0%

Beta-lactamase	17.1	22.5%	1.4	0.4	7.4%
resistant					
penicillins					
			• •		
Carbapenems	3.6	4.7%	2,6	1.6	7.4%
Penicillins with	6.1	8.0%	1.8	0.8	6.2%
extended					
spectrum					
Combinations of	10.6	14.0%	1.4	0.4	6.0%
penicillin					
Fluoroquinolones	6.8	9.0%	1.34	0.34	3.0%
Glycopeptides	1.4	1.8%	1.9	0.9	1.7%
Lincosamides	1.9	2.5%	1.56	0.56	1.4%
Macrolides	1.2	1.6%	1.2	0.2	0.3%
Beta-lactamase	7.1	9.4%	1	0	0%
sensitive					
penicillins					
Tetracyclins	4.1	5.4%	1	0	0%
Trimethoprim-	3.6	4.7%	1	0	0%
sulfamethoxazole					
Aminoglycosides	2.8	1.1%	1	0	0%
Other	2.9	3.8%	1	0	0%
Total	75.9	100.0%			44.1%

Table 3. Maximal effects of interventions. Grey background indicates deviation from the baseline value. For definitions of parameters, see Table 1.

Parameter	Baseline	Antibiotic	Increased	Increased	All
	value	stewardship	CDI isolation	cleaning	interventions
				and	combined
				disinfection	
N	500	500	500	500	500
F	к(N-I)	κ(N-I)	к(N-I)	κ(N-I)	к(N-I)
ευ	0.949	0.949	0.949	0.949	0.949
ε _C	0.05	0.05	0.05	0.05	0.05
ει	0.001	0.001	0.001	0.001	0.001
κ	0.2	0.2	0.2	0.2	0.2
β	0.017	0.017	0.017	0.017	0.017
θ	0.004	0.00233	0.004	0.004	0.00233
ρ	0.16	0.16	0.16	0.16	0.16
δ	0.0175	0.0175	0.0175	0.02275	0.02275
αι	0.001	0.001	0.0001	0.001	0.0001
α _C	0.00025	0.00025	0.00025	0.00025	0.00025
α _R	0.0001	0.0001	0.0001	0.0001	0.0001
Output					
Environmental	0.998	0.936	0.781	0.601	0.484
spores					
Unexposed	87.1%	87.6%	88.6%	90.0%	90.9%
Colonized	12.1%	11.8%	10.7%	9.39%	8.63%
Infected	0.43%	0.30%	0.39%	0.36%	0.25%
Recovering	0.34%	0.23%	0.31%	0.29%	0.20%

Output, change relative to baseline								
Environmental	-	-6.2%	-21.8%	-39.7%	-51.5%			
spores								
Unexposed	-	+0.6%	+1.8%	+3.3%	+4.4%			
Colonized	-	-2.5%	-12.2%	-22.5%	-28.9%			
Infected	-	-30.6%	-8.6%	-16.0%	-41.5%			
Recovering	-	-30.6%	-8.6%	-16.0%	-41.5%			



Figure 1





