1Study demands and health status among medical students in two German2universities.

- 3 Amanda Voss¹ (ORCID: 0000-0002-0116-343X), Susanne Dettmer² (ORCID: 0000-0002-
- 4 3916-6492), Mira Tschorn^{3,4} (ORCID: 0000-0002-9128-3987), Jan C. Zoellick^{2,4*} (ORCID:
- 5 0000-0003-1053-8819)
- 6 ¹ FAU Erlangen-Nürnberg, Institut u. Poliklinik für Arbeits-, Sozial- und Umweltmedizin,
- 7 Henkestraße 9-11, 91054 Erlangen, Germany
- 8 ² Charité Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and
- 9 Humboldt-Universität zu Berlin, Institute of Medical Sociology and Rehabilitation Science,
- 10 Charitéplatz 1, 10117 Berlin, Germany
- ³ Social and Preventive Medicine, Department of Sports and Health Sciences, University of
- 12 Potsdam, Potsdam, Germany.
- 13 ⁴ German Center for Mental Health (DZPG), partner site Berlin/Potsdam
- 14 * Corresponding author: Jan C. Zoellick

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

15

Abstract

16 Background and objectives

Medical students regularly report high study demands and low mental health. We thus studied medical students in two different curricula in Germany investigating their study demands, study and life satisfaction, and overall and mental health with emphasis on contrasting beginners and advanced students.

21 Design and methods

We used online surveys with convenience samples through university mailing lists and student union channels. Uni1 participants (*N*=357; 70% female, *M*=24.83 years) were older than Uni2 participants (*N*=126; 79% female; *M*=23.39 years), but the cohorts did not differ further regarding sociodemographic variables. For analyses, we used *t*-tests for comparisons and correlations for associations.

27 Results

28 Students of both universities reported good mental and overall health as well as high

29 satisfaction with their studies and lives, yet sleep difficulties were prevalent. Study demands

30 were low to medium with the highest demands being learning activities (*M*=5.31; *SD*=2.19,

scale 0-10), self-structuring (*M*=4.61; *SD*=2.01), and performance pressure (*M*=4.45;

32 *SD*=2.27). The students in the reformed degree programme reported fewer issues with

integrating theory and practice compared to those in the regular degree programme

34 $(M_{Uni1}=3.38; SD_{Uni1}=2.05; M_{Uni2}=4.17; SD_{Uni2}=2.06; t(430)=-3.53, p<.001).$

35 Discussion and conclusions

36 Our sample was rather resilient regarding mental health and coping with study demands.

37 The two universities showed little to no differences. With relative increased demands

regarding learning activities, self-structuring, and performance pressure we suggest to focus

- 39 on learning strategies for beginners and strategies for stress reduction to address, among
- 40 other things, the reported sleep difficulties.

41

Keywords: student satisfaction; study demands; mental and overall health; medical
students; online survey; stress

44

1. Introduction

Medical students regularly report excessive prevalence of depression [1], emotional
exhaustion [2], burnout [3], and sleep difficulties [4] leading to low life satisfaction [5] as a
result of stress. Impaired mental health is highly prevalent in university students with
elevated levels of stress and depressive and anxiety symptoms [6].

49 Substance use as a malevolent coping mechanism is common in medical students [7] and 50 translates into stress-burdened professional careers [8, 9]. In addition to these known 51 factors, social support and structural conditions appear to be specific factors that can also 52 impact the health of students [10, 11]. Understanding stress trajectories in their studies and 53 creating an environment for better and healthier academic and professional achievements is one of the great challenges for medical faculties [12, 13]. Supporting resilience in students 54 55 via specific intervention programmes has been proposed to mitigate stress trajectories [14-56 17]. Students burdened with poor mental health suffer disproportionately under high study 57 demands. Furthermore, study demands seem to be a crucial source for stress [11]. An 58 established measure for study demands in the German academic context differentiates six 59 areas of demands: mode of science, learning activities, self-structuring, contact and 60 cooperation, performance pressure and failure, and combining theory and practice [18]. 61 These areas cover the broad range of expectations towards students ranging from

organising oneself to performing in examinations or academic thinking and writing.
Students interact with the learning environment and perceive it in different ways based on

64 their individual resources and characteristics [19]. Medical education in Germany is 65 undergoing a dynamic reform process. After the introduction of the model clause in the Medical Licensing Regulations (ÄAppO) in 1999, numerous initiatives for the reform of 66 67 medical studies have been implemented [20]. Currently, there are 13 reformed degree 68 programs in Germany, including University 1 (Uni1), as well as 26 traditional degree 69 programs, including University 2 (Uni2). A university curriculum can be understood as a 70 learning environment that enables students to have learning experiences in order to achieve 71 specific educational goals [21], and curricula can impact student satisfaction [22-24].

72 The purpose of this paper is to analyse medical students' self-reported difficulties to cope 73 with study demands, depression, exhaustion, sleep difficulties, general health status, and life 74 as well as study satisfaction in two German universities, one structured following the 75 reformed curriculum and one with a traditional curriculum. Exploratory studies have shown 76 that these constructs are closely related to form stress trajectories - students with anxiety and stress tend to be overburdened by the workload leading to disengagement from their 77 78 studies, depression, and burnout [25, 26]. The COVID-pandemic presented an additional 79 burden potentially alleviating mental and overall health in students [27]. We suspect that the

80 workload and anxiety is higher for beginners compared to advanced students. Since

- 81 beginners have little experience in university exams, they lack a frame of reference for
- demands and realistic assessments of their own performance [28, 29]. Although previous
- 83 research indicates no evidence for different rates of depressive symptoms between
- 84 beginners compared to advanced students [30], this lack of a frame of reference might lead
- 85 to inadequate expectations and additional stress.

86 **1.1 Research questions**

The following research questions are formulated: How high are the levels of study
demands, depression, exhaustion, sleep difficulties, general health status, and life
satisfaction in the whole sample? How do Uni1 and Uni2 students differ regarding their
levels of study demands, depression, exhaustion, sleep difficulties, general health status,
and life satisfaction? Are these constructs correlated? What are differences between cohorts

- 92 (beginners and advanced students) regarding study demands and mental health outcomes?
- 93

2. Methods

94 2.1 Measures and definitions

We included the constructs depression, general health status, emotional exhaustion, sleep difficulties, life satisfaction, study satisfaction, and difficulties to cope with study demands in our survey. Table 1 defines the constructs and provides their measures. All constructs were transformed into POMP-scores [31] with a range 0 (low depression scores, bad health, few sleep difficulties, low life and study satisfaction, few difficulties in study demands) to 10 (high depression scores, good health, many sleep difficulties, high life and study satisfaction, many difficulties in study demands).

102 ---TABLE 1 HERE---

103 2.2 Procedure

2.2.1 Survey administration. We programmed an online questionnaire in German hosted
on in-house servers of the respective university in accordance with European data protection
laws. Participation was enabled in spring 2022. We invited students via the dean of studies
mailing list and via the student union's social media appearance.

2.2.2 Ethical considerations. The ethics committee voted that no detailed examination of
the pseudonymous study design was necessary (vote 21-393-ANF from 21 November 2021).
Our study was carried out in accordance with the Declaration of Helsinki, including, but not
limited to the anonymity of participants being guaranteed. Participants provided informed
consent.

2.2.3 Statistical analysis. Data were analysed using IBM SPSS Version 28. We
 calculated POMP scores for each variable in this analysis using the formula

115 $(\frac{(x_i - \min)}{(\max - \min)}) * 10$

116 with x_i as individual observed values, *min* as the minimum value and *max* as the maximum value of the scale [31]. With their range between 0 and 10, POMP-scores can be intuitively 117 118 interpreted in contrast to the varying original scores (e.g., 1 to 5 [18] or 0 to 3 [32]). The 119 underlying linear transformation does not change the rank of values and is thus robust 120 concerning parametric analyses, e.g. for *t*-tests, correlations, or Cronbach's a [31]. We used 121 descriptive statistics, t-tests, and correlations; results were considered statistically significant 122 at $p \leq .05$. For differences between beginners and advanced students, we created a dummy 123 variable with values 0 (semesters 1-3) and 1 (semesters 8-10). We used pairwise deletion for 124 missing data.

125 **2.3 Sample characteristics**

126 2.3.1 Inclusion criteria. Medical students at Uni1 and Uni2 were invited to participate.
127 We included participants who provided answers to ≥20% of the questions.

2.3.2 Exclusion criteria. We excluded students of other subjects and other universities
 from participation. We also excluded datasets with answers to less than 20% of questions or
 incomplete datasets in cases of multiple participations.

131 2.3.3 Sample size calculation. We did not perform a sample size calculation as this data132 collection was exploratory.

133 **2.3.4 Respondent characteristics.** At Uni1, the welcome page was visited 656 times.

134 187 visitors were excluded immediately, because they provided no consent for participation

135 or a subject of study other than medicine. We excluded 42 datasets for multiple

136 participations. We additionally excluded 70 datasets with answers to <20% of questions.

137 Thus, 357 medical students (70% female, M = 24.83 years, SD = 4.99 years) across all ten

semesters of study (*Median* = 5th semester) remained for analysis. At Uni1, 4,837 medical

139 students were enrolled in the winter term 2021/22 resulting in a response rate of 7%.

At Uni2, the welcome page was visited 175 times. 33 visitors were excluded immediately, because they provided no consent for participation or a subject of study other than medicine. We excluded 4 datasets for multiple participations. We additionally excluded 12 datasets with answers to less than 20% of questions. Thus, 126 medical students (79% female, M = 24.41years, SD = 3.47 years) across all ten semesters of study (*Median* = 5th semester) remained for analysis. At Uni2, 2,828 medical students were enrolled in the winter term 2021/22

resulting in a response rate of 4%. Table 2 shows sample characteristics with tests fordifferences between universities.

148 ---TABLE 2 HERE---

149

3. Results

150 Our first research question aimed to assess mental and overall health as well as study 151 demands in medical students. Students generally reported being mentally and overall 152 healthy. They also reported being satisfied with their lives and their studies. Despite low mean depression scores (M = 3.16; SD = 2.75; range 0-10), 111 students (26% of valid 153 154 responses) reported depression values above the clinically relevant threshold. Similarly, 155 mean sleeping difficulty was low (M = 3.95; SD = 2.55), but only 161 participants (38% of 156 valid responses) never or rarely experienced difficulties at all: 115 participants (27% of valid 157 responses) reported *either* difficulties to fall asleep or sleep through the night; and the 158 remaining 148 participants (35% of valid responses) reported difficulties of falling asleep and 159 sleep through the night. Thus, reduced sleep quality was reported at least sometimes by the 160 majority of the sample. Students also reported feeling medium exhausted in their studies. 161 Table 3 shows descriptive statistics of the constructs. Reliability coefficients for all constructs 162 were satisfying with the exception of self-structuring (Cronbach's α = .49).

163 ---TABLE 3 HERE---

164 Our second exploratory question aimed at differences between Uni1 and Uni2. Uni2 165 students reported higher exhaustion that Uni1 students just short of statistical significance 166 (p = .055). Students reported low to medium study demands with $3.38 \le M \le 5.39$ on a scale 167 ranging between 0 and 10. The most challenging aspects for both student groups were self-168 structuring, learning activities, and performance pressure, however in different orders. 169 Significant differences were found in combining theory and practice which was more difficult 170 for Uni2 students than for Uni1 students.

171 Our third exploratory question addressed correlations of constructs. All constructs 172 regarding mental and overall health were closely related $(.33 \le |r| \le .67; p < .01)$ indicating 173 that respondents with low overall health also reported worse mental health, more exhaustion, 174 and more sleep difficulties along with lower life and study satisfaction. All constructs 175 regarding study demands were closely related (.24 \leq |r| \leq .61, p < .01), indicating that some 176 students do well generally in the academic field while others struggle with learning, 177 organising, and the scientific mode simultaneously. Communication and cooperation with 178 fellow students showed the lowest correlations with other study demands. Comparing health 179 and study constructs, mode of science was least associated with mental and overall health. 180 Emotional exhaustion showed highest correlations with difficulties to cope with study

demands. Study satisfaction was more closely related to practical application and study
organisation than to contact with peers. Table 4 shows correlations between the constructs.

183 ---TABLE 4 HERE---

Lastly, our fourth research question addressed differences in academic progression. Advanced medical students reported fewer difficulties to organise their learning and they suffered less from performance pressure than beginners – but only for Uni1 students. No differences were observed for Uni2 students. The other study demands did not differ between beginners and advanced students. Differences between beginners and advanced students of medicine are shown in Table 5.

- 190 ---TABLE 5 HERE----
- 191

4. Discussion

We studied medical students in two different universities in Germany regarding theiroverall and mental health, coping with study demands, and their study and life satisfaction.

194 Results revealed students of both universities reporting good overall and mental health as 195 well as high satisfaction with their studies and lives. However, 26% students reported 196 depressive symptoms above the clinically relevant cut-point. This rate was slightly lower than 197 in a study investigating students at five German universities during the COVID-pandemic 198 [33], but substantially higher to the general German population, where 10.3% (17-29-year-199 olds) and 10.9% (30-39-year-olds) reported elevated depressive symptoms measured with 200 the PHQ-2 [34]. Sleep difficulties as a particular aspect of mental health were prevalent in the 201 majority with 62% of respondents reporting sleep disturbances at least sometimes. This 202 corresponds to 52.7% of 25,735 medical students reporting poor sleep quality that were 203 included in a meta-analysis [4]. In comparison, 36% of the general German population 204 reported poor sleep quality [35]. Thus, students reported seemingly conflicting findings of 205 good overall health and simultaneously more sleep difficulties and depressive symptoms 206 than the general population. However, sleep as a specific measure can be seen as an early 207 indicator of mental health difficulties compared to manifest depressive symptoms or declined 208 overall health. We thus suggest to intervene early to interrupt a pathway from current early 209 indicators of stress and struggle towards chronified versions of stress and struggle.

Study demands most difficult to cope with were learning activities (M = 5.31), selfstructuring (M = 4.61), and performance pressure (M = 4.45). These three demands were also the most difficult to cope with in the representative samples of Jänsch and Bosse [18] across degree programmes with somewhat lower POMP-scores ($M_{LA} = 4.60$; $M_{SS} = 3.93$; $M_{PP} = 4.30$). We observed significant differences between the two universities only regarding

Uni1 students reporting lower difficulties to combine theory and practice than Uni2 students. For both universities, more difficulties combining theory and practice were relatively strongly correlated with depression (r = .31), emotional exhaustion (r = .43), and study satisfaction (r = .48).

Differences between beginners and advanced students to cope with study demands were inconclusive. Advanced students had lower difficulties to cope with learning activities and performance pressure at Uni1, but not at Uni2 – all other study demands did not yield differences in study progression. These differences can be explained via experience – advanced students have built up a wealth of experience in exam situations and can better assess their expectations and competencies compared to beginners.

Surprisingly, academic progress did not correspond to better assessment of contact and cooperation between students at either university. First to third semester students in winter term 2021 had primarily experienced online teaching whereas their more experienced counterparts studied the majority of their academic career in-person. Yet, online teaching with fewer opportunities to get in contact with fellow students did not compromise the social embeddedness in beginners compared to advanced students.

With our findings, we did not replicate the results of previous studies presenting medical students as depressed, overworked, exhausted, and with low quality of life [1-3, 5]. However, sleep difficulties reported in the literature [4] were also prevalent in our sample. We thus suggest to develop and offer programmes for students supporting their mental health with particular focus on sleep.

236 Mental and overall health were correlated, as well as the constructs of study demands. 237 This suggests that some students struggle particularly whereas others are able to cope with 238 study demands more easily. This supports the call to identify vulnerable, struggling students 239 to provide adequate, individual support in their studies.

240 4.1 Limitations

Convenience samples were used for both universities; we invited students to participate via e-mail lists and social media activities of the student unions. Thus, particularly students who are keen to provide feedback on their study experience were inclined to answer presenting selection bias. Despite convenience sampling, gender ratios, age, and median semester of studies were representative of the student body and did not differ between the two universities. We are thus confident that our findings are informative for the respective study populations.

8

248	5. Conclusions
249	Our sample was rather resilient regarding mental health and coping with study demands.
250	Curricular influence on our outcomes seemed comparatively minor, which might result from
251	increased online teaching during the pandemic. We suggest to focus on learning strategies
252	for beginners and strategies for stress reduction to address, among other things, the reported
253	sleep difficulties.
254	Acknowledgements
255	The authors would like to thank Luise Chassol from the Uni1 student union for support in
256	finalising the questionnaire and Wiebke Höppner from the Uni1 dean's office for supporting
257	student recruitment. The authors would also like to thank the Uni1 student assistants Anton
258	Horeis and Janek Wilhelm for survey programming and data cleaning.
259	Contributions statement
260	AV, SD, and JZ developed the study design and questionnaire, and collected the data; JZ
261	analysed the data and created the first manuscript draft; all authors interpreted the data and
262	substantively revised the manuscript; all authors read and approved the final manuscript.
263	Declaration of Interest Statement
264	The authors report there are no competing interests to declare.
265	Funding
266	None.

267

References

Puthran R, Zhang MW, Tam WW, Ho RC. Prevalence of depression amongst medical
 students: A meta-analysis. Medical Education. 2016;50(4):456-68.

Voltmer E, Kieschke U, Spahn C. Studienbezogenes psychosoziales Verhalten und
 Erleben von Medizinstudenten im ersten und fünften Studienjahr. Das Gesundheitswesen.
 2008;70(02):98-104.

Frajerman A, Morvan Y, Krebs M-O, Gorwood P, Chaumette B. Burnout in medical
 students before residency: A systematic review and meta-analysis. European Psychiatry.
 2019;55:36-42.

Rao W-W, Li W, Qi H, Hong L, Chen C, Li C-Y, et al. Sleep quality in medical
 students: A comprehensive meta-analysis of observational studies. Sleep and Breathing.
 2020;24(3):1151-65.

Kurth RA, Klier S, Pokorny D, Jurkat HB, Reimer C. Studienbezogene Belastungen,
 Lebensqualität und Beziehungserleben bei Medizinstudenten. Psychotherapeut.
 2007;52(5):355-61.

Kroher M, Beuße M, Isleib S, Becker K, Ehrhardt M-C, Gerdes F, et al. Die
 Studierendenbefragung in Deutschland: 22. Sozialerhebung. Die wirtschaftliche und soziale
 Lage der Studierenden in Deutschland 2021 Berlin: Bundesministerium für Bildung und
 Forschung (BMBF); 2023 [Available from:

https://www.bmbf.de/SharedDocs/Publikationen/de/bmbf/4/31790_22_Sozialerhebung_2021.
 pdf? blob=publicationFile&v=9.

7. Molodynski A, Lewis T, Kadhum M, Farrell SM, Lemtiri Chelieh M, Falcão De Almeida
T, et al. Cultural variations in wellbeing, burnout and substance use amongst medical
students in twelve countries. International Review of Psychiatry. 2021;33(1-2):37-42.

8. Bransi A, Winter L, Glahn A, Kahl KG. Abhängigkeitserkrankungen bei Ärzten. Der
Nervenarzt. 2020;91(1):77-90.

Liu RQ, Davidson J, Van Hooren TA, Van Koughnett JAM, Jones S, Ott MC.
 Impostorism and anxiety contribute to burnout among resident physicians. Medical Teacher.
 2022;44(7):758-64.

Hennig L, Strack M, Boos M, Reich G. Soziale Unterstützung und psychisches
Befinden von Studierenden. Psychotherapeut. 2017;62(5):431–5.

11. Kirsch A-S, Laemmert P, Tittlbach S. Gesundheitliche Anforderungen und
Ressourcen von Studierenden. Eine Studie mit Studierenden der Sportökonomie. Prävention
und Gesundheitsförderung. 2017;12:181-8.

301 12. Garriott PO, Nisle S. Stress, coping, and perceived academic goal progress in first 302 generation college students: The role of institutional supports. Journal of Diversity in Higher
 303 Education. 2018;11(4):436.

304 13. Ferguson E, James D, Madeley L. Factors associated with success in medical school:
 305 systematic review of the literature. BMJ. 2002;324(7343):952-7.

Healy C, Ryan Á, Moran CN, Harkin DW, Doyle F, Hickey A. Medical students, mental
health and the role of resilience – A cross-sectional study. Medical Teacher. 2023;45(1):40-8.

Backmann J, Weiss M, Schippers MC, Hoegl M. Personality factors, student
 resiliency, and the moderating role of achievement values in study progress. Learning and
 Individual Differences. 2019;72:39-48.

311 16. Neufeld A, Mossière A, Malin G. Basic psychological needs, more than mindfulness
312 and resilience, relate to medical student stress: A case for shifting the focus of wellness
313 curricula. Medical Teacher. 2020;42(12):1401-12.

314 17. Waechter R, Stahl G, Rabie S, Colak B, Johnson-Rais D, Landon B, et al. Mitigating
315 medical student stress and anxiety: Should schools mandate participation in wellness
316 intervention programs? Medical Teacher. 2021;43(8):945-55.

317 18. Jänsch VK, Bosse E. Messinstrument für die Wahrnehmung von

Studienanforderungen. Zusammenstellung sozialwissenschaftlicher Items und Skalen (ZIS).2018.

Billett S. Curriculum considerations. The integration of experiences. In: Billett S,
 editor. Integrating practice-based experiences into higher education. New York: Springer;
 2015. p. 161-94.

323 20. Kiessling C, Rotthoff T, Schnabel KP, Stosch C, Begenau J. 20 years of model
324 curricula in German-speaking countries. GMS Journal for Medical Education.
325 2019;36(5):Doc65.

326 21. Biggs JB. From theory to practice: A cognitive systems approach. Higher Education327 Research and Development. 1993;12(1):73-85.

Schauber SK, Hecht M, Nouns ZM, Kuhlmey A, Dettmer S. The role of environmental
and individual characteristics in the development of student achievement: A comparison
between a traditional and a problem-based-learning curriculum. Advances in Health
Sciences Education. 2015;20(4):1033-52.

332 23. Bosch J. Erfolgreich studieren in der Humanmedizin. Ein Vergleich von traditionellen
333 und reformierten Studiengängen an der Charité–Universitätsmedizin Berlin. Berlin: Freie
334 Universität Berlin (Germany); 2020.

24. Dettmer S, Kuhlmey A. Studienzufriedenheit und berufliche Zukunftsplanung von
Medizinstudierenden - ein Vergleich zweier Ausbildungskonzepte. In: Schwartz FW, Angerer
P, editors. Arbeitsbedingungen und Befinden von Ärztinnen und Ärzten Report
Versorgungsforschung. 2. Berlin: Deutscher Ärzte-Verlag; 2010. p. 103-15.

25. Carrard V, Bourquin C, Berney S, Schlegel K, Gaume J, Bart P-A, et al. The
relationship between medical students' empathy, mental health, and burnout: A crosssectional study. Medical Teacher. 2022;44(12):1392-9.

342 26. Agarwal G, Mosquera M, Ring M, Victorson D. Work engagement in medical
343 students: An exploratory analysis of the relationship between engagement, burnout,
344 perceived stress, lifestyle factors, and medical student attitudes. Medical Teacher.
345 2020;42(3):299-305.

Veer IM, Riepenhausen A, Zerban M, Wackerhagen C, Puhlmann L, Engen H, et al.
Psycho-social factors associated with mental resilience in the Corona lockdown.
Translational Psychiatry. 2021;11(1):1-11.

349 28. Bewick B, Koutsopoulou G, Miles J, Slaa E, Barkham M. Changes in undergraduate
350 students' psychological well-being as they progress through university. Studies in Higher
351 Education. 2010;35(6):633-45.

Pluut H, Curşeu PL, Ilies R. Social and study related stressors and resources among
 university entrants: Effects on well-being and academic performance. Learning and Individual
 Differences. 2015;37:262-8.

355 30. Grohne P, Wenig V, Pischke CR, Matos Fialho PM, Negash S, Niephaus Y, et al.
356 Genderspezifische Unterschiede depressiver Symptome von Studienanfänger*innen
357 während der COVID-19-Pandemie. Prävention und Gesundheitsförderung. 2024.

358 31. Cohen P, Cohen J, Aiken LS, West SG. The problem of units and the circumstance 359 for POMP. Multivariate Behavioral Research. 1999;34(3):315-46.

360 32. Löwe B, Kroenke K, Gräfe K. Detecting and monitoring depression with a two-item 361 questionnaire (PHQ-2). Journal of Psychosomatic Research. 2005;58(2):163-71.

362 33. Heumann E, Helmer SM, Busse H, Negash S, Horn J, Pischke CR, et al. Anxiety and
363 depressive symptoms of German university students 20 months after the COVID-19 outbreak
364 - A cross-sectional study. Journal of Affective Disorders. 2023;320:568-75.

365 34. Hajek A, König H-H. Prevalence and correlates of individuals screening positive for 366 depression and anxiety on the PHQ-4 in the German general population: Findings from the 367 nationally representative German Socio-Economic Panel (GSOEP). International Journal of 368 Environmental Research and Public Health. 2020;17(21):7865.

369 35. Hinz A, Glaesmer H, Brähler E, Löffler M, Engel C, Enzenbach C, et al. Sleep quality
370 in the general population: Psychometric properties of the Pittsburgh Sleep Quality Index,
371 derived from a German community sample of 9284 people. Sleep Medicine. 2017;30:57-63.

36. American Psychiatric Association. Diagnostic and statistical manual of mental
 disorders: American Psychiatric Publishing Inc.; 2000.

374 37. Demerouti E, Bakker AB, Vardakou I, Kantas A. The convergent validity of two
375 burnout instruments: A multitrait-multimethod analysis. European Journal of Psychological
376 Assessment. 2003;19(1):12-23.

377 38. Beierlein C, Kovaleva A, László Z, Kemper CJ, Rammstedt B. Kurzskala zur
378 Erfassung der Allgemeinen Lebenszufriedenheit (L-1). Zusammenstellung
370 anzielwissenschoftlicher Items und Skelen (ZIS). 2015

379 sozialwissenschaftlicher Items und Skalen (ZIS). 2015.

380

Table 1.

Definition of constructs.

Construct	Method	Range	Definition
Difficulties regarding study demands in			
mode of science	MWS [18]	1 to 5	Demands regarding the study content related to scientific language and thinking [18]
learning activities	MWS [18]	1 to 5	Personal demands linked to organising learning and studying [18]
self-structuring	MWS [18]	1 to 5	Organisational demands to structure contextual aspects of studying, e.g., assembling a study schedule [18]
contact and cooperation	MWS [18]	1 to 5	Social demands resulting from teamwork activities and forming bonds with other students [18]
performance pressure and failure	MWS [18]	1 to 5	Personal demands to cope with exam results or stress during exam preparation [18]
combining theory and practice	MWS [18]	1 to 5	Demands related to linking study contents to future occupation topics [18]
Depression	PHQ-2 [32]	0 to 3	"a period of at least 2 weeks during which there is either depressed mood or the loss of interest or pleasure in nearly all activities." [36]
Emotional exhaustion	OLBI [37]	1 to 4	"a consequence of intensive physical, affective, and cognitive strain, i. e., as a longterm consequence of prolonged exposure to certain job demands" [37]
Sleep difficulty	Self- devised	1 to 5	"Dyssomnias are primary disorders of initiating or maintaining sleep or of excessive sleepiness and are characterized by a disturbance in the amount, quality, or timing of sleep." [36]
Health status	Single item	1 to 5	Assessment of one's own health ("How is your health status generally?")
Life satisfaction	L-1 [38]	0 to 100	"cognitive process of assessing one's own wellbeing" [38]
Study satisfaction	Single item	1 to 5	Appraisal of students' study experience ("How satisfied are you overall with your study?")

Note. The column range depicts the value range used in this study. After data collection, all
constructs were transformed into POMP-scores [31] with a range 0 (few difficulties in study
demands, low depression scores, low exhaustion, few sleep difficulties, bad health, and low
life and study satisfaction,) to 10 (many difficulties in study demands, high depression
scores, high exhaustion, many sleep difficulties, good health, and high life and study
satisfaction).

Table 2

Sample characteristics

	Uni1	Uni2	t / χ²	р	d
N	357	126			
Age (M, SD)	24.83 (4.99)	23.39 (3.47)	3.53	.000	0.31
Gender (% women)	71%	79%	3.98	.137	-
Semester (M, SD)	5.31 (3.13)	4.98 (2.62)	1.15	.252	-

387 *Note.* M = mean; SD = standard deviation; t = t-value for age and semester; $\chi^2 =$ Chi-square 388 value for gender; p = p-value test statistics; d = Cohen's d

Table 3

Descriptive re Construct	Range	N	М	SD	α	t	р
Mode of science	0-10	318ª 113 ^b	4.06 3.87	2.26 2.33	.78	0.78	.437
Learning activities	0-10	319ª 113 ^b	5.28 5.39	2.18 2.21	.78	-0.44	.660
Self- structuring	0-10	319ª 113 ^b	4.67 4.42	2.06 1.86	.49	1.14	.254
Contact and cooperation	0-10	319ª 113 ^b	4.36 4.02	2.10 2.17	.69	1.47	.143
Performance pressure and failure	0-10	319ª 113 ^b	4.35 4.73	2.26 2.26	.74	-1.51	.132
Combining theory and practice	0-10	319 ^a 113 ^b	3.38 4.17	2.05 2.06	.70	-3.53	.000
Depression	0-10	312ª 113 ^b	3.19 3.10	2.82 2.53	.87	0.30	.761
Emotional exhaustion	0-10	315ª 114 ^ь	5.35 5.75	2.03 1.84	.87	-1.84	.055
Sleep difficulty	0-10	312ª 114	3.91 4.06	2.57 2.49	.69	-0.54	.589
Health status	0-10	317ª 113 ^b	7.33 7.48	2.34 1.96	-	-0.63	.527
Life satisfaction	0-10	319ª 113 ^b	6.58 6.55	2.28 2.11	-	0.11	.910
Study satisfaction	0-10	331 115	6.75 6.50	2.37 2.29	-	0.99	.332

Descriptive results of mental and overall health, study satisfaction, and study demands

389

Note. M = mean; SD = standard deviation; α = internal consistency (Cronbach's alpha); t = t-value for difference between Uni1 and Uni2 students; p = p-value for t-tests; 390 ^a = Uni1 students; ^b = Uni2 students; Effect size for combining theory and practice: d = 0.39. 391 Health status, life satisfaction, and study satisfaction were each measured with a single item, 392 so Cronbach's α could not be calculated. 393

Table 4

		1	2	3	4	5	6	7	8	9	10	11	12
1	Mode of science	-											
2	Learning activities	.37**	-										
3	Self- structuring	.35**	.49**	-									
4	Contact and cooperation	.26**	.35**	.34**	-								
5	Performance pressure and failure	.39**	.61**	.45**	.27**	-							
6	Combining theory and practice	.31**	.38**	.37**	.24**	.35**	-						
7	Depression	12 [*]	.40**	.30**	.30**	.35**	.31**	-					
3	Emotional exhaustion	.29**	.65**	.46**	.28**	.54**	.43**	.55**	-				
9	Sleep difficulty	.11*	.30**	.21**	.20**	.29**	.16**	.40**	.41**	-			
10	Health status	12 [*]	30**	19**	23**	30**	17**	46**	44**	41**	-		
11	Life satisfaction	18**	41**	33**	30**	30**	34*	67**	50**	33**	.52**	-	
12	Study satisfaction	21**	34**	40**	21**	29**	48**	32**	44**	24**	.32**	.51**	-

Correlations between study demands, mental and overall health, and study satisfaction.
--

Table 5

Construct	University	Ν	Μ	SD	t	р	d
Mode of science	Uni1	108 ^a 94 ^b	3.94 4.17	2.03 2.40	-0.75	.457	-
	Uni2	34 ^a 22 ^b	3.76 4.81	2.46 2.02	-1.67	.102	-
Learning activities	Uni1	108 ^a 94 ^b	5.63 4.83	2.02 2.24	2.66	.008	0.38
	Uni2	34 ^a 22 ^b	5.48 5.42	2.19 2.12	0.09	.930	-
Self-structuring	Uni1	108 ^a 94 ^b	4.63 4.70	2.14 2.10	-0.23	.818	-
	Uni2	34 ^a 22 ^b	4.20 4.66	1.80 1.79	-0.93	.358	-
Contact and cooperation	Uni1	108 ^a 94 ^b	4.12 4.40	2.03 2.35	-0.89	.377	-
	Uni2	34 ^a 22 ^b	4.48 4.13	2.31 2.39	0.56	.581	-
Performance pressure and failure	Uni1	108 ^a 94 ^b	4.74 3.82	2.26 2.35	2.85	.005	0.40
	Uni2	34 ^a 22 ^b	4.64 4.70	2.56 2.14	-0.08	.468	-
Combining theory and practice	Uni1	108ª 94 ^b	3.16 3.48	1.95 2.22	-1.09	.282	-
	Uni2	34 ^a 22 ^b	3.85 4.72	2.11 2.37	-1.43	.079	-

D:# 4 dadu . . , . ,

Note. N_{beginners} = 142; N_{advanced} = 116; M = mean; SD = standard deviation; t = t-value for difference between beginners and advanced of medical studies; p = p-value for t-tests; d = Cohen's d; ^a = beginners; ^b = advanced. 395 396 397