

**Effects of nationwide alerting for acute kidney injury on healthcare and patient outcomes:
population based, regression discontinuity analysis**

Supplement

This supplement contains:

- **Additional Methods**
- **Figure S1 to S4**
- **Tables S1 to S8**

Additional methods


Running variable derivation:

For this analysis, we defined a centered running variable (RV) based on the AKI algorithm as implemented in Wales: for an index serum creatinine measurement on a given date (C), the baseline value (B) was determined as the minimum serum creatinine level recorded in the preceding seven days. If no tests were conducted in this timeframe, the baseline value was calculated as the median value from the preceding 8–365 days. The RV was then calculated as $RV = C - (1.5 \times B)$. Thus, the RV represented the change in creatinine level relative to the 50% rise threshold. Following the AKI alert algorithm, an alert would be triggered when $RV \geq 0$ on the date of the index test.

Local linear regression:

To quantify the discontinuity, we modelled the outcomes as a linear function of the centered RV, separately for observations on either side of the threshold. The discontinuity was defined as the difference between the two regression lines at the threshold. To minimize bias from misspecifying the functional form, we restricted the analysis to observations within a narrow bandwidth around the threshold. While a narrow bandwidth reduces bias, it increases the variance of the estimate. Following the standard practice, we selected a bandwidth optimized to balance this bias-variance trade-off. Asymmetry in the running variable's distribution was accounted for by allowing different bandwidths on either side of the threshold. We further applied a triangular kernel, assigning linearly greater weight to observations closer to the threshold.

Panel A: Hospital Setting



Panel B: Community Setting




Fig S1. Illustration of index test selection and timing of follow-up^{1,2}

¹ The triangles indicate the serum creatinine tests taken.

² The section in grey indicates the duration of an emergency inpatient stay.

STOP AKI	Response	
S epsis	Deliver Sepsis Six (if appropriate)	<input type="checkbox"/>
T oxins	Review Medications	<input type="checkbox"/>
	Stop / Avoid potential nephrotoxins *	<input type="checkbox"/>
O ptimise BP	Assess volume status: give fluids (if appropriate)	<input type="checkbox"/>
	Start a fluid balance chart	<input type="checkbox"/>
	Review BP lowering medication *	<input type="checkbox"/>
P revent harm	Treat complications	
	Monitor renal function to assess response	
	Record cause of AKI:	
* Review daily - consider restarting within clinical context (including at discharge from hospital)		
Urine dipstick result:		<input type="checkbox"/>
USS KUB urgent if:	<ul style="list-style-type: none"> • No identified cause of AKI • Possibility of renal tract obstruction • Deteriorating renal function 	
Date:	Time:	
Name:	Grade:	Sign:

Fig S2. Care bundle that was completed and recorded for interruptive alerts.

AKI Telephoning Guidelines




Fig S3. Algorithm used for the interruptive telephoning of AKI alerts.




Fig S4. Distribution of the centred running variable for the hospital and community setting.

	Hospital setting				Community setting			
	N	Mean			N	Mean		
		Full sample	Sample near threshold			Full sample	Sample near threshold	
			below	above			below	above
Baseline variables								
Age (year)	861,494	64.64	66.94	68.28	354,505	64.25	64.55	66.06
Female (%)	861,491	54.47	60.68	57.00	354,505	54.41	63.60	60.26
Rural area (%)	861,494	32.61	33.64	33.46	354,505	33.30	34.61	34.22
WIMD decile (1 = most deprived)	816,252	5.05	4.97	4.98	335,528	5.13	5.07	5.05
Baseline eGFR (ml/min/1.73m ²)	861,494	78.20	85.78	81.66	354,505	81.15	92.78	89.51
Had diabetes at baseline (%)	861,494	23.60	27.63	30.50	354,505	24.70	24.78	27.90
Had CKD3+ at baseline (%)	861,494	29.26	23.28	27.60	354,505	24.69	17.55	21.07
Had heart failure at baseline (%)	861,494	16.67	19.84	21.66	354,505	16.26	18.28	20.27
Had cerebrovascular disease in the past 180 days (%)	861,494	2.82	2.39	2.60	354,505	3.88	2.75	3.36
Had peripheral artery disease in the past 180 days (%)	861,494	3.15	3.89	3.81	354,505	2.86	3.75	4.16
Had coronary heart disease in the past 180 days (%)	861,494	19.06	19.34	19.98	354,505	19.84	17.55	17.86
Had stroke in the past 180 days (%)	861,494	4.12	3.84	4.00	354,505	3.88	2.82	3.61
RAASI use in the past 180 days (%)	861,494	28.04	30.47	33.07	354,505	31.23	28.43	30.21
Diuretic use in the past 180 days (%)	861,494	17.01	20.93	23.04	354,505	18.47	23.29	23.83
NSAID use in the past 180 days (%)	861,494	8.99	8.35	8.01	354,505	10.07	9.06	8.88
PPI use in the past 180 days (%)	861,494	38.60	40.80	40.78	354,505	42.84	43.76	43.75
Statin use in the past 180 days (%)	861,494	31.45	32.68	34.69	354,505	34.31	29.56	32.36
Interruptive alert setting (%)	861,494	14.06	15.55	15.54				
With history of vascular disease (%)	861,494	31.27	33.80	35.73	354,505	32.48	31.25	33.82
With major surgery during admission (%)	861,494	0.78	0.92	0.94				
Exposure variable								
AKI alert (%)	861,494	5.75	16.52	92.70	354,505	2.01	5.84	91.22
Outcome variables								
Death in 365 days (%)	861,494	21.14	29.81	32.54	354,505	15.79	25.32	28.65
Emergency hospitalization or emergency department attendance in 180 days (%)	808,890	49.26	52.98	53.56	354,505	51.06	61.18	66.13
Death in 30 days (%)	861,494	6.33	11.66	13.83	354,505	2.60	6.64	8.63
Emergency hospitalization or emergency department attendance in 30 days (%)	808,890	23.48	26.00	26.13	354,505	26.68	40.15	45.86
Having an inpatient creatinine retest (%)	861,494	54.88	70.95	77.31				
Peak inpatient creatinine after test (micromol/L)	472,763	104.76	106.80	116.23				
Updated eGFR at 90 days (ml/min/1.73m ²)	589,545	79.08	80.04	77.10	212,213	78.77	81.51	78.21
Length of inpatient stay >=7 days (%)	843,778	37.00	47.21	50.28				
Any PPI prescription in 365 days (%)	675,317	43.16	44.95	45.37	298,524	45.63	47.80	46.77
Any NSAID prescription in 365 days (%)	675,317	11.08	9.91	9.34	298,524	10.97	8.97	10.06
RAASI continuation in 180 days among prior users	198,243	84.82	80.93	80.45	99,965	88.36	85.90	83.55
Diuretic continuation in 180 days among prior users	105,081	82.79	79.72	82.37	54,709	84.39	82.99	82.84
Statin continuation in 180 days among prior users	220,755	91.82	90.03	92.07	108,962	93.07	92.19	89.21
Having cancer in 365 days (%)	808,890	3.64	3.55	3.70	354,505	3.02	2.69	3.31
Proteinuria check in 365 days (%)	675,317	20.00	21.84	24.34	298,524	21.94	21.50	24.61
Creatinine test in 90 days (%)	752,412	52.78	57.98	59.08	332,115	55.19	66.29	68.19
Blood pressure measurement in 90 days (%)	752,412	34.02	35.38	37.65	332,115	36.43	36.56	36.32
AKI coding in primary care in 90 days (%)	760,553	1.54	3.30	4.67	332,115	0.59	1.97	5.45
AKI coding for hospitalization (%)	808,890	6.18	16.08	26.48				

Table S1. sample characteristics¹

¹“Sample near threshold” is defined as observations with a change in serum creatinine relative to its 50% increase threshold within [-5, 0) and [0, 5) micromol/L for “below” and “above” the threshold, respectively.

²N is the sample size. There are 861,494 observations included in the hospital setting, among with, 52,604 observations are excluded for emergency hospitalization or emergency department attendance and AKI coding for hospitalization outcomes because they did not survive until discharge; 388,731 are excluded for peak inpatient creatinine outcome because they did not receive another creatinine test during the hospitalization; 271,949 are excluded for having updated eGFR outcome because they did not test for eGFR or survive within 90 days; 17,716 are excluded for length of stay outcome because they did not survive for at least 7 days; 186,177 are excluded for PPI and NSAID prescription, and proteinuria check outcomes because they did not survive for at least 365 days; 663,251, 756,413 and 640,739 are excluded for RAASI, diuretic and statin continuation outcomes, respectively, because they were not prescribed with the respective medication within 180 days before test or did not survive for at least 180 days after discharge; 109,082

are excluded for creatinine test and blood pressure measurement because they did not survive for at least 90 days after discharge; 100,941 are excluded for AKI coding in primary care because they did not survive until discharge or for at least 90 days after test.

³There are 354,505 observations included in the community setting, among with, 142,292 are excluded for having updated eGFR outcome because they did not test for eGFR or survive within 90 days; 55,981 are excluded for PPI and NSAID prescription, and proteinuria check outcomes because they did not survive for at least 365 days; 254,540, 299,796 and 245,543 are excluded for RAASI, diuretic and statin continuation outcomes, respectively, because they were not prescribed with the respective medication within 180 days before test or did not survive for at least 180 days after test; 22,390 are excluded for creatinine test, blood pressure measurement and AKI coding in primary care because they did not survive for at least 90 days after test.

	Hospital setting						Community setting					
	Mean below threshold	Difference at the cutoff		Bandwidth (micromol/L)		Effective sample size	Mean below threshold	Difference at the cutoff		Bandwidth (micromol/L)		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Exposure variable												
AKI alert (%)	16.52	71.52 (68.63, 73.70)	<0.001***	5	38	30,796	5.84	82.88 (79.78, 85.00)	<0.001***	7	38	8,872
Baseline variable												
Had diabetes at baseline (%)	27.63	1.50 (-0.40, 2.83)	0.140	14	68	66,391	24.78	1.50 (-2.18, 4.06)	0.554	12	55	17,572
Had CKD3+ at baseline (%)	23.28	2.02 (-0.48, 4.00)	0.123	8	36	37,488	17.55	-0.98 (-5.90, 2.38)	0.405	6	36	8,792
Had heart failure at baseline (%)	19.84	0.60 (-1.32, 1.86)	0.743	11	55	55,230	18.28	-0.41 (-3.96, 1.92)	0.497	11	46	15,485
Had cerebrovascular disease in the past 180 days (%)	2.39	0.10 (-0.38, 0.51)	0.779	18	167	104,929	2.75	0.57 (-0.42, 1.42)	0.287	23	87	54,476
Had peripheral artery disease in the past 180 days (%)	3.89	-0.20 (-0.94, 0.33)	0.351	17	78	85,473	3.75	0.30 (-1.08, 1.31)	0.852	16	72	24,875
Had coronary heart disease in the past 180 days (%)	19.34	0.68 (-0.98, 2.06)	0.486	12	100	64,211	17.55	-0.90 (-4.35, 1.48)	0.333	11	46	15,518
Had stroke in the past 180 days (%)	3.84	0.29 (-0.34, 0.92)	0.369	16	111	81,136	2.82	0.80 (-0.25, 1.69)	0.145	22	81	54,442
Age (year)	66.94	0.20 (-0.78, 0.84)	0.937	11	37	51,775	64.55	0.19 (-1.52, 1.29)	0.873	11	44	15,429
Female (%)	60.68	-2.34 (-4.54, 0.55)	0.125	7	62	42,460	63.60	-3.35 (-7.26, 1.54)	0.203	8	67	10,574
RAASI use in the past 180 days (%)	30.47	1.28 (-1.01, 2.95)	0.337	10	65	52,650	28.43	-0.16 (-4.32, 2.86)	0.690	10	53	14,008
Diuretic use in the past 180 days (%)	20.93	0.38 (-1.54, 1.62)	0.960	12	54	59,464	23.29	-2.50 (-6.39, 0.01)	0.051	11	44	15,457
NSAID use in the past 180 days (%)	8.35	-0.62 (-1.59, 0.28)	0.168	14	137	76,389	9.06	0.21 (-1.47, 1.98)	0.773	17	90	27,947
PPI use in the past 180 days (%)	40.80	0.01 (-1.76, 1.32)	0.780	17	96	94,002	43.76	0.68 (-2.43, 3.29)	0.770	18	97	31,293
Statin use in the past 180 days (%)	32.68	1.71 (-0.38, 3.44)	0.117	11	81	58,412	29.56	1.04 (-2.95, 4.00)	0.767	11	60	15,831
Baseline eGFR (ml/min/1.73m ²)	85.78	-1.26 (-2.40, 0.55)	0.217	7	28	35,004	92.78	1.08 (-1.11, 4.48)	0.237	6	31	8,627
Rural area (%)	33.64	-0.09 (-1.63, 1.77)	0.936	12	156	66,575	34.61	-1.19 (-4.11, 2.22)	0.560	13	86	20,056
WIMD decile (1 = most deprived)	4.97	0.03 (-0.08, 0.11)	0.721	15	107	76,416	5.07	0.03 (-0.12, 0.20)	0.616	22	80	51,291

Table S2. Balance tests for baseline variables^{1,2,3}

¹Mean below threshold is calculated as the mean within the 5 micromol/L below the threshold.

²Effective sample size is the number of observations within the MSE-optimal bandwidth.

³Abbreviations: AKI = acute kidney injury; CKD3+ = chronic kidney disease of stage 3 or higher; WIMD decile = Welsh Index of Multiple Deprivation decile (1 = most deprived).

	Hospital setting						Community setting					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Negative outcome control												
Statin continuation in 180 days among prior users (%)	90.03	1.37 (-1.52, 4.53)	0.330	11	77	14,390	92.19	-3.44 (-7.70, 0.77)	0.108	14	69	5,437
Having cancer in 365 days (%)	3.55	-0.09 (-1.24, 0.86)	0.720	11	98	47,454	2.69	0.51 (-0.84, 1.76)	0.489	13	101	20,184

Table S3. Negative outcome controls^{1,2,3}

¹Mean below threshold is calculated as the mean within the 5 micromol/L below the threshold.

²Effective sample size is the number of observations within the MSE-optimal bandwidth.

³Abbreviations: AKI = acute kidney injury

	Hospital setting						Community setting					
	Mean below threshold	Effect of AKI alert		Bandwidth		Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Primary outcome												
Death in 365 days (%)	31.60	1.57 (-0.93, 3.16)	0.284	17	44	49,077	24.51	1.62 (-3.20, 4.96)	0.672	13	58	10,440
Emergency hospitalization or emergency department attendance in 180 days (%)	50.43	1.78 (-0.40, 4.25)	0.105	13	149	37,354	54.94	2.35 (-2.49, 5.53)	0.458	17	62	16,057
Secondary outcome												
Health outcome												
Death in 30 days (%)	12.95	1.83 (-0.12, 3.16)	0.069	14	45	41,874	5.87	1.76 (-0.93, 3.67)	0.243	14	67	11,730
Emergency hospitalization or emergency department attendance in 30 days (%)	24.50	-0.58 (-2.14, 0.72)	0.33	30	133	138,110	33.19	3.97 (-0.95, 7.40)	0.131	15	68	11,740
Kidney recovery outcome												
Peak inpatient creatinine after test (micromol/L)	111.40	3.21 (-1.91, 7.19)	0.255	12	62	26,424						
Updated eGFR at 90 days (ml/min/1.73m ²)	77.64	-1.24 (-2.97, 0.86)	0.282	10	45	20,663	77.08	3.15 (-0.52, 8.67)	0.082	7	43	3,417
Care process outcome												
Length of inpatient stay >=7 days (%)	51.98	1.45 (-1.06, 3.56)	0.290	15	47	42,929						
Having an inpatient creatinine retest (%)	69.46	1.29 (-0.96, 3.11)	0.301	16	39	44,595						
Proteinuria check in 365 days (%)	25.58	0.32 (-2.66, 2.44)	0.933	12	74	24,784	24.68	0.64 (-5.13, 4.97)	0.975	11	75	5,542
Creatinine test in 90 days (%)	50.89	0.41 (-2.52, 2.40)	0.962	14	73	33,627	57.45	1.78 (-3.50, 5.48)	0.666	15	69	11,434
Blood pressure measurement in 90 days (%)	36.33	-0.43 (-3.21, 1.60)	0.513	14	92	34,331	37.23	0.98 (-4.26, 4.80)	0.907	15	70	10,231
AKI coding in primary care in 90 days (%)	0.37	0.03 (-0.30, 0.28)	0.937	23	69	74,663	0.34	0.21 (-0.25, 0.75)	0.330	32	56	63,817
AKI coding for hospitalization (%)	4.72	1.30 (-0.31, 2.41)	0.130	10	53	26,937						
Any PPI prescription in 365 days (%)	43.43	2.46 (-0.01, 4.62)	0.051	20	102	45,801	45.00	4.25 (-0.48, 8.15)	0.082	20	107	18,011
Any NSAID prescription in 365 days (%)	13.24	-0.24 (-1.79, 1.63)	0.927	17	88	34,906	12.86	0.73 (-2.14, 4.08)	0.541	17	114	12,630
RAASI continuation in 180 days among prior users (%)	83.37	-0.15 (-2.44, 3.21)	0.790	20	85	15,567	84.69	3.78 (-0.94, 9.59)	0.108	19	84	4,952
Diuretic continuation in 180 days among prior users (%)	84.65	1.87 (-1.13, 4.60)	0.236	34	101	23,177	87.70	1.49 (-4.38, 6.18)	0.738	25	75	5,519

Table S4. Falsification test (Negative exposure control)^{1,2,3}

¹Mean below threshold is calculated as the mean within the 5 micromol/L below the threshold.

²Effective sample size is the number of observations within the MSE-optimal bandwidth.

³Abbreviations: AKI = acute kidney injury

A

	Interruptive alert						Passive alert					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size		Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	
		Estimate (95% CI)	P value					Estimate (95% CI)	P value			
Primary outcome												
Death in 365 days (%)	26.73	1.30 (-5.56, 6.26)	0.907	11	62	8,641	30.38	1.19 (-3.42, 4.87)	0.733	6	50	30,057
Emergency hospitalization or emergency department attendance in 180 days (%)	53.27	3.45 (-2.54, 9.65)	0.253	13	112	8,530	52.92	-0.82 (-4.89, 3.36)	0.716	6	100	30,658
Secondary outcome												
Health outcome												
Death in 30 days (%)	11.45	0.24 (-4.14, 2.60)	0.654	19	59	15,149	11.70	0.46 (-2.48, 2.72)	0.926	7	48	34,237
Emergency hospitalization or emergency department attendance in 30 days (%)	26.36	1.60 (-3.47, 6.54)	0.548	14	121	9,273	25.93	-1.72 (-4.87, 1.30)	0.258	8	114	35,498
Kidney recovery outcome												
Peak inpatient creatinine after test (micromol/L)	106.22	-1.98 (-15.36, 5.99)	0.389	13	83	7,127	106.90	8.21 (1.05, 13.16)	0.021*	7	86	29,056
Updated eGFR at 90 days (ml/min/1.73m ²)	82.76	-3.92 (-7.68, 1.32)	0.166	10	61	5,239	79.55	-0.88 (-3.30, 2.20)	0.694	6	43	20,401
Care process outcome												
Length of inpatient stay >=7 days (%)	42.47	2.88 (-4.73, 8.19)	0.6	12	65	8,190	48.08	-3.71 (-8.93, 0.33)	0.069	5	51	28,157
Having an inpatient creatinine retest (%)	68.27	8.12 (0.70, 13.39)	0.030*	9	71	7,609	71.45	-0.65 (-5.31, 2.76)	0.535	5	46	29,598
Proteinuria check in 365 days (%)	22.56	2.95 (-2.51, 7.78)	0.315	15	134	8,200	21.70	4.73 (0.93, 7.82)	0.013*	8	81	23,891
Creatinine test in 90 days (%)	54.73	4.07 (-3.26, 9.53)	0.336	13	88	7,571	58.59	1.64 (-3.47, 5.76)	0.627	6	79	24,472
Blood pressure measurement in 90 days (%)	39.44	-4.12 (-10.47, 1.16)	0.117	14	115	9,145	34.63	5.55 (2.10, 8.91)	0.002**	9	129	31,731
AKI coding in primary care in 90 days (%)	5.18	3.29 (0.17, 4.14)	0.033*	36	88	48,305	2.95	0.69 (-0.70, 1.37)	0.526	17	47	55,783
AKI coding for hospitalization (%)	15.55	12.66 (6.23, 14.94)	<0.001***	21	44	17,529	16.17	5.39 (1.50, 7.13)	0.003**	10	30	33,597
Any PPI prescription in 365 days (%)	51.69	1.74 (-4.29, 6.97)	0.641	17	128	9,705	43.64	1.53 (-2.42, 5.30)	0.465	8	100	26,287
Any NSAID prescription in 365 days (%)	12.50	-1.46 (-5.28, 2.92)	0.571	13	147	6,949	9.41	-0.08 (-2.19, 2.38)	0.936	8	94	26,142
RAASI continuation in 180 days among prior users (%)	82.47	1.25 (-5.47, 9.56)	0.594	16	83	3,424	80.59	2.63 (-2.19, 8.23)	0.255	10	70	10,510
Diuretic continuation in 180 days among prior users (%)	83.41	4.88 (-3.86, 13.82)	0.270	23	81	2,913	78.98	5.39 (-0.25, 11.03)	0.061	14	85	7,830

B

	Most deprived quintile						Least deprived quintile					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size		Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	
		Estimate (95% CI)	P value					Estimate (95% CI)	P value			
Primary outcome												
Death in 365 days (%)	25.64	0.45 (-6.00, 5.57)	0.942	8	55	10,113	33.08	0.16 (-7.75, 6.77)	0.895	10	92	7,483
Emergency hospitalization or emergency department attendance in 180 days (%)	55.56	-1.17 (-6.44, 4.49)	0.727	11	106	11,060	48.70	5.28 (-2.23, 12.74)	0.169	11	113	7,409
Secondary outcome												
Health outcome												
Death in 30 days (%)	9.14	2.55 (-1.17, 5.08)	0.22	13	54	13,671	12.74	-0.74 (-5.95, 2.59)	0.441	16	66	11,106
Emergency hospitalization or emergency department attendance in 30 days (%)	25.60	2.50 (-1.91, 7.09)	0.259	12	137	12,176	24.53	-1.29 (-7.02, 4.45)	0.66	13	119	8,612
Kidney recovery outcome												
Peak inpatient creatinine after test (micromol/L)	103.72	3.71 (-6.96, 11.16)	0.65	11	96	9,579	109.95	3.29 (-9.55, 12.78)	0.777	12	93	6,675
Updated eGFR at 90 days (ml/min/1.73m ²)	83.48	-1.40 (-4.93, 3.19)	0.674	8	56	7,197	75.94	-3.06 (-7.03, 2.30)	0.321	9	81	5,086
Care process outcome												
Length of inpatient stay >=7 days (%)	44.06	2.07 (-5.09, 7.62)	0.697	9	62	9,651	52.30	-4.46 (-13.48, 2.24)	0.161	10	71	7,189
Having an inpatient creatinine retest (%)	68.41	0.60 (-6.21, 5.54)	0.912	8	56	9,458	72.57	7.38 (-0.85, 13.67)	0.083	9	62	6,604
Proteinuria check in 365 days (%)	21.89	3.13 (-2.19, 7.29)	0.292	12	84	10,063	22.05	3.39 (-3.78, 9.35)	0.405	13	99	5,991
Creatinine test in 90 days (%)	56.34	-2.21 (-9.19, 3.12)	0.334	10	77	8,943	61.55	0.30 (-8.55, 7.39)	0.887	11	83	5,882
Blood pressure measurement in 90 days (%)	34.45	0.90 (-4.35, 5.84)	0.775	12	126	11,878	34.49	4.72 (-2.31, 11.74)	0.188	12	108	7,035
AKI coding in primary care in 90 days (%)	3.75	1.69 (-0.79, 2.35)	0.329	32	48	68,213	3.43	0.70 (-2.24, 1.70)	0.789	32	60	41,180
AKI coding for hospitalization (%)	15.38	8.59 (2.92, 10.78)	<0.001***	17	39	16,585	19.34	5.98 (-1.60, 8.39)	0.182	20	47	14,732
Any PPI prescription in 365 days (%)	49.45	-3.64 (-9.27, 1.35)	0.144	13	126	11,245	39.55	6.58 (-0.93, 14.08)	0.086	13	113	6,046
Any NSAID prescription in 365 days (%)	13.07	-3.38 (-7.02, 0.73)	0.112	11	152	8,919	8.30	-0.69 (-4.81, 3.67)	0.791	12	136	5,671
RAASI continuation in 180 days among prior users (%)	79.96	5.89 (-1.61, 14.32)	0.118	15	103	4,089	79.42	8.56 (0.22, 19.28)	0.045*	16	72	2,554
Diuretic continuation in 180 days among prior users (%)	77.96	11.34 (2.89, 20.68)	0.009**	19	84	3,120	76.10	10.26 (-0.67, 23.09)	0.064	20	78	1,890

C

	Rural						Urban					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size		Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	
		Estimate (95% CI)	P value					Estimate (95% CI)	P value			
Primary outcome												
Death in 365 days (%)	31.87	-0.34 (-6.37, 4.28)	0.701	8	64	14,387	28.77	1.39 (-3.58, 5.41)	0.69	6	49	23,459
Emergency hospitalization or emergency department attendance in 30 days (%)	51.67	0.79 (-4.46, 5.88)	0.788	9	123	15,056	53.63	0.04 (-4.28, 4.57)	0.948	7	95	23,907
Secondary outcome												
Health outcome												
Death in 30 days (%)	13.56	-2.27 (-5.96, -0.05)	0.046*	13	59	21,944	10.69	2.25 (-0.78, 4.57)	0.164	7	48	26,708
Emergency hospitalization or emergency department attendance in 30 days (%)	27.27	-2.98 (-7.02, 1.04)	0.145	12	115	17,324	25.36	1.28 (-2.16, 4.45)	0.496	9	122	28,034
Kidney recovery outcome												
Peak inpatient creatinine after test (micromol/L)	105.89	3.69 (-6.17, 9.78)	0.658	10	76	13,312	107.27	6.63 (-1.22, 12.39)	0.108	7	93	24,092
Updated eGFR at 90 days (ml/min/1.73m ²)	79.58	-2.31 (-5.42, 1.95)	0.357	8	51	9,593	80.27	-0.19 (-2.90, 3.16)	0.934	6	48	16,550
Care process outcome												
Length of inpatient stay >=7 days (%)	47.84	3.07 (-3.32, 8.06)	0.415	8	68	13,445	46.89	-4.71 (-10.40, -0.36)	0.036*	6	53	22,254
Having an inpatient creatinine retest (%)	72.32	7.03 (1.19, 11.41)	0.016*	7	69	14,592	70.26	-1.41 (-6.50, 2.32)	0.353	5	47	23,197
Proteinuria check in 365 days (%)	21.17	1.73 (-3.20, 5.65)	0.588	11	95	12,941	22.17	5.55 (1.65, 8.78)	0.004**	8	96	20,806
Creatinine test in 90 days (%)	58.08	4.05 (-2.03, 9.12)	0.213	9	127	12,412	57.94	0.85 (-4.52, 5.07)	0.912	7	74	20,452
Blood pressure measurement in 90 days (%)	34.99	3.26 (-1.66, 7.81)	0.203	11	177	14,592	35.58	3.62 (-0.05, 7.09)	0.054	9	174	27,447
AKI coding in primary care in 90 days (%)	3.07	1.78 (-0.20, 2.47)	0.097	27	67	60,527	3.42	1.02 (-0.59, 1.62)	0.358	21	47	70,816
AKI coding for hospitalization (%)	14.65	10.09 (5.06, 12.00)	<0.001***	16	38	23,717	16.79	4.63 (0.40, 6.52)	0.027*	11	32	26,761
Any PPI prescription in 365 days (%)	43.19	-0.68 (-6.29, 4.56)	0.755	10	107	12,054	45.81	2.63 (-1.61, 6.34)	0.243	9	145	23,236
Any NSAID prescription in 365 days (%)	8.99	0.15 (-2.75, 3.61)	0.791	10	86	11,818	10.36	0.16 (-2.13, 2.83)	0.784	9	114	21,180
RAASI continuation in 180 days among prior users (%)	82.38	-2.69 (-8.87, 4.78)	0.557	14	73	4,778	80.30	2.94 (-1.94, 8.68)	0.214	11	75	9,523
Diuretic continuation in 180 days among prior users (%)	80.93	2.09 (-4.12, 8.68)	0.485	24	93	5,969	79.10	6.43 (0.29, 12.71)	0.040*	14	89	6,335

D

	Male					Female						
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size		
		Estimate (95% CI)	P value				Estimate (95% CI)	P value				
Primary outcome												
Death in 365 days (%)	33.39	-0.51 (-6.17, 4.02)	0.678	8	66	18,981	27.50	1.63 (-3.68, 5.86)	0.655	5	48	19,564
Emergency hospitalization or emergency department attendance in 180 days (%)	54.71	-0.99 (-5.92, 4.18)	0.736	9	109	18,318	51.88	0.83 (-4.00, 5.82)	0.718	6	99	19,899
Secondary outcome												
Health outcome												
Death in 30 days (%)	13.31	-0.26 (-3.86, 2.35)	0.634	11	56	21,962	10.59	1.46 (-1.84, 4.09)	0.457	6	53	21,382
Emergency hospitalization or emergency department attendance in 30 days (%)	28.43	-2.21 (-5.97, 1.66)	0.268	12	112	21,948	24.46	0.89 (-2.83, 4.42)	0.667	8	112	23,381
Kidney recovery outcome												
Peak inpatient creatinine after test (micromol/L)	120.76	9.59 (0.10, 16.19)	0.047*	9	87	18,235	97.05	0.34 (-7.67, 6.20)	0.836	7	87	18,339
Updated eGFR at 90 days (ml/min/1.73m ²)	81.84	-3.30 (-6.14, 0.40)	0.085	8	47	12,502	78.85	1.02 (-1.97, 4.78)	0.415	5	47	13,568
Care process outcome												
Length of inpatient stay >=7 days (%)	46.58	0.13 (-5.98, 5.07)	0.871	8	62	17,388	47.61	-3.34 (-9.61, 1.26)	0.132	5	52	18,434
Having an inpatient creatinine retest (%)	74.17	4.91 (-0.47, 9.19)	0.077	7	73	18,436	68.87	-1.07 (-6.82, 2.79)	0.412	5	42	18,981
Proteinuria check in 365 days (%)	22.30	2.76 (-1.85, 6.37)	0.281	12	99	16,053	21.57	5.46 (1.07, 9.29)	0.014**	7	81	16,208
Creatinine test in 90 days (%)	62.21	-1.14 (-7.08, 3.57)	0.518	8	89	15,632	55.35	4.51 (-1.58, 9.64)	0.159	5	90	16,318
Blood pressure measurement in 90 days (%)	36.16	2.06 (-2.39, 6.15)	0.389	11	185	20,133	34.90	4.64 (0.34, 8.82)	0.034*	8	113	19,336
AKI coding in primary care in 90 days (%)	4.24	1.46 (-0.61, 1.94)	0.309	32	57	87,617	2.72	1.17 (-0.40, 1.97)	0.195	17	49	40,783
AKI coding for hospitalization (%)	19.42	9.80 (4.85, 12.11)	<0.001***	14	41	21,896	13.97	3.79 (-0.58, 5.70)	0.11	10	30	23,552
Any PPI prescription in 365 days (%)	45.29	2.06 (-2.83, 6.80)	0.419	11	128	16,478	44.75	0.10 (-4.62, 4.13)	0.913	9	91	17,835
Any NSAID prescription in 365 days (%)	7.83	0.54 (-1.90, 3.13)	0.634	12	124	16,418	11.15	0.40 (-2.59, 3.67)	0.735	7	86	15,045
RAASI continuation in 180 days among prior users (%)	81.56	2.56 (-2.83, 9.08)	0.303	13	73	7,122	80.47	0.60 (-5.22, 7.14)	0.76	10	75	6,858
Diuretic continuation in 180 days among prior users (%)	80.91	3.21 (-3.05, 9.57)	0.312	21	84	5,488	79.04	5.45 (-0.99, 11.98)	0.097	14	92	5,732

E

	With major surgery during admission						Without major surgery during admission					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size		Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	
		Estimate (95% CI)	P value					Estimate (95% CI)	P value			
Primary outcome												
Death in 365 days (%)	26.74	10.14 (-10.73, 27.76)	0.386	23	107	1,223	29.84	1.28 (-3.12, 4.73)	0.688	5	46	34,332
Emergency hospitalization or emergency department attendance in 180 days (%)	41.56	3.85 (-21.37, 28.03)	0.792	24	68	1,160	53.08	0.01 (-3.94, 4.10)	0.968	6	98	33,534
Secondary outcome												
<i>Health outcome</i>												
Death in 30 days (%)	9.30	6.23 (-7.58, 19.44)	0.39	27	77	1,624	11.68	0.32 (-2.45, 2.42)	0.991	7	45	36,744
Emergency hospitalization or emergency department attendance in 30 days (%)	18.18	-0.90 (-21.88, 19.29)	0.902	26	65	1,269	26.07	-0.42 (-3.39, 2.48)	0.761	8	119	38,951
<i>Kidney recovery outcome</i>												
Peak inpatient creatinine after test (micromol/L)	132.36	16.90 (-22.49, 34.64)	0.677	37	87	2,938	106.48	6.85 (-0.02, 11.59)	0.051	7	84	33,564
Updated eGFR at 90 days (ml/min/1.73m ²)	82.30	12.30 (1.23, 30.54)	0.034*	21	56	799	80.02	-0.54 (-2.90, 2.41)	0.857	5	41	23,394
<i>Care process outcome</i>												
Length of inpatient stay >=7 days (%)	96.34	2.84 (-6.11, 10.95)	0.578	20	75	991	46.75	-3.02 (-8.05, 0.86)	0.114	5	49	32,448
Having an inpatient creatinine retest (%)	96.51	2.10 (-6.39, 10.27)	0.648	23	76	1,283	70.72	0.69 (-3.94, 4.13)	0.963	5	44	31,616
Proteinuria check in 365 days (%)	27.87	-6.18 (-28.84, 11.21)	0.388	36	69	2,126	21.78	4.65 (1.07, 7.62)	0.009**	7	85	28,214
Creatinine test in 90 days (%)	71.43	-27.00 (-55.49, -8.76)	0.007**	29	65	1,585	57.86	1.92 (-2.90, 5.78)	0.516	6	76	28,412
Blood pressure measurement in 90 days (%)	34.29	-4.22 (-24.46, 14.24)	0.605	33	97	1,985	35.39	3.78 (0.57, 6.79)	0.020*	9	153	37,622
AKI coding in primary care in 90 days (%)	4.23	-2.36 (-6.18, 0.89)	0.142	39	40	2,760	3.29	0.77 (-0.63, 1.35)	0.473	17	44	71,463
AKI coding for hospitalization (%)	14.29	14.97 (-3.14, 32.17)	0.107	34	65	2,220	16.09	5.79 (2.11, 7.48)	<0.001***	10	29	35,830
Any PPI prescription in 365 days (%)	52.46	-2.93 (-31.15, 22.80)	0.761	25	82	1,064	44.88	2.12 (-1.62, 5.45)	0.288	8	129	31,689
Any NSAID prescription in 365 days (%)	9.84	-10.92 (-25.27, 2.18)	0.099	27	75	1,232	9.91	0.33 (-1.63, 2.63)	0.646	8	98	28,722
RAASI continuation in 180 days among prior users (%)	76.67	9.09 (-13.69, 39.54)	0.341	35	107	636	80.99	2.29 (-2.12, 7.42)	0.277	10	68	11,804
Diuretic continuation in 180 days among prior users (%)	75.00	-16.43 (-65.22, 19.64)	0.292	50	63	485	79.76	4.93 (-0.10, 10.01)	0.055	14	81	9,211

F

	With history of vascular disease						Without history of vascular disease					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size		Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	
		Estimate (95% CI)	P value					Estimate (95% CI)	P value			
Primary outcome												
Death in 365 days (%)	39.53	4.22 (-1.90, 9.13)	0.199	8	69	16,796	24.85	-1.90 (-6.70, 1.91)	0.276	6	49	22,229
Emergency hospitalization or emergency department attendance in 180 days (%)	57.83	0.82 (-4.13, 5.85)	0.735	10	135	16,419	50.72	-0.11 (-4.28, 4.28)	1	7	104	25,400
Secondary outcome												
<i>Health outcome</i>												
Death in 30 days (%)	15.92	1.23 (-2.95, 4.20)	0.733	11	61	19,798	9.48	0.13 (-2.66, 2.20)	0.855	8	48	25,620
Emergency hospitalization or emergency department attendance in 30 days (%)	28.67	-1.64 (-6.40, 2.37)	0.367	12	104	17,042	24.75	-0.08 (-3.19, 3.14)	0.989	9	120	27,644
<i>Kidney recovery outcome</i>												
Peak inpatient creatinine after test (micromol/L)	122.21	7.50 (-2.34, 14.43)	0.158	10	89	15,709	97.46	3.88 (-3.65, 8.81)	0.417	8	81	21,569
Updated eGFR at 90 days (ml/min/1.73m ²)	69.40	-2.53 (-5.58, 1.15)	0.197	8	48	9,921	85.45	-0.02 (-2.60, 3.35)	0.805	6	47	15,841
<i>Care process outcome</i>												
Length of inpatient stay >=7 days (%)	57.20	4.03 (-2.28, 8.96)	0.244	8	79	15,583	42.29	-5.38 (-11.00, -1.00)	0.019*	6	50	21,161
Having an inpatient creatinine retest (%)	79.19	4.40 (-0.74, 7.96)	0.103	8	99	17,003	66.75	0.12 (-5.37, 4.28)	0.824	5	43	21,483
Proteinuria check in 365 days (%)	26.21	3.74 (-1.30, 7.98)	0.159	13	127	12,898	20.06	3.56 (-0.36, 6.86)	0.078	8	89	19,730
Creatinine test in 90 days (%)	67.18	-0.60 (-6.48, 3.87)	0.621	11	106	13,840	53.86	1.12 (-4.13, 5.38)	0.796	6	76	20,355
Blood pressure measurement in 90 days (%)	43.92	3.27 (-1.22, 8.15)	0.147	14	88	17,790	31.56	2.84 (-0.78, 6.11)	0.13	9	134	27,036
AKI coding in primary care in 90 days (%)	5.03	2.20 (-0.38, 3.18)	0.123	26	60	41,333	2.52	0.71 (-0.68, 1.24)	0.568	21	48	68,995
AKI coding for hospitalization (%)	24.87	8.03 (1.88, 9.95)	0.004**	17	42	23,597	11.98	6.50 (2.82, 8.18)	<0.001***	11	30	26,722
Any PPI prescription in 365 days (%)	55.43	0.96 (-3.84, 5.94)	0.673	15	121	14,757	40.67	0.73 (-3.09, 4.19)	0.767	10	106	25,334
Any NSAID prescription in 365 days (%)	4.32	0.74 (-1.31, 2.82)	0.473	13	130	12,911	12.19	0.11 (-2.34, 3.02)	0.805	9	83	21,119
RAASI continuation in 180 days among prior users (%)	81.03	2.66 (-2.99, 9.37)	0.312	13	65	7,030	80.84	-0.46 (-5.71, 5.73)	0.997	11	82	7,080
Diuretic continuation in 180 days among prior users (%)	83.17	5.74 (0.77, 11.24)	0.025*	17	75	7,312	73.32	3.83 (-4.07, 11.51)	0.349	18	119	4,692

G

	With history of diabetes						Without history of diabetes					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Primary outcome												
Death in 365 days (%)	30.81	-0.48 (-6.38, 4.04)	0.659	10	73	15,833	29.43	0.24 (-4.67, 4.09)	0.896	6	45	23,950
Emergency hospitalization or emergency department attendance in 180 days (%)	55.32	0.84 (-4.46, 6.21)	0.748	11	90	14,853	52.08	0.62 (-3.50, 4.82)	0.756	7	116	25,270
Secondary outcome												
Heath outcome												
Death in 30 days (%)	11.03	0.85 (-2.77, 3.16)	0.897	15	60	21,150	11.90	0.60 (-2.57, 3.02)	0.876	7	47	26,018
Emergency hospitalization or emergency department attendance in 30 days (%)	25.69	1.77 (-2.69, 6.37)	0.426	12	105	16,056	26.11	-1.56 (-4.77, 1.45)	0.294	9	127	29,306
Kidney recovery outcome												
Peak inpatient creatinine after test (micromol/L)	118.93	0.26 (-11.21, 8.33)	0.773	11	94	14,318	101.70	9.31 (2.53, 13.96)	0.005**	7	80	24,017
Updated eGFR at 90 days (ml/min/1.73m ²)	74.21	-2.01 (-5.35, 2.18)	0.41	9	65	10,259	82.48	-0.02 (-2.54, 3.28)	0.804	6	43	16,235
Care process outcome												
Length of inpatient stay >=7 days (%)	49.31	0.88 (-5.31, 5.52)	0.97	11	96	15,418	46.40	-2.17 (-7.69, 2.13)	0.267	5	45	22,091
Having an inpatient creatinine retest (%)	76.05	4.74 (-1.10, 8.51)	0.13	9	98	14,864	69.01	0.32 (-4.75, 4.18)	0.901	5	43	23,644
Proteinuria check in 365 days (%)	43.18	3.25 (-2.15, 8.01)	0.258	15	134	14,175	13.85	2.65 (-0.29, 5.19)	0.079	9	115	22,054
Creatinine test in 90 days (%)	65.05	1.61 (-4.25, 6.24)	0.71	12	114	13,573	55.26	1.90 (-3.21, 6.02)	0.552	6	73	20,954
Blood pressure measurement in 90 days (%)	41.96	2.33 (-1.83, 6.36)	0.278	19	132	21,819	32.85	3.70 (0.21, 7.15)	0.037*	9	136	26,073
AKI coding in primary care in 90 days (%)	5.08	2.83 (-0.06, 3.20)	0.059	36	65	73,407	2.62	0.76 (-0.59, 1.42)	0.422	18	48	53,579
AKI coding for hospitalization (%)	23.96	8.40 (1.85, 9.69)	0.004**	21	43	25,755	13.07	6.72 (2.97, 8.53)	<0.001***	10	30	28,496
Any PPI prescription in 365 days (%)	50.14	1.02 (-3.99, 5.85)	0.711	15	180	15,415	43.01	0.59 (-3.47, 4.18)	0.855	9	119	23,881
Any NSAID prescription in 365 days (%)	7.97	0.87 (-1.98, 4.08)	0.497	12	115	11,544	10.64	-0.20 (-2.48, 2.47)	0.997	8	101	21,770
RAASI continuation in 180 days among prior users (%)	82.22	1.33 (-4.24, 7.71)	0.569	14	75	6,763	79.98	1.62 (-3.68, 7.96)	0.471	11	80	7,276
Diuretic continuation in 180 days among prior users (%)	81.03	5.00 (-1.82, 12.30)	0.146	18	104	5,014	78.85	4.51 (-1.30, 10.43)	0.127	15	87	6,479

Table S5: Effects of AKI alerts in emergency hospitalization setting by (A) alert approach, (B) resident neighbourhood measures of deprivation (WIMD), (C) rurality, (D) sex, (E) presence of major surgery during the admission, (F) history of vascular disease and (G) diabetes.

A

	Most deprived quintile					Least deprived quintile						
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Primary outcome												
Death in 365 days (%)	26.13	-3.43 (-12.59, 3.56)	0.273	11	67	3,558	26.10	6.54 (-2.26, 14.26)	0.155	15	86	3,758
Emergency hospitalization or emergency department attendance in 180 days (%)	64.41	3.94 (-4.17, 10.46)	0.399	13	118	4,511	59.43	-0.90 (-11.12, 6.95)	0.652	15	87	3,333
Secondary outcome												
Heath outcome												
Death in 30 days (%)	6.91	-0.04 (-4.87, 3.44)	0.736	18	59	6,892	6.80	4.17 (-1.53, 7.91)	0.185	22	68	8,332
Emergency hospitalization or emergency department attendance in 30 days (%)	45.05	0.47 (-9.54, 6.97)	0.761	13	61	4,405	40.13	1.10 (-8.95, 8.24)	0.935	17	86	4,711
Kidney recovery outcome												
Updated eGFR at 90 days (ml/min/1.73m ²)	84.76	1.96 (-3.18, 9.20)	0.34	11	54	2,121	79.06	-2.00 (-7.40, 6.34)	0.88	11	41	1,599
Care process outcome												
Proteinuria check in 365 days (%)	22.76	1.72 (-6.36, 8.24)	0.801	16	76	4,787	21.66	-0.78 (-12.06, 7.22)	0.623	15	49	2,813
Creatinine test in 90 days (%)	66.32	-1.11 (-10.44, 6.12)	0.609	13	86	3,866	70.10	1.10 (-10.35, 9.06)	0.897	15	67	2,884
Blood pressure measurement in 90 days (%)	36.63	-1.29 (-9.05, 5.31)	0.609	17	88	6,201	37.94	-0.24 (-10.31, 9.89)	0.968	15	75	2,890
AKI coding in primary care in 90 days (%)	2.26	2.71 (-0.27, 5.07)	0.078	26	81	15,626	2.01	6.59 (2.43, 9.72)	0.001**	31	71	19,103
Any PPI prescription in 365 days (%)	53.86	0.61 (-7.01, 8.75)	0.829	19	70	6,906	41.84	1.40 (-8.80, 12.11)	0.756	17	79	3,206
Any NSAID prescription in 365 days (%)	11.18	0.38 (-5.41, 6.67)	0.838	13	80	2,965	8.01	2.95 (-2.42, 10.34)	0.224	16	58	2,827
RAASI continuation in 180 days among prior users (%)	86.31	-4.91 (-14.29, 6.46)	0.459	18	71	1,541	87.63	3.99 (-5.47, 16.10)	0.334	20	93	1,540
Diuretic continuation in 180 days among prior users (%)	83.76	3.91 (-8.46, 15.53)	0.564	23	72	1,940	85.07	0.59 (-12.36, 12.34)	0.999	31	108	2,309

B

	Rural						Urban					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	P value	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	P value
		Estimate (95% CI)	P value					Estimate (95% CI)	P value			
Primary outcome												
Death in 365 days (%)	26.17	0.38 (-6.60, 5.75)	0.893	12	76	6,119	24.86	2.42 (-3.96, 7.75)	0.525	7	70	7,002
Emergency hospitalization or emergency department attendance in 180 days (%)	58.49	4.79 (-2.82, 10.02)	0.271	13	85	6,150	62.61	3.66 (-3.04, 8.50)	0.353	9	68	7,693
Secondary outcome												
Heath outcome												
Death in 30 days (%)	6.52	0.21 (-3.32, 2.61)	0.814	20	65	13,384	6.70	0.86 (-2.26, 2.88)	0.815	14	56	14,391
Emergency hospitalization or emergency department attendance in 30 days (%)	37.20	5.46 (-2.25, 10.42)	0.206	14	71	6,814	41.70	2.07 (-4.97, 6.80)	0.761	9	50	8,276
Kidney recovery outcome												
Updated eGFR at 90 days (ml/min/1.73m2)	82.31	-1.99 (-5.91, 4.23)	0.746	11	44	3,507	81.11	1.14 (-2.82, 6.45)	0.442	7	45	4,138
Care process outcome												
Proteinuria check in 365 days (%)	21.17	3.70 (-2.70, 9.12)	0.287	16	87	7,299	21.67	0.39 (-5.44, 5.02)	0.937	11	70	6,889
Creatinine test in 90 days (%)	63.36	9.56 (2.29, 15.71)	0.009**	12	102	5,292	67.85	-2.27 (-9.60, 2.70)	0.272	9	58	6,360
Blood pressure measurement in 90 days (%)	35.99	4.45 (-2.59, 10.32)	0.241	15	79	6,647	36.87	-2.94 (-8.88, 1.76)	0.19	12	75	9,019
AKI coding in primary care in 90 days (%)	1.20	3.70 (1.21, 5.68)	0.002**	26	70	23,252	2.39	3.20 (1.22, 4.32)	<0.001***	29	78	62,094
Any PPI prescription in 365 days (%)	45.19	-1.92 (-9.08, 4.63)	0.525	16	86	7,298	49.16	-0.99 (-6.62, 4.29)	0.676	13	110	8,776
Any NSAID prescription in 365 days (%)	8.70	2.37 (-1.11, 6.60)	0.163	14	120	5,730	9.12	2.80 (-0.54, 6.88)	0.094	10	69	6,135
RAASI continuation in 180 days among prior users (%)	87.39	-3.95 (-11.66, 5.22)	0.454	19	75	2,616	85.21	-0.64 (-7.53, 6.41)	0.875	12	95	2,863
Diuretic continuation in 180 days among prior users (%)	81.66	0.40 (-9.94, 9.77)	0.987	25	61	3,160	83.61	2.67 (-4.13, 9.12)	0.461	19	107	3,989

C

	Male						Female					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	P value	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)	Effective sample size	P value
		Estimate (95% CI)	P value					Estimate (95% CI)	P value			
Primary outcome												
Death in 365 days (%)	30.08	1.54 (-5.51, 8.15)	0.704	10	85	5,524	22.59	0.13 (-6.61, 5.79)	0.897	7	61	5,662
Emergency hospitalization or emergency department attendance in 180 days (%)	64.08	3.50 (-4.50, 9.14)	0.505	11	67	5,427	59.52	3.32 (-3.60, 8.95)	0.404	8	88	6,471
Secondary outcome												
Heath outcome												
Death in 30 days (%)	7.57	1.92 (-1.78, 4.64)	0.383	17	77	10,906	6.10	-0.33 (-3.55, 1.88)	0.547	12	52	9,820
Emergency hospitalization or emergency department attendance in 30 days (%)	45.03	1.59 (-6.61, 6.88)	0.969	12	57	6,460	37.35	3.32 (-3.95, 8.70)	0.462	8	56	7,002
Kidney recovery outcome												
Updated eGFR at 90 days (ml/min/1.73m2)	82.40	1.38 (-3.63, 7.69)	0.481	9	54	3,041	80.96	0.09 (-3.48, 5.45)	0.666	8	35	4,080
Care process outcome												
Proteinuria check in 365 days (%)	21.12	2.59 (-3.94, 8.13)	0.496	14	108	5,144	21.70	0.76 (-5.20, 5.73)	0.924	10	73	6,691
Creatinine test in 90 days (%)	72.09	-2.37 (-10.96, 4.05)	0.367	10	77	4,496	63.10	4.83 (-2.64, 10.44)	0.243	8	64	5,404
Blood pressure measurement in 90 days (%)	40.64	-3.27 (-9.50, 2.38)	0.24	16	101	8,458	34.31	1.75 (-4.44, 6.35)	0.729	11	58	8,659
AKI coding in primary care in 90 days (%)	2.67	4.24 (1.77, 6.16)	<0.001***	30	76	35,855	1.59	2.79 (0.72, 4.00)	0.005**	25	61	41,765
Any PPI prescription in 365 days (%)	48.63	-2.17 (-9.35, 5.44)	0.605	14	76	5,657	47.37	0.04 (-5.97, 5.27)	0.903	12	109	8,661
Any NSAID prescription in 365 days (%)	6.65	4.59 (0.92, 9.07)	0.016*	13	69	5,076	10.18	3.65 (-0.02, 7.99)	0.051	8	74	5,266
RAASI continuation in 180 days among prior users (%)	85.67	-0.69 (-7.92, 7.88)	0.996	15	70	2,309	86.06	-3.39 (-10.97, 3.92)	0.354	13	98	2,534
Diuretic continuation in 180 days among prior users (%)	83.07	5.69 (-2.70, 14.05)	0.184	21	58	2,434	82.94	-0.70 (-8.64, 5.86)	0.707	19	129	3,604

D

	With history of vascular disease						Without history of vascular disease					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Primary outcome												
Death in 365 days (%)	31.00	3.57 (-3.29, 10.76)	0.297	11	77	4,850	22.73	-0.79 (-3.08, 4.12)	0.605	8	59	6,790
Emergency hospitalization or emergency department attendance in 180 days (%)	65.71	1.97 (-5.52, 8.12)	0.709	11	89	5,356	59.12	4.73 (-1.75, 9.08)	0.185	10	63	8,454
Secondary outcome												
Heath outcome												
Death in 30 days (%)	7.96	2.51 (-1.36, 5.74)	0.227	18	77	9,321	6.03	-0.90 (-3.76, 0.88)	0.223	15	51	14,957
Emergency hospitalization or emergency department attendance in 30 days (%)	43.95	1.62 (-6.57, 7.83)	0.865	12	76	5,302	38.42	3.35 (-3.37, 7.55)	0.454	10	46	9,225
Kidney recovery outcome												
Updated eGFR at 90 days (ml/min/1.73m ²)	66.89	1.71 (-2.28, 7.09)	0.314	10	50	3,004	88.91	-1.59 (-5.43, 3.43)	0.659	7	46	4,426
Care process outcome												
Proteinuria check in 365 days (%)	27.54	-0.92 (-8.50, 5.84)	0.717	14	83	4,373	19.05	3.00 (-2.20, 7.13)	0.301	12	77	9,252
Creatinine test in 90 days (%)	74.72	-1.64 (-9.73, 4.06)	0.42	12	77	4,364	62.62	4.25 (-2.62, 9.34)	0.271	9	76	7,375
Blood pressure measurement in 90 days (%)	47.56	0.20 (-6.40, 5.88)	0.934	20	79	9,769	31.77	-0.20 (-5.17, 4.18)	0.836	13	86	10,630
AKI coding in primary care in 90 days (%)	2.88	4.56 (1.50, 6.83)	0.002**	25	78	16,468	1.58	2.64 (0.89, 3.81)	0.002**	27	68	62,326
Any PPI prescription in 365 days (%)	55.38	-2.47 (-10.20, 5.24)	0.529	15	77	5,377	44.72	0.11 (-4.22, 5.04)	0.863	17	74	17,370
Any NSAID prescription in 365 days (%)	3.69	1.17 (-1.60, 4.40)	0.361	15	86	4,867	11.12	3.85 (0.43, 8.04)	0.029*	10	70	7,198
RAASI continuation in 180 days among prior users (%)	86.47	-2.95 (-9.98, 3.70)	0.368	14	107	2,652	85.15	-2.00 (-9.79, 7.32)	0.777	16	58	2,575
Diuretic continuation in 180 days among prior users (%)	88.34	1.61 (-4.83, 7.65)	0.658	20	72	4,132	73.12	0.96 (-8.87, 9.89)	0.915	27	109	4,776

E

	With history of diabetes						Without history of diabetes					
	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth (micromol/L)		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Primary outcome												
Death in 365 days (%)	28.11	2.55 (-4.96, 10.35)	0.49	12	71	4,144	24.39	0.80 (-5.20, 5.55)	0.949	7	59	7,556
Emergency hospitalization or emergency department attendance in 180 days (%)	67.87	2.89 (-6.21, 9.78)	0.662	11	73	4,153	58.98	4.24 (-1.73, 8.64)	0.192	9	79	9,534
Secondary outcome												
Kidney recovery outcome												
Last eGFR in 90 days (ml/min/1.73m ²)	74.51	4.11 (-1.79, 11.73)	0.149	9	53	2,225	84.05	-0.85 (-4.14, 3.84)	0.942	8	39	4,863
Care process outcome												
Proteinuria check in 365 days (%)	46.18	-1.28 (-9.42, 6.85)	0.757	18	70	5,428	13.77	2.51 (-1.36, 5.83)	0.224	14	83	12,722
Creatinine test in 90 days (%)	73.13	1.99 (-6.99, 9.53)	0.763	11	96	3,542	64.09	2.04 (-4.61, 6.63)	0.725	9	59	7,963
Blood pressure measurement in 90 days (%)	46.25	-1.54 (-8.97, 5.20)	0.602	18	98	6,451	33.45	-0.35 (-5.74, 4.05)	0.735	11	76	10,212
AKI coding in primary care in 90 days (%)	3.13	5.31 (1.84, 8.34)	0.002**	24	64	13,103	1.60	2.61 (0.86, 3.68)	0.002**	28	63	73,854
PPI prescription in 365 days (%)	52.70	3.07 (-3.42, 10.91)	0.306	24	62	11,335	46.27	-1.52 (-6.69, 3.09)	0.471	14	96	11,316
NSAID prescription in 365 days (%)	8.01	1.36 (-2.97, 6.22)	0.489	15	99	3,989	9.28	3.64 (0.69, 7.47)	0.018*	10	63	7,709
RAASI prescription in 180 days (%)	86.42	-0.88 (-8.13, 8.17)	0.996	16	60	2,352	85.58	-3.53 (-11.11, 3.37)	0.295	14	75	2,934
Diuretic prescription in 180 days (%)	83.89	2.12 (-6.18, 9.63)	0.669	30	81	4,638	82.52	0.73 (-7.02, 6.95)	0.992	20	95	3,972

Table S6: Effects of AKI alerts in community setting by (A) resident neighbourhood measures of deprivation (WIMD), (B) rurality (C) sex, (D) history of vascular disease and (E) diabetes.

	Hospital setting					Community setting						
	Mean below threshold	Effect of AKI alert		Bandwidth		Effective sample size	Mean below threshold	Effect of AKI alert		Bandwidth		Effective sample size
		Estimate (95% CI)	P value	below	above			Estimate (95% CI)	P value	below	above	
Primary outcome												
Death in 365 days (%)	29.81	1.31 (-3.07, 4.74)	0.674	5	45	34,685	25.32	2.07 (-3.44, 6.65)	0.534	7	70	9,590
Emergency hospitalization or emergency department attendance in 180 days (%)	52.98	0.13 (-3.82, 4.21)	0.924	6	97	33,793	61.18	4.07 (-1.84, 8.27)	0.212	8	68	10,592
Secondary outcome												
<i>Health outcome</i>												
Death in 30 days (%)	11.66	0.43 (-2.33, 2.52)	0.941	7	45	37,112	6.64	0.58 (-2.02, 2.30)	0.9	13	55	17,598
Emergency hospitalization or emergency department attendance in 30 days (%)	26.00	-0.35 (-3.33, 2.54)	0.791	8	116	39,261	40.15	2.98 (-3.11, 7.12)	0.442	8	48	11,284
<i>Kidney recovery outcome</i>												
Peak inpatient creatinine after test (micromol/L)	106.80	6.45 (-0.46, 11.16)	0.071	7	84	33,993						
Updated eGFR at 90 days (ml/min/1.73m ²)	80.04	-0.40 (-2.75, 2.55)	0.941	5	40	23,648	81.51	0.80 (-2.48, 5.30)	0.479	7	38	6,130
<i>Care process outcome</i>												
Length of inpatient stay >=7 days (%)	47.21	-3.09 (-8.12, 0.78)	0.106	5	49	32,779						
Having an inpatient creatinine retest (%)	70.95	0.80 (-3.80, 4.21)	0.919	5	44	31,945						
Proteinuria check in 365 days (%)	21.84	4.33 (0.76, 7.28)	0.016*	7	85	28,551	21.50	1.15 (-3.76, 5.03)	0.777	10	73	10,417
Creatinine test in 90 days (%)	57.98	1.71 (-3.12, 5.54)	0.584	6	73	28,577	66.29	3.59 (-2.67, 8.05)	0.326	7	66	8,788
Blood pressure measurement in 90 days (%)	35.38	3.79 (0.58, 6.79)	0.020*	9	150	37,947	36.56	-0.56 (-5.63, 3.42)	0.633	10	76	12,225
AKI coding in primary care in 90 days (%)	3.30	0.72 (-0.67, 1.30)	0.527	17	44	72,088	1.97	3.27 (1.56, 4.25)	<0.001***	25	64	61,535
AKI coding for hospitalization (%)	16.08	5.88 (2.22, 7.58)	<0.001***	9	29	36,152						
Any PPI prescription in 365 days (%)	44.95	2.10 (-1.65, 5.43)	0.295	8	130	32,021	47.80	-1.52 (-6.28, 3.07)	0.501	12	86	11,762
Any NSAID prescription in 365 days (%)	9.91	0.23 (-1.73, 2.53)	0.712	8	98	29,016	8.97	3.19 (0.57, 6.56)	0.020*	9	63	9,215
RAASI continuation in 180 days among prior users (%)	80.93	2.29 (-2.09, 7.39)	0.274	10	68	11,913	85.90	-2.71 (-8.74, 3.39)	0.387	11	68	3,779
Diuretic continuation in 180 days among prior users (%)	79.72	5.07 (0.01, 10.18)	0.050*	13	81	9,286	82.99	1.76 (-4.94, 6.73)	0.764	18	84	5,457

Table S7: Effect of AKI alert, including the MSE-optimal bandwidths^{1,2,3}

¹Mean below threshold is calculated as the mean within the 5 micromol/L below the threshold.

² Effective sample size is the number of observations within the MSE-optimal bandwidth.

³ Abbreviations: AKI = acute kidney injury

Outcome	Robustness check	Effect of AKI alert			
		Hospital setting		Community setting	
		Estimate (95% CI)	P value	Estimate (95% CI)	P value
Death in 365 days (%)	Main specification	1.31 (-3.07, 4.74)	0.674	2.07 (-3.44, 6.65)	0.534
	Quadratic fit	0.18 (-2.67, 2.71)	0.989	1.72 (-1.93, 5.48)	0.348
	0.75*MSE-optimal bandwidth	1.76 (-3.82, 6.58)	0.604	3.16 (-3.58, 9.07)	0.395
	1.25*MSE-optimal bandwidth	1.22 (-2.65, 4.04)	0.683	1.88 (-3.13, 5.61)	0.577
	1.5*MSE-optimal bandwidth	1.16 (-2.44, 3.53)	0.72	1.91 (-2.99, 4.82)	0.645
	2*MSE-optimal bandwidth	1.78 (-1.56, 3.34)	0.478	2.84 (-1.93, 4.59)	0.423
	Alternative index test	3.29 (-1.33, 7.03)	0.182		
Emergency hospitalization or emergency department attendance in 180 days (%)	Main specification	0.13 (-3.82, 4.21)	0.924	4.07 (-1.84, 8.27)	0.212
	Quadratic fit	0.95 (-1.62, 3.76)	0.435	3.06 (-1.31, 6.45)	0.195
	0.75*MSE-optimal bandwidth	-0.25 (-5.06, 4.67)	0.938	3.29 (-3.54, 8.84)	0.401
	1.25*MSE-optimal bandwidth	0.53 (-2.79, 3.97)	0.731	4.42 (-0.81, 7.91)	0.111
	1.5*MSE-optimal bandwidth	0.25 (-2.76, 3.28)	0.864	4.92 (0.15, 7.90)	0.042*
	2*MSE-optimal bandwidth	0.23 (-2.37, 2.61)	0.924	5.79 (1.44, 7.91)	0.005**
	Alternative index test	-0.68 (-4.68, 3.53)	0.784		
Death in 30 days (%)	Main specification	0.43 (-2.33, 2.52)	0.941	0.58 (-2.02, 2.30)	0.9
	Quadratic fit	0.26 (-1.75, 2.08)	0.868	0.11 (-2.09, 2.08)	0.993
	0.75*MSE-optimal bandwidth	0.56 (-2.67, 3.40)	0.816	0.48 (-2.46, 2.61)	0.954
	1.25*MSE-optimal bandwidth	0.66 (-1.91, 2.27)	0.866	0.80 (-1.65, 2.17)	0.787
	1.5*MSE-optimal bandwidth	0.95 (-1.53, 2.20)	0.723	1.12 (-1.26, 2.21)	0.594
	2*MSE-optimal bandwidth	1.24 (-1.13, 1.96)	0.595	1.65 (-0.82, 2.20)	0.37
	Alternative index test	1.75 (-1.74, 4.42)	0.393		
Emergency hospitalization or emergency department attendance in 30 days (%)	Main specification	-0.35 (-3.33, 2.54)	0.791	2.98 (-3.11, 7.12)	0.442
	Quadratic fit	-0.39 (-2.47, 1.73)	0.73	2.12 (-2.59, 5.89)	0.447
	0.75*MSE-optimal bandwidth	-0.81 (-4.44, 2.71)	0.635	1.86 (-5.00, 7.46)	0.699
	1.25*MSE-optimal bandwidth	-0.36 (-2.94, 2.11)	0.745	3.26 (-2.39, 6.47)	0.366
	1.5*MSE-optimal bandwidth	-0.23 (-2.57, 1.90)	0.77	3.86 (-1.41, 6.45)	0.209
	2*MSE-optimal bandwidth	0.00 (-2.00, 1.69)	0.865	4.96 (0.14, 6.72)	0.041*
	Alternative index test	0.44 (-2.75, 3.73)	0.767		
Having an inpatient creatinine retest (%)	Main specification	0.80 (-3.80, 4.21)	0.919		
	Quadratic fit	1.88 (-1.01, 4.49)	0.216		
	0.75*MSE-optimal bandwidth	0.31 (-5.31, 4.67)	0.9		
	1.25*MSE-optimal bandwidth	1.50 (-2.42, 4.20)	0.599		
	1.5*MSE-optimal bandwidth	3.13 (-0.42, 5.38)	0.093		
	2*MSE-optimal bandwidth	4.76 (1.41, 6.22)	0.002**		
	Alternative index test	5.13 (1.55, 7.68)	0.003**		
Updated eGFR at 90 days (ml/min/1.73m2)	Main specification	-0.40 (-2.75, 2.55)	0.941	0.80 (-2.48, 5.30)	0.479
	Quadratic fit	0.10 (-1.57, 2.12)	0.772	1.01 (-1.93, 4.26)	0.46
	0.75*MSE-optimal bandwidth	0.65 (-2.66, 4.40)	0.629	0.96 (-3.32, 6.35)	0.539
	1.25*MSE-optimal bandwidth	-1.20 (-3.10, 1.45)	0.478	0.07 (-2.57, 4.17)	0.643
	1.5*MSE-optimal bandwidth	-1.71 (-3.27, 0.81)	0.238	-0.76 (-2.92, 3.13)	0.946
	2*MSE-optimal bandwidth	-2.50 (-3.41, -0.07)	0.042*	-2.42 (-3.63, 1.39)	0.381
	Alternative index test	-1.35 (-3.59, 1.70)	0.485		

Peak inpatient creatinine after test (micromol/L)	Main specification	6.45 (-0.46, 11.16)	0.071		
	Quadratic fit	2.88 (-2.12, 7.19)	0.285		
	0.75*MSE-optimal bandwidth	6.75 (-1.23, 13.34)	0.103		
	1.25*MSE-optimal bandwidth	6.01 (-0.35, 9.85)	0.068		
	1.5*MSE-optimal bandwidth	6.16 (-0.06, 9.10)	0.053		
	2*MSE-optimal bandwidth	6.86 (0.62, 8.36)	0.023*		
	Alternative index test	7.50 (0.53, 11.95)	0.032*		
Diuretic continuation in 180 days among prior users (%)	Main specification	5.07 (0.01, 10.18)	0.050*	1.76 (-4.94, 6.73)	0.764
	Quadratic fit	4.22 (-0.84, 9.25)	0.102	-0.88 (-8.00, 4.84)	0.629
	0.75*MSE-optimal bandwidth	4.45 (-1.75, 10.44)	0.162	2.02 (-5.54, 8.11)	0.712
	1.25*MSE-optimal bandwidth	5.05 (0.69, 9.53)	0.024*	1.24 (-4.48, 5.80)	0.802
	1.5*MSE-optimal bandwidth	4.14 (0.28, 8.14)	0.036*	1.13 (-3.94, 5.38)	0.761
	2*MSE-optimal bandwidth	2.66 (-0.52, 5.97)	0.1	1.91 (-2.29, 5.91)	0.387
	Inclusion of non-survivors	4.95 (0.51, 9.85)	0.030*	0.25 (-5.07, 5.49)	0.938
	Alternative index test	5.53 (-0.02, 11.27)	0.051		
Any NSAID prescription in 365 days (%)	Main specification	0.23 (-1.73, 2.53)	0.712	3.19 (0.57, 6.56)	0.020*
	Quadratic fit	-0.24 (-2.19, 1.12)	0.525	3.22 (0.59, 5.99)	0.017*
	0.75*MSE-optimal bandwidth	1.02 (-1.42, 3.72)	0.381	4.20 (0.86, 7.99)	0.015*
	1.25*MSE-optimal bandwidth	-0.01 (-1.63, 2.07)	0.817	2.87 (0.70, 5.99)	0.013*
	1.5*MSE-optimal bandwidth	-0.16 (-1.55, 1.75)	0.906	2.63 (0.76, 5.55)	0.010**
	2*MSE-optimal bandwidth	-0.55 (-1.65, 1.09)	0.691	1.86 (0.41, 4.53)	0.019*
	Inclusion of non-survivors	0.26 (-1.47, 2.25)	0.68	2.24 (0.01, 5.10)	0.050*
	Alternative index test	-0.48 (-2.49, 1.73)	0.725		
Any PPI prescription in 365 days (%)	Main specification	2.10 (-1.65, 5.43)	0.295	-1.52 (-6.28, 3.07)	0.501
	Quadratic fit	0.23 (-2.74, 3.23)	0.872	-1.07 (-5.49, 3.25)	0.615
	0.75*MSE-optimal bandwidth	4.03 (-0.35, 8.21)	0.072	-2.23 (-7.77, 3.45)	0.451
	1.25*MSE-optimal bandwidth	1.32 (-2.07, 4.04)	0.528	-0.78 (-5.11, 3.07)	0.624
	1.5*MSE-optimal bandwidth	1.05 (-2.06, 3.36)	0.638	-0.84 (-4.79, 2.56)	0.554
	2*MSE-optimal bandwidth	1.00 (-1.67, 2.82)	0.615	-0.59 (-4.02, 2.23)	0.573
	Inclusion of non-survivors	2.64 (-0.67, 5.79)	0.121	-1.66 (-5.29, 2.31)	0.443
	Alternative index test	-1.18 (-5.32, 2.53)	0.486		
RAASI continuation in 180 days among prior users (%)	Main specification	2.29 (-2.09, 7.39)	0.274	-2.71 (-8.74, 3.39)	0.387
	Quadratic fit	0.58 (-3.34, 4.55)	0.764	-1.85 (-7.19, 3.21)	0.454
	0.75*MSE-optimal bandwidth	4.25 (-1.21, 10.27)	0.122	-3.66 (-10.99, 3.40)	0.301
	1.25*MSE-optimal bandwidth	1.00 (-2.58, 5.59)	0.47	-2.78 (-7.80, 2.88)	0.366
	1.5*MSE-optimal bandwidth	0.25 (-2.71, 4.55)	0.62	-2.67 (-6.94, 2.74)	0.395
	2*MSE-optimal bandwidth	-0.82 (-2.96, 3.02)	0.983	-1.57 (-4.78, 3.54)	0.77
	Inclusion of non-survivors	2.37 (-2.37, 7.83)	0.294	-6.01 (-12.62, 0.35)	0.064
	Alternative index test	2.41 (-2.53, 8.07)	0.305		
AKI coding for hospitalization (%)	Main specification	5.88 (2.22, 7.58)	<0.001***		
	Quadratic fit	5.87 (3.07, 7.20)	<0.001***		
	0.75*MSE-optimal bandwidth	4.71 (0.78, 7.27)	0.015*		
	1.25*MSE-optimal bandwidth	6.67 (3.14, 7.80)	<0.001***		
	1.5*MSE-optimal bandwidth	7.53 (4.07, 8.22)	<0.001***		
	2*MSE-optimal bandwidth	9.38 (5.96, 9.44)	<0.001***		
	Alternative index test	6.23 (2.50, 7.28)	<0.001***		

AKI coding in primary care in 90 days (%)	Main specification	0.72 (-0.67, 1.30)	0.527	3.27 (1.56, 4.25)	<0.001***
	Quadratic fit	0.67 (-0.44, 1.32)	0.329	3.06 (1.38, 4.28)	<0.001***
	0.75*MSE-optimal bandwidth	0.38 (-1.13, 1.22)	0.935	3.10 (1.45, 4.47)	<0.001***
	1.25*MSE-optimal bandwidth	1.22 (-0.12, 1.61)	0.091	3.46 (1.86, 4.33)	<0.001***
	1.5*MSE-optimal bandwidth	1.63 (0.30, 1.86)	0.006**	3.69 (2.15, 4.50)	<0.001***
	2*MSE-optimal bandwidth	2.47 (1.11, 2.44)	<0.001***	4.26 (2.64, 4.84)	<0.001***
	Inclusion of non-survivors	0.75 (-0.59, 1.32)	0.456	3.35 (1.78, 4.19)	<0.001***
	Alternative index test	0.99 (-0.48, 1.47)	0.322		
Blood pressure measurement in 90 days (%)	Main specification	3.79 (0.58, 6.79)	0.020*	-0.56 (-5.63, 3.42)	0.633
	Quadratic fit	2.92 (0.13, 5.72)	0.040*	-0.97 (-5.38, 3.01)	0.579
	0.75*MSE-optimal bandwidth	4.40 (0.58, 8.09)	0.024*	-1.03 (-6.87, 4.00)	0.605
	1.25*MSE-optimal bandwidth	3.26 (0.43, 5.79)	0.023*	-0.06 (-4.68, 3.24)	0.721
	1.5*MSE-optimal bandwidth	3.04 (0.47, 5.22)	0.019*	0.26 (-3.96, 3.11)	0.813
	2*MSE-optimal bandwidth	2.94 (0.76, 4.69)	0.007**	1.07 (-2.60, 3.40)	0.795
	Inclusion of non-survivors	4.21 (1.20, 7.28)	0.006**	-0.85 (-5.14, 2.90)	0.584
	Alternative index test	3.13 (-0.60, 6.77)	0.101		
Creatinine test in 90 days (%)	Main specification	1.71 (-3.12, 5.54)	0.584	3.59 (-2.67, 8.05)	0.326
	Quadratic fit	0.14 (-3.12, 2.90)	0.944	0.26 (-4.37, 4.01)	0.933
	0.75*MSE-optimal bandwidth	1.76 (-4.09, 6.72)	0.633	4.48 (-2.72, 10.33)	0.253
	1.25*MSE-optimal bandwidth	1.95 (-2.35, 5.14)	0.466	3.07 (-2.65, 6.65)	0.398
	1.5*MSE-optimal bandwidth	1.82 (-2.10, 4.46)	0.48	2.47 (-2.77, 5.53)	0.514
	2*MSE-optimal bandwidth	2.22 (-1.28, 4.13)	0.302	3.01 (-1.60, 5.33)	0.291
	Inclusion of non-survivors	1.57 (-2.92, 5.20)	0.581	3.26 (-2.07, 7.90)	0.251
	Alternative index test	2.62 (-2.44, 6.84)	0.353		
Length of inpatient stay >=7 days (%)	Main specification	-3.09 (-8.12, 0.78)	0.106		
	Quadratic fit	-0.92 (-4.20, 2.10)	0.515		
	0.75*MSE-optimal bandwidth	-2.92 (-9.18, 2.30)	0.24		
	1.25*MSE-optimal bandwidth	-1.67 (-6.07, 1.40)	0.221		
	1.5*MSE-optimal bandwidth	-0.28 (-4.35, 2.29)	0.543		
	2*MSE-optimal bandwidth	1.41 (-2.42, 3.05)	0.822		
	Inclusion of non-survivors	-3.18 (-7.95, 0.61)	0.093		
	Alternative index test	0.90 (-3.86, 4.55)	0.872		
Proteinuria check in 365 days (%)	Main specification	4.33 (0.76, 7.28)	0.016*	1.15 (-3.76, 5.03)	0.777
	Quadratic fit	2.33 (-0.29, 4.63)	0.084	1.31 (-2.66, 4.73)	0.582
	0.75*MSE-optimal bandwidth	4.02 (-0.14, 7.69)	0.059	-0.52 (-6.30, 4.23)	0.701
	1.25*MSE-optimal bandwidth	4.34 (1.10, 6.80)	0.007**	2.24 (-2.16, 5.49)	0.394
	1.5*MSE-optimal bandwidth	4.23 (1.31, 6.35)	0.003**	2.67 (-1.37, 5.47)	0.24
	2*MSE-optimal bandwidth	3.56 (1.05, 5.23)	0.003**	3.08 (-0.61, 5.20)	0.121
	Inclusion of non-survivors	3.73 (0.94, 6.03)	0.007**	1.55 (-1.87, 4.48)	0.42
	Alternative index test	3.50 (-0.27, 6.53)	0.071		

Table S8: Robustness checks for the effects of AKI alert in hospital and community setting^{1,2,3}

¹Alternative index test” refers to the test with the highest reading in the first two days of inpatient stay, and inpatient stays of less than two days are excluded from this specific robustness check.

² Abbreviations: AKI = acute kidney injury