

**Supplementary Table 1. Patient Selection**

Quality Metric	Inclusion criteria	Patients included			Patients excluded		Final cohort		
		Initial number included			Exclusion criteria	Number excluded (%)	N	Uninsured/Medicaid	Insured
		N	Uninsured/Medicaid	Insured					
<b>Bladder</b>									
Appropriate lymph node removal	Patients under 80 with bladder cancer undergoing partial or radical cystectomy	17,297	1,326 (7.7%)	15,971 (92.3%)	Unknown number of lymph nodes resected	418 (2.4%)	16,879	1,304 (7.7%)	15,575 (92.3%)
Appropriate surgery	Patients with clinical T2-4N0M0 bladder cancer	26,759	2,098 (7.8%)	24,661 (92.2%)	Unknown surgery; or unknown chemotherapy and radiation if local tumor destruction/excision	99 (0.4%)	26,660	2,089 (7.8%)	24,571 (92.2%)
Appropriate chemotherapy	Patients with muscle-invasive bladder cancer undergoing radical cystectomy	12,237	968 (7.9%)	11,269 (92.1%)	Unknown chemotherapy details	219 (1.8%)	12,018	947 (7.9%)	11,071 (92.1%)
<b>Breast</b>									
Appropriate radiation after breast conserving surgery	Patients under 70 receiving breast conserving surgery	519,267	46,164 (8.9%)	473,103 (91.1%)	Unknown radiation therapy details	13,209 (2.5%)	506,058	44,846 (8.9%)	461,212 (91.1%)
Appropriate chemotherapy	Patients under 70 with T1cN0M0 or stage IB-III breast cancer	125,130	18,565 (14.8%)	106,565 (85.2%)	Unknown hormone receptor status; or unknown chemotherapy	15,772 (12.6%)	109,358	16,536 (15.1%)	92,822 (84.9%)
Appropriate hormone therapy	Patients with T1cN0M0 or stage IB-III breast cancer	584,809	52,150 (8.9%)	532,659 (91.1%)	Unknown hormone receptor status; or unknown hormone therapy	60,610 (10.4%)	524,199	46,733 (8.9%)	477,466 (91.1%)
Appropriate radiation after mastectomy	Patients with breast cancer undergoing mastectomy	84,041	11,863 (14.1%)	72,178 (85.9%)	Unknown number of lymph nodes resected; or unknown radiation	6,504 (7.7%)	77,537	10,770 (13.9%)	66,767 (86.1%)
<b>Cervix</b>									
Appropriate brachytherapy	Patients with cervical cancer treated with curative intent radiation	32,953	12,326 (37.4%)	20,627 (62.6%)	Unknown brachytherapy	338 (1.0%)	32,615	12,175 (37.3%)	20,440 (62.7%)
Appropriate radiation timing	Patients with cervical cancer treated with radiation	32,296	11,495 (35.6%)	20,801 (64.4%)	Unknown radiation therapy details	1,385 (4.3%)	30,911	10,860 (35.1%)	20,051 (64.9%)
Appropriate chemotherapy and radiation	Patients with stage IB2-IV cervical cancer treated with radiation, or treated with surgery	41,249	15,161 (36.8%)	26,088 (63.2%)	Unknown chemotherapy; or if patients received surgery unknown lymph nodes, or surgical margin status	451 (1.1 %)	40,798	14,998 (36.8%)	25,800 (63.2%)
<b>Colon</b>									
Appropriate chemotherapy	Patients under 80 with stage III (lymph node positive) colon cancer treated with surgery	25,501	2,620 (10.3%)	22,881 (89.7%)	Unknown chemotherapy details	1,707 (6.7%)	23,794	2,470 (10.4%)	21,324 (89.6%)
Appropriate lymph node removal	Patients with colon cancer treated with surgery	114,902	6,116 (5.3%)	108,786 (94.7%)	Unknown number of lymph nodes resected	327 (0.3%)	114,575	6,101 (5.3%)	108,474 (94.7%)
<b>Endometrium</b>									
Appropriate chemotherapy and radiation	Patients with stage IIIC or IV endometrial cancer	27,298	3,247 (11.9%)	24,051 (88.1%)	Unknown chemotherapy; or unknown radiation	128 (0.5%)	27,170	3,230 (11.9%)	23,940 (88.1%)
Appropriate surgery	Patients with stage I-IIIC endometrial cancer	152,668	14,633 (9.6%)	138,035 (90.4%)	Unknown surgery	25 (0.0%)	152,643	14,630 (9.6%)	138,013 (90.4%)
<b>Gastric</b>									

Appropriate lymph node removal	Patients with gastric cancer treated with surgery	32,527	3,782 (11.6%)	28,745 (88.4%)	Unknown number of lymph nodes resected	326 (1.0 %)	32,201	3,736 (11.6%)	28,465 (88.4%)	
<b>Non-Small Cell Lung</b>										
Appropriate lymph node removal	Patients with stage IA-IIIB NSCLC treated with surgery	161,431	9,995 (6.2%)	151,436 (93.8%)	Unknown number of lymph nodes resected	10,402 (6.4%)	151,029	9,471 (6.3%)	141,558 (93.7%)	
Appropriate chemotherapy	Patients with pN1 or pN2 NSCLC	44,115	3,528 (8.0%)	40,587 (92.0%)	Unknown chemotherapy details	2,214 (5.0%)	41,901	3,388 (8.1%)	38,513 (91.9%)	
Appropriate surgery	Patients with cN2 NSCLC	107,109	11,945 (11.2%)	95,164 (88.8%)	Unknown surgery	473 (0.4%)	106,636	11,898 (11.2%)	94,738 (88.8%)	
<b>Ovary</b>										
Appropriate surgery	Patients with stage I-IIIc ovarian cancer	71,529	7,916 (11.1%)	63,613 (88.9%)	Unknown surgery	91 (0.1%)	71,438	7,906 (11.1%)	63,532 (88.9%)	
<b>Rectum</b>										
Appropriate chemotherapy and radiation	Patients with cT3-4N0-3M0, or cT1-4N1-3M0 rectal cancer treated with surgery; or patients with cT1-2N0M0 who did not receive preoperative therapy though with pT3, pT4, or pN1-3 rectal cancer	55,975	7,066 (12.6%)	48,909 (87.4%)	Unknown chemotherapy; or unknown radiation	2 (0.0%)	55,973	7,066 (12.6%)	48,907 (87.4%)	

---