

Appendix A

Table A1. Hazard Ratios (HR) with 95% Confidence Interval (CI) for Outcomes of Incidence Fracture According to race group, adjusted for T-score diagnosis, age, BMI, previous fracture and PGS groups: results from cox proportional hazard model, individuals with previous fractures were excluded)

	Major osteoporotic fracture	Any fracture
	HR (95 % CI)	HR (95 % CI)
Adjusted for T-score diagnosis		
low	1(reference)	1(reference)
medium	0.83 (0.65-1.06)	0.83 (0.68-1.00)
high	0.83 (0.68-1.02)	0.81 (0.65-1.02)

Significant results are in boldface.

Table A2. Hazard Ratios (HR) with 95% Confidence Interval (CI) for Outcomes of Incidence Fracture According to race group, adjusted for T-score diagnosis, age, BMI, previous fracture and PGS groups: results from cox proportional hazard model, individuals with previous fractures were excluded)

	Major osteoporotic fracture	Any fracture
	OR (95 % CI)	OR (95 % CI)
Adjusted for T-score diagnosis		
Caucasian	1(reference)	1(reference)
American Indian	0.77 (0.49-1.21)	0.71 (0.46-1.11)
African American	0.57 (0.47-0.70)	0.58 (0.48-0.70)
Hispanic	0.45 (0.35-0.58)	0.45(0.35-0.57)
Adjusted for T-score diagnosis + weighted PGS		
Caucasian	1(reference)	1(reference)
American Indian	0.77 (0.48-1.23)	0.71 (0.45-1.12)
African American	0.57 (0.47-0.70)	0.58 (0.48-0.70)
Hispanic	0.45 (0.35-0.59)	0.46 (0.36-0.58)

Significant results are in boldface.