

Appendix 1

IPE Study Simulation: VF

Clinical setting and history

- S:** Nurse on acute admissions unit, calling about the patient in side room 6.
- B:** 60-year-old patient admitted with chest pain and shortness of breath. Previous history of MI treated by coronary stenting.
- A:** Chest pain has returned and seems to be increasing in severity, he is becoming hypoxic and his level of consciousness is decreasing.
- R:** Started high-flow oxygen. Assessment is requested urgently as he may need further treatment.

Clinical course

- agonal gasping/no signs of circulation
- initial rhythm is VF
- after 3rd shock rhythm should change to sinus tachycardia
- P 100 min⁻¹, BP 100/60 mmHg
- patient breathing and rousable.

INTERVENTIONS – KEY TREATMENT POINTS IN BOLD		EXCELLENT	ACCEPTABLE	BORDERLINE	NOT ACHIEVED
<i>Initial approach</i>					
<i>Cardiac arrest management</i>					
VF	Confirm cardiac arrest / check rhythm				
	Call resuscitation team / help				
	Commence CPR until Defib is attached				
	Check monitor / confirm rhythm				
	1st shock at appropriate energy				
	2 min CPR (continuous chest compression / ventilation)				
VF	Check monitor / confirm rhythm				
	Get IV/IO access				
	Minimise interruptions in CPR				
	2nd shock at appropriate energy				
	2 min CPR (continuous chest compression / ventilation)				
VF	Check monitor / confirm rhythm				
	Minimise interruptions in CPR				
	3rd shock at appropriate energy				
	2 min CPR (continuous chest compression / ventilation)				
	Administer 1mg IV adrenaline				
	Administer 300mg Amioderone				
STach	Check monitor / confirm rhythm				
	Check patient (signs of life / pulse)				
	ABCDE assessment				
	Post-arrest investigations				