

ONLINE APPENDIX

TABLE OF CONTENTS

Table 1 Semi-structured interview schedule for IAPT patients	2
Table 2 Semi-structured interview schedule for IAPT psychological wellbeing practitioners (PWPs)	3
Table 3 Semi-structured interview schedule for stop smoking service advisors	5
Table 4 Examples of text that were coded in more than one way, both inductively and deductively	6
Table 5 IAPT patient characteristics	7
Table 6 IAPT psychological wellbeing practitioner (PWP) characteristics	7
Table 7 Smoking cessation advisor characteristics	7

Table 1 Semi-structured interview schedule for IAPT patients

Gain an understanding of patients' experience of smoking and quitting.

- Do you remember the reasons why you started smoking? What were they?
- Have you ever tried to quit smoking?
- What motivated you to quit?
- What helped you in quitting?
- What made it difficult to quit/stay quit?
- If you've never tried to quit, do you mind me asking the reasons why you've not ever tried?

Perspectives about treatments for smoking cessation.

- Have you ever used prescription or over-the-counter medication, or e-cigs to help you quit smoking?
- Have you ever been offered behavioural support?
- What do you think the benefits or risks of using medications/e-cigs are?

Experience of comorbid smoking and mental health.

- Can you explain the ways that your mental health and smoking are related?
 - Can you describe a time/example?
- Can you explain the ways in which smoking might be good or bad for mental health?
- Explain how the withdrawal cycle can mimic mental health symptoms.
 - How does this relate to your own mental health?

Perspectives about parallel treatments of smoking cessation and depression.

- Do you see any issues or benefits of delivering a smoking cessation intervention as an adjunct to usual depression treatment?
- What would be the best way to help someone quit smoking using medications/e-cigs while they are receiving depression treatment?
- What would be good and less good about helping someone quit smoking while they were receiving depression treatment?
- How do you think smoking cessation treatment may be best introduced into therapy settings?

How to optimise recruitment, retention, treatment adherence?

- What have been the good/less good things about the care you received in IAPT?

Is there anything else that you would like to mention?

Table 2 Semi-structured interview schedule for IAPT psychological wellbeing practitioners (PWPs)

Explore PWP skills

- In what ways do you help patients change unhealthy lifestyle behaviours?
- What types of therapy/techniques do you use in IAPT for treating depression?
- Which interventions do you offer?
- Do you sometime address lifestyle behaviours in your treatment plan? What prompts you to do so and what approach do you take.
- Do you do any other addiction work?

Explore professional identity

- Do you feel that part of your role is to help people change lifestyle behaviours, like smoking?

Explore PWP knowledge

- Can you explain the ways that smoking might be good or bad for mental health?
- How do you think that stopping smoking can affect mental health?
- Does it matter that a person tries to stop smoking when they are depressed?
- Do you ever give advice about using smoking cessation medications? If so what?
- Are there any benefits or risks of smoking cessation medications?

Perspectives about the smoking cessation intervention

- Do you see any issues/benefits of delivering smoking cessation intervention as an adjunct to usual IAPT care?
- How do you feel about your patients trying to quit smoking while their using IAPT services?
- What would motivate you to deliver the smoking intervention?

Integration into IAPT treatment

- How do you feel this intervention would best fit into your practice?
- Do you notice any over-lapping techniques/theories between your practice and the behavioural stop smoking treatment?
- How do you think it's best to integrate the stop smoking treatment?
- When to you think it would be best to start the SS treatment during an IAPT therapy session?
- Do you see any barriers/facilitators to introducing SS treatment in to IAPT treatment?
- Are there any service level barriers/ facilitators to its delivery?

Retention

- What are some of the main reasons patients drop out of /remain in treatment?
- Do you think there are any solutions to this?
- Do you have any suggestions about how to retain participants in the treatment and study?

Therapist training needs

- How could we help you in delivering the intervention?
- What training do you think you need?

Is there anything else that you would like to mention?

Table 3 Semi-structured interview schedule for stop smoking service advisors

Elicit views about co-morbid smoking and mental health

- Do you think that smoking and mental health are related? If so, in what ways?
- Can you tell me about how smoking might be good or bad for mental health?
- What are the current barriers and facilitators to retaining patients in the Stop Smoking Services?
- Thinking about your patients with low mood/depression – can you tell me about the reasons that they drop out/remain in treatment?
- Do you think there are any solutions to this?

Talking to patients about the relationship between their mental health and smoking

- How would you go about reassuring a patient w/low mood about psychological withdrawal symptoms?
- How would you go about reassuring a patient who uses smoking to cope with stress and mood? What would you say?
- How do you target beliefs that smoking is mood enhancer? What would you say?
- What skills do you draw upon when talking to patients about the relationship between smoking and mental health?

Intervention manual behavioural components

- Which of these items are most/least useful? And In people who rely heavily on tobacco for mood enhancement, pleasure, stress?
- How long does each component take you?
- Do you combine any techniques together?
- What feedback could we use instead of CO monitoring?
- How do you use the Heaviness of Smoking Index to inform your practice/prescribing?

Intervention manual pharmacological components

- If someone reports depression/mood disorders, what's the standard medication procedure?
- What's the standard medication procedure for high dependency patients?
- How do you address rumors around varenicline and mental health symptoms?

Trial specific

- Do you see any issues or benefits of delivering a smoking cessation intervention as adjunct to usual depression treatment?
- Do you have any suggestions about how to motivate participants to stay in treatment?

Is there anything else that you would like to mention?

Table 4 Examples of text that were coded in more than one way, both inductively and deductively

Deductive: PWP Knowledge/skills

Inductive: PWPs express their ability provide behavioural support

“If they wanted to change it (smoking)... there has to be a desire to change it, so in the assessment we’re checking three factors, so it’s called COMBI ... so it’s their, um, capability to change, the opportunity they’ve got to change and the motivation they’ve got to change their behaviour, so those three factors, if they’re keen to change it, then we would look at integrating it and look at making those changes. If they’re not in a place where they can change, then we look at kind of trying to motivate them to get to that point. So they have to be ready to change, ‘cause it’s got to be their choice, so they may not be in a place where they’re able to do it, and actually, we may just be encouraging them, supporting them to do the basics like get out of bed, have a shower, that kind of thing, if they’re particularly low. Um, if they are motivated and they’re really keen to make some change and actually we’ve done the COMBI assessment and checked on their capabilities, opportunities and motivation and they’re ready to make that change, we then push them a little bit further.”

- Female PWP, aged 30 years, 4 years’ experience

“I think it would be harder because of their low motivation and I think that’s why exploring the beliefs around it might be helpful and um exploring with them their readiness for change, so in terms of motivational – their motivation, where they are on that... so when somebody is thinking about making a change, they go through different stages. So, off the top of my head, you go through a pre-contemplation, a contemplation and then like a part where you are then ready to change, and if you’re in this part here, this stuff would be really good because you move them from this kind of precontemplation to action.”

- Female PWP, aged 28 years, 4 years’ experience

Deductive: Using smoking as a coping mechanism

Inductive: Smoking as a vicious cycle

“If I have one (cigarette) it’s sort of a bit of a relief, not only just maybe five, ten minutes to myself to think it’s more that my body is getting what it’s craving as well and it just eases off and it’s a calmer feeling anyway but that’s probably me just not recognising the fact that I don’t get the nicotine or something”

- Male IAPT patient, aged 32 years, generalised anxiety disorder

“Well it just makes me feel less anxious. Makes me feel a little bit calmer but then obviously you know you’re craving the next cigarette; you know you’re craving the next hit to keep you feeling calm.”

- Female IAPT patient, aged 44 years, mixed anxiety and depressive disorder

Table 5 IAPT patient characteristics

ID	Age	Primary diagnosis	PHQ-9 score	GAD-7 score	Sex
1	44	Mixed anxiety and depressive disorder	14	14	Female
2	54	Depressive episode	20	13	Female
3	26	Social phobias	15	15	Female
4	60	Mixed anxiety and depressive disorder	17	19	Female
5	32	Generalised anxiety disorder	13	15	Male
6	34	Social anxiety	9	8	Male

Table 6 IAPT psychological wellbeing practitioner (PWP) characteristics

ID	Age	Years working as PWP	Site	Sex
01	27	1	1	Female
02	24	2	1	Female
03	30	4	1	Female
04	28	2	1	Female
05	28	3	1	Male
06	35	4	1	Male
01	32	3	2	Female
02	28	1	2	Female
03	30	1	2	Female
04	28	4	2	Female
05	27	2	2	Female

Table 7 Smoking cessation advisor characteristics

ID	Age	Years working as a smoking cessation advisor	Site	Sex
01	28	2	1	Female
02	35	4	1	Male
03	34	2	1	Female
04	36	1	1	Female
05	33	2	1	Female
06	39	2	1	Male