

1 **Air Pollution Reduction and Mortality Benefit during the COVID-19 Outbreak in China**

2 Kai Chen,¹ Meng Wang,² Conghong Huang,² Patrick L. Kinney,³ Paul T. Anastas^{1,4}

3 ¹ Yale School of Public Health, New Haven, CT, USA.

4 ² University at Buffalo School of Public Health and Health Professions, Buffalo, NY, USA.

5 ³ Boston University School of Public Health, Boston, MA, USA

6 ⁴ Yale School of Forestry and Environmental Studies, New Haven, CT, USA.

7

8 **Supplementary Appendix**

9 **Table of contents**

10 Supplementary material and methods page 2-4

11 Table S1 page 5-6

12 Table S2 page 7

13 Table S3 page 8

14 Supplementary references page 9

15 **Supplementary material and methods**

16 ***Daily confirmed COVID-19 cases in China***

17 Daily number of newly confirmed cases for Wuhan, Hubei Province, and China were collected
18 from the official websites of the China National Health Commission
19 (http://www.nhc.gov.cn/xcs/yqtb/list_gzbd.shtml) and the Health Commission of Hubei Province
20 (<http://wjw.hubei.gov.cn/bmdt/ztl/fkxxgzbdgrfyyq/>). A confirmed case was defined as a
21 suspected case with a positive test result based on Novel Coronavirus Pneumonia Diagnosis and
22 Treatment Plan.¹ We collected data from January 10 to March 15, 2020 (Table S1). On February
23 12, there were abnormally high number of cases in Wuhan (13436) and Hubei Province (14840)
24 due to a change in definition of confirmed cases in Hubei Province, which included clinical
25 diagnosed cases with pneumonia imaging features. This additional inclusion of clinical
26 diagnosed cases was only for Hubei Province and was later abolished on February 19 in an
27 updated version of diagnosis plan.

28 ***Air pollution data and analysis***

29 Daily 24-h average concentrations of NO₂ and PM_{2.5} for 367 major Chinese cities from January
30 1, 2016 to March 14, 2020 were obtained from the <http://beijingair.sinaapp.com>. Averaged
31 concentrations from multiple ground-level air quality monitoring stations within each city were
32 used to represent the daily level of air pollutant for each city. Using a difference-in-difference
33 approach, we quantified the air pollution changes due to the quarantine from February 10 to
34 March 14, 2020. The air pollution changes corresponding to Figure 1B can be found in Table S2.
35 Because of the large decreasing trend in PM_{2.5} concentrations during the corresponding
36 quarantine period (17 to 50 days after the Chinese New Year) vs. the corresponding before
37 quarantine period (5 to 20 days before the Chinese New Year) in Wuhan and Hubei Province in

38 2016-2019, the PM_{2.5} reductions due to the 2020 quarantine generally diminished for Wuhan and
39 Hubei Province. However, for the total of 367 Chinese cities, the PM_{2.5} reductions due to the
40 2020 quarantine remained large (18.9 µg/m³).

41 ***Mean NO₂ tropospheric column density calculation with Sentinel-5P satellite images***

42 We used satellite images from the Tropospheric Monitoring Instrument (TROPOMI) which is
43 onboard Sentinel-5 Precursor (Sentinel-5P) satellite² to derive the mean NO₂ tropospheric
44 column density of during and before quarantine periods. The Sentinel-5P is a satellite launched
45 on 13 October 2017 by the European Space Agency to monitor air pollution including NO₂. The
46 NO₂ satellite image analysis was based on Google Earth Engine (GEE), a cloud-based platform
47 for planetary-scale geospatial analysis³. The GEE team has converted the original Sentinel-5P
48 Level 2 NO₂ data which is binned by time, not by latitude/longitude, to a new Level 3 NO₂ data
49 which keeps a single grid per orbit ([https://developers.google.com/earth-](https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S5P_NRTI_L3_NO2)
50 [engine/datasets/catalog/COPERNICUS_S5P_NRTI_L3_NO2](https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S5P_NRTI_L3_NO2)). The new Level 3 NO₂ data is
51 filtered to remove any values with QA values of less than 50. The spatial resolution of the new
52 Level 3 NO₂ data is 0.01 degree (~1km), and the data has been ingested to the GEE platform for
53 analysis. On the GEE platform, we selected all available NO₂ satellite images of during and
54 before quarantine periods, then we averaged them to obtain the mean NO₂ tropospheric column
55 density maps of during and before quarantine periods.

56 ***Concentration-response function and baseline mortality***

57 We applied the cause-specific linear concentration-response functions for NO₂ and PM_{2.5} from
58 two previous time-series studies in 272 Chinese cities using data from 2013-2015.^{4,5} Nation-
59 average mortality rate for total nonaccidental, hypertensive disease, coronary heart disease,
60 stroke, and chronic obstructive pulmonary disease (COPD) were obtained from the China Health

61 and Family Planning Statistical Yearbook 2018. National-averaged cause-specific mortality rate
62 was then multiplied by the total Chinese population to estimate the baseline cause-specific deaths
63 in China. We then used the average daily number of deaths as the baseline daily mortality.

64 ***Health impact assessment***

65 We used the attributable fraction (AF) method to estimate the daily avoided cause-specific
66 mortality from the air pollution reduction. AF is defined as follows:

$$67 \quad AF = 1 - \exp^{-\beta \times \Delta c}$$

68 Where β is the cause-specific coefficient of the concentration-response function and Δc is the air
69 pollution changes due to the quarantine. We then multiply AF with the daily cause-specific
70 number of deaths and the total number of days during the quarantine period (34 days) to estimate
71 the cause-specific avoided deaths. Results corresponding to Figure 1D can be found in Table S3.

72 **Table S1.** Daily number of confirmed COVID-19 cases in Wuhan, Hubei Province, and China
 73 from January 10 to March 14, 2020.

Date	Wuhan	Hubei	China
1/10/2020	41	41	41
1/11/2020	0	0	0
1/12/2020	0	0	0
1/13/2020	0	0	0
1/14/2020	0	0	0
1/15/2020	0	0	0
1/16/2020	6	6	6
1/17/2020	17	17	17
1/18/2020	59	59	59
1/19/2020	77	77	77
1/20/2020	60	72	75
1/21/2020	105	105	149
1/22/2020	62	69	131
1/23/2020	70	105	259
1/24/2020	77	180	444
1/25/2020	46	323	688
1/26/2020	80	371	769
1/27/2020	892	1291	1771
1/28/2020	315	840	1459
1/29/2020	356	1032	1737
1/30/2020	378	1220	1982
1/31/2020	576	1347	2102
2/1/2020	894	1921	2590
2/2/2020	1033	2103	2829
2/3/2020	1242	2345	3235
2/4/2020	1967	3156	3887
2/5/2020	1766	2987	3694
2/6/2020	1501	2447	3143
2/7/2020	1985	2841	3399
2/8/2020	1379	2147	2656
2/9/2020	1921	2618	3062
2/10/2020	1552	2097	2478
2/11/2020	1104	1638	2015
2/12/2020	13436	14840	15152
2/13/2020	3910	4823	5090
2/14/2020	1923	2420	2641
2/15/2020	1548	1843	2009
2/16/2020	1690	1933	2048
2/17/2020	1600	1807	1886
2/18/2020	1660	1693	1749
2/19/2020	615	349	394

Date	Wuhan	Hubei	China
2/20/2020	319	631	889
2/21/2020	314	366	397
2/22/2020	541	630	648
2/23/2020	348	398	409
2/24/2020	464	499	508
2/25/2020	370	401	406
2/26/2020	383	409	433
2/27/2020	313	318	327
2/28/2020	420	423	427
2/29/2020	565	570	573
3/1/2020	193	196	202
3/2/2020	111	114	125
3/3/2020	114	115	119
3/4/2020	131	134	139
3/5/2020	126	126	143
3/6/2020	74	74	99
3/7/2020	41	41	44
3/8/2020	36	36	40
3/9/2020	17	17	19
3/10/2020	13	13	24
3/11/2020	8	8	15
3/12/2020	5	5	8
3/13/2020	4	4	11
3/14/2020	4	4	20
3/15/2020	4	4	16

74

75

76 **Table S2.** Air pollution changes due to the quarantine in Wuhan, Hubei, and China.

	During vs before in 2020	During vs before in 2016-2019	Difference in difference
<i>NO₂ (μg/m³)</i>			
Wuhan	-21.7	1.1	-22.8
Hubei (Outside Wuhan)	-16.9	-4.7	-12.2
China (Outside Hubei)	-15.6	-2.7	-12.9
China	-15.7	-2.7	-12.9
<i>PM_{2.5} (μg/m³)</i>			
Wuhan	-20.0	-18.6	-1.4
Hubei (Outside Wuhan)	-18.5	-22.2	3.7
China (Outside Hubei)	-26.3	-6.6	-19.7
China	-26.0	-7.1	-18.9

77 Note: During quarantine was defined as the period from 17 to 50 days after the Chinese Lunar
78 New Year, whereas before quarantine was defined as the period from 5 to 20 days before the
79 Chinese Lunar New Year.

80 **Table S3.** Avoided cause-specific deaths (95% confidence interval) due to air pollution
81 reduction due to the quarantine.

	NO ₂	PM _{2.5}
Total	8911 (6950, 10866)	3214 (2340, 4087)
Hypertensive disease	490 (283, 695)	202 (68, 335)
Coronary heart disease	1757 (1177, 2335)	863 (577, 1150)
Stroke	2074 (1390, 2756)	782 (409, 1154)
COPD	1427 (1076, 1776)	503 (305, 700)

82

83

84

85

86 **Supplementary references**

- 87 1. Policy and regulatory documents. National Health Commission of the People's Republic
88 of China,. (Accessed March 18, 2020, at <http://www.nhc.gov.cn/wjw/gfxwj/list.shtml>.)
- 89 2. Veefkind J, Aben I, McMullan K, et al. TROPOMI on the ESA Sentinel-5 Precursor: A
90 GMES mission for global observations of the atmospheric composition for climate, air quality
91 and ozone layer applications. *Remote Sensing of Environment* 2012;120:70-83.
- 92 3. Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. Google Earth
93 Engine: Planetary-scale geospatial analysis for everyone. *Remote sensing of Environment*
94 2017;202:18-27.
- 95 4. Chen R, Yin P, Meng X, et al. Fine particulate air pollution and daily mortality. A
96 nationwide analysis in 272 Chinese cities. *American Journal of Respiratory and Critical Care*
97 *Medicine* 2017;196:73-81.
- 98 5. Chen R, Yin P, Meng X, et al. Associations between ambient nitrogen dioxide and daily
99 cause-specific mortality: evidence from 272 Chinese cities. *Epidemiology* 2018;29:482-9.

100