

Supplementary table 1 Baseline characteristics of PPP and PPN patients

No. (%)	PPP (n=19)	PPN (n=18)	P value
Symptoms			
Fever	16 (84.2)	13 (74.2)	0.38
Cough	17(89.5)	12(66.7)	0.09
Expectoration	7(16.8)	2(11.1)	0.07
Hemoptysis	1 (5.3)	0 (0)	0.32
Dyspnea	3 (15.8)	8 (44.4)	0.06
Weep tears	1 (5.3)	0 (0)	0.32
Pharyngalgia	0 (0)	1 (5.3)	0.30
Diarrhea	3 (15.8)	2 (11.1)	0.68
Comorbidities			
CVD	3 (15.8)	2 (11.1)	0.68
Diabetes	2 (10.5)	0 (0)	0.16
Hepatitis	1 (5.3)	1 (5.6)	1
Other	0 (0)	1 (5.6)	0.30
Treatments			
GC	0 (0)	2 (11.1)	0.14
Antiviral	15 (78.9)	11 (61.1)	0.23
Antibiotics	7 (36.8)	10 (55.6)	0.52
TCM	17 (89.5)	13 (72.2)	0.18
Others			
Smoke	0 (0)	3 (16.7)	0.06
Drink	0 (0)	3 (16.7)	0.06

Data are median (IQR) or n (%). *P* values were obtained from χ^2 tests, Fisher's exact tests, T tests or Mann-Whitney U tests, when appropriate. *P* < 0.05 was considered statistically significant (in bold).

Abbreviations: COVID-19, coronavirus disease 19; PP (persistently positive); PPP, PP patients tested positive again; PPN: PP patients tested negative; CVD: cardiovascular diseases; GC, glucocorticoids; TCM, traditional Chinese medicine.

Supplementary table 2 Laboratory testing results of PPP and PPN patients.

Median (IQR)	Normal range	PPP (n=19)	PPN (n=18)	P Value
Blood cells				
WBC (x10 ⁹ /L)	3.5-9.5	5.0 (4.6-5.6)	4.9 (4.2-6.3)	0.59
Neutrophils (x10 ⁹ /L)	1.8-6.3	2.9 (2.3-3.2)	2.7 (2.3-3.7)	0.54
Lymphocytes (x10 ⁹ /L)	1.1-3.2	1.6 (1.2-1.9)	1.5 (1.4-1.7)	0.84
PLTs (x10 ⁹ /L)	125-350	185 (169-227)	176 (148-212)	0.68
Blood biochemicals				
Hb (g/L)	130-175	138 (123-146)	129 (116-140)	0.31
ALB (g/L)	40-55	43.4 (41.7-44.8)	40.2 (37.3-42.7)	0.06
ALT (U/L)	9-50	26.0 (20.5-35.5)	29.5 (18.5-47.3)	0.76
AST (U/L)	15-40	23.0 (19.0-30.0)	25.5 (14.3-32.5)	0.99
Total Bilirubin (μmol/L)	5-21	12.9 (10.9-16.0)	12.9 (9.5-16.6)	
Inflammatory biomarkers				
CRP (g/L)	0-10	1.8 (0.9-2.4)	1.9 (1.3-3.2)	0.14
SAA (mg/L)	0-10	6.5 (4.5-10.1)	6.2 (5.3-10.2)	0.35
IL-6 (pg/mL)	0-7	2.2 (1.5-2.5)	2.7 (1.8-4.7)	0.06

Data are median (IQR) or n (%). *P* values were obtained from χ^2 tests, Fisher's exact tests, T tests or Mann-Whitney U tests, when appropriate. *P* < 0.05 was considered statistically significant (in bold).

Abbreviations: COVID-19, coronavirus disease 19; PP (persistently positive); PPP, PP patients tested positive again; PPN: PP patients tested negative; NLR: neutrophil-to-lymphocyte ratio; PLTs, platelets; Hb, hemoglobin; ALB, albumin; ALT, alanine aminotransferase; AST, aspartate aminotransferase; CRP, C-reactive protein; SAA, serum amyloid A..