

Supplemental Table 1. Association between ethnicity and hospitalization for COVID-19 in UK Biobank – according to sex

		Men n=155,913					Women n=185,053				
		Cases	OR	95% CI	p-value	% Attenuation	Cases	OR	95% CI	p-value	% Attenuation
Black	Age&sex	18	5.41	(3.34 - 8.76)	<0.001		13	3.33	(1.9 - 5.85)	<0.001	
Black	+SES		3.64	(2.22 - 5.97)	<0.001	-23.4		2.44	(1.37 - 4.34)	0.002	-25.8
Black	+Lifestyle		4.94	(3.03 - 8.06)	<0.001	-5.3		2.97	(1.68 - 5.25)	<0.001	-9.6
Black	+Comorbidities		5.41	(3.33 - 8.79)	<0.001	0.0		2.49	(1.41 - 4.40)	0.002	-24.3
Black	+Biomarkers		4.48	(2.75 - 7.30)	<0.001	-11.2		2.73	(1.54 - 4.84)	0.001	-16.5
Black	All		3.51	(2.11 - 5.81)	<0.001	-25.7		1.93	(1.07 - 3.48)	0.029	-45.5
Asian	Age&sex	10	1.68	(0.89 - 3.15)	0.109		11	2.84	(1.54 - 5.20)	0.001	
Asian	+SES		1.32	(0.69 - 2.51)	0.398	-46.3		2.27	(1.22 - 4.20)	0.009	-21.5
Asian	+Lifestyle		1.51	(0.80 - 2.86)	0.209	-20.7		2.53	(1.36 - 4.71)	0.003	-10.8
Asian	+Comorbidities		1.65	(0.87 - 3.12)	0.124	-3.3		2.49	(1.35 - 4.59)	0.004	-12.6
Asian	+Biomarkers		1.35	(0.71 - 2.55)	0.359	-42.4		2.33	(1.26 - 4.31)	0.007	-18.9
Asian	All		1.16	(0.60 - 2.23)	0.669	-72.1		1.91	(1.01 - 3.62)	0.047	-37.9
Others	Age&sex	10	2.42	(1.28 - 4.55)	0.006		7	1.33	(0.62 - 2.82)	0.462	
Others	+SES		1.95	(1.03 - 3.69)	0.040	-24.2		1.15	(0.54 - 2.45)	0.725	-51.8
Others	+Lifestyle		2.11	(1.12 - 4.00)	0.021	-15.1		1.17	(0.55 - 2.49)	0.687	-44.8
Others	+Comorbidities		2.45	(1.30 - 4.61)	0.006	1.4		1.25	(0.59 - 2.66)	0.562	-21.1
Others	+Biomarkers		2.19	(1.16 - 4.13)	0.016	-11.2		1.25	(0.59 - 2.67)	0.557	-20.0
Others	All		1.82	(0.95 - 3.46)	0.069	-32.4		1.04	(0.48 - 2.23)	0.925	-86.8

Supplemental Table 2. Association between ethnicity and hospitalization for COVID-19 in UK Biobank – maximum sample for each set of covariates

		N total sample	OR	95% CI	Attenuation (%)
Black	Age&sex	428,494	4.07	(3.08 - 5.39)	
Black	+SES	417,511	2.65	(1.94 - 3.61)	-30.6
Black	+Lifestyle	396,353	3.74	(2.76 - 5.08)	-6.0
Black	+Comorbidities	415,705	3.95	(2.97 - 5.26)	-2.2
Black	+Biomarkers	377,117	3.35	(2.40 - 4.69)	-13.9
Black	All	341,593	2.63	(1.79 - 3.86)	-31.2
Asian	Age&sex	428,494	2.68	(1.99 - 3.62)	
Asian	+SES	417,511	2.10	(1.52 - 2.90)	-24.8
Asian	+Lifestyle	396,353	2.25	(1.58 - 3.19)	-18.0
Asian	+Comorbidities	415,705	2.47	(1.79 - 3.42)	-8.4
Asian	+Biomarkers	377,117	1.87	(1.31 - 2.67)	-36.8
Asian	All	341,593	1.42	(0.90 - 2.25)	-64.5
Others	Age&sex	428,494	1.97	(1.35 - 2.88)	
Others	+SES	417,511	1.49	(0.99 - 2.26)	-40.7
Others	+Lifestyle	396,353	1.70	(1.13 - 2.57)	-21.3
Others	+Comorbidities	415,705	1.74	(1.15 - 2.65)	-17.8
Others	+Biomarkers	377,117	1.52	(0.96 - 2.41)	-38.1
Others	All	341,593	1.40	(0.86 - 2.29)	-50.1

Supplemental Table 3. Association between ethnicity and hospitalization for COVID-19 in UK Biobank – based on subgroup with data on verbal numerical reasoning

N cases / N total 206 / 116,990	Age- and sex-adjusted			Multiply-adjusted a			Multiply-adjusted + cognition		
	OR	95% CI	p-value	OR	95% CI	p-value			
Ethnicity (reference=White)									
Black (13/ 2,013)	4.61	(2.6 - 8.17)	<0.001	3.63	(1.96 - 6.74)	<0.001	3.06	(1.64 - 5.73)	<0.001
Asian (7/ 2,738)	1.64	(0.77 - 3.49)	0.204	1.15	(0.51 - 2.59)	0.730	0.94	(0.42 - 2.14)	0.889
Other (7/ 2,311)	2.13	(1 - 4.56)	0.051	1.74	(0.8 - 3.79)	0.163	1.47	(0.67 - 3.24)	0.335
Age (years)	1.02	(1.01 - 1.04)	0.013	1.02	(0.99 - 1.04)	0.182	1.02	(0.99 - 1.04)	0.204
Male (vs female)	1.66	(1.25 - 2.19)	<0.001	0.93	(0.6 - 1.45)	0.758	0.94	(0.6 - 1.46)	0.768
Verbal numerical reasoning							0.90	(0.83 - 0.96)	0.003
Lower education				1.06	(0.78 - 1.44)	0.695	0.93	(0.68 - 1.28)	0.659
Household composition (ref=2 people)									
One person				1.17	(0.8 - 1.7)	0.413	1.16	(0.8 - 1.69)	0.429
3 people				1.18	(0.78 - 1.78)	0.427	1.18	(0.78 - 1.78)	0.432
4 people or more				1.27	(0.84 - 1.93)	0.265	1.27	(0.84 - 1.93)	0.264
Townsend deprivation (ref=least deprived Q1)									
Q2				1.12	(0.71 - 1.76)	0.640	1.11	(0.7 - 1.75)	0.660
Q3				1.03	(0.65 - 1.64)	0.896	1.01	(0.64 - 1.61)	0.957
Q4				0.90	(0.56 - 1.44)	0.646	0.88	(0.55 - 1.41)	0.589
Q5				1.06	(0.67 - 1.68)	0.793	1.01	(0.64 - 1.6)	0.965
Physical activity (ref=meeting guideline)									
Active >10min not reaching guideline				0.96	(0.69 - 1.35)	0.826	1.00	(0.71 - 1.4)	0.987
Inactive				1.46	(1.04 - 2.05)	0.031	1.48	(1.05 - 2.08)	0.025
Alcohol (ref=within guideline)									
Never/very rarely drink				1.62	(1.13 - 2.31)	0.009	1.58	(1.11 - 2.26)	0.012
Dangerous intake above guideline				1.36	(0.94 - 1.96)	0.102	1.39	(0.96 - 2)	0.083
Smoking (ref=never smoker)									
Ex-smoker				1.35	(0.99 - 1.84)	0.057	1.36	(1 - 1.85)	0.053
Current smoker				1.50	(0.95 - 2.35)	0.080	1.47	(0.94 - 2.32)	0.092

Body mass index (kg/m ²)	1.00	(0.97 - 1.03)	0.957	1.00	(0.97 - 1.04)	0.994
Waist-to-hip ratio (+0.1 unit)	1.45	(1.15 - 1.84)	0.002	1.44	(1.13 - 1.82)	0.003
Hypertension	1.13	(0.82 - 1.56)	0.453	1.12	(0.81 - 1.55)	0.483
Cardiovascular disease	0.98	(0.58 - 1.66)	0.939	0.97	(0.57 - 1.64)	0.898
Chronic bronchitis	2.39	(1.1 - 5.2)	0.028	2.34	(1.08 - 5.09)	0.032
Ever seen a psychiatrist	1.32	(0.89 - 1.95)	0.164	1.30	(0.88 - 1.92)	0.186
log-CRP	1.08	(0.87 - 1.36)	0.483	1.08	(0.86 - 1.36)	0.494
log-HbA1c	1.46	(0.65 - 3.28)	0.353	1.46	(0.65 - 3.27)	0.357
Cholesterol (mmol/L)	0.86	(0.75 - 0.98)	0.026	0.86	(0.75 - 0.98)	0.027
HDL-cholesterol (mmol/L)	0.65	(0.39 - 1.09)	0.103	0.64	(0.38 - 1.08)	0.094
Forced expiratory volume (L)	0.94	(0.54 - 1.64)	0.829	0.99	(0.56 - 1.75)	0.979

Supplemental Table 4. Association between ethnicity and hospitalization for COVID-19 in UK Biobank - using multiple imputation for covariates

N cases / N total 900 / 428,494	Age and sex			All predictors			Attenuation (%)
	OR	95% CI	p-value	OR	95% CI	p-value	
Ethnicity (reference=White)							
Black (54 / 7,734)	4.07	(3.08 - 5.39)	<.001	2.61	(1.93 - 3.55)	<.001	-31.6
Asian (46 / 9,260)	2.68	(1.99 - 3.62)	<.001	1.68	(1.20 - 2.34)	0.003	-47.7
Other (28 / 8,304)	1.97	(1.35 - 2.88)	<.001	1.40	(0.93 - 2.09)	0.105	-50.7
Age (years)	1.02	(1.01 - 1.03)	<.001	1.02	(1.01 - 1.03)	0.004	
Male (vs female)	1.52	(1.33 - 1.73)	<.001	1.15	(0.93 - 1.42)	0.194	
Lower education				1.14	(0.96 - 1.34)	0.125	
Household composition (ref=2 people)							
One person				1.10	(0.91 - 1.33)	0.319	
3 people				1.13	(0.92 - 1.39)	0.236	
4 people or more				1.49	(1.22 - 1.82)	<.001	
Townsend deprivation (ref=least deprived Q1)							
Q2				1.12	(0.87 - 1.43)	0.383	
Q3				1.13	(0.88 - 1.44)	0.339	
Q4				1.33	(1.05 - 1.68)	0.017	
Q5				1.69	(1.34 - 2.13)	<.001	
Low income (Q1 vs others)				1.15	(0.93 - 1.42)	0.179	
Manual occupation (vs non-manual)				1.05	(0.88 - 1.26)	0.555	
Physical activity (ref=meeting guideline)							
Active >10min not reaching guideline				0.95	(0.79 - 1.14)	0.547	
Inactive				1.20	(1.02 - 1.42)	0.031	
Alcohol (ref=within guideline)							
Never/very rarely drink				1.27	(1.07 - 1.51)	0.007	
Intake above guideline				1.21	(1.01 - 1.44)	0.037	
Smoking (ref=never smoker)							
Ex-smoker				1.31	(1.13 - 1.52)	<.001	
Current smoker				1.01	(0.80 - 1.28)	0.913	
Body mass index (kg/m ²)				1.02	(1.00 - 1.03)	0.026	
Waist-to-hip ratio (+0.1 unit)				1.21	(1.08 - 1.36)	0.001	
Hypertension				1.02	(0.87 - 1.19)	0.814	
Cardiovascular disease				1.09	(0.85 - 1.39)	0.49	
Chronic bronchitis				1.52	(1.03 - 2.24)	0.034	
Psychological distress (ref= low)							
Moderate				1.20	(1.00 - 1.44)	0.053	
High				1.14	(0.95 - 1.36)	0.163	
Ever seen a psychiatrist				1.25	(1.04 - 1.51)	0.019	
log-CRP				1.10	(0.99 - 1.23)	0.075	
log-HbA1c				1.30	(0.86 - 1.95)	0.213	
Cholesterol (mmol/L)				0.69	(0.53 - 0.89)	0.004	
HDL-cholesterol (mmol/L)				0.94	(0.86 - 1.02)	0.121	
Forced expiratory volume (L)				0.82	(0.65 - 1.04)	0.109	

