



Supplementary Material

The King's College London Coronavirus Health and Experiences of Colleagues at King's Study (KCL CHECK) protocol paper: a platform for study of the effects of coronavirus pandemic on staff and postgraduate students.

Katrina A.S. Davis*, Sharon A.M. Stevelink*, Ammar Al-Chalabi, Gabriella Bergin-Cartwright, Rupa Bhundia, Howard Burdett, Ewan Carr, Grace Lavelle, Daniel Leightley, Candice Middleton, Timothy Nicholson, Lucy O'Neill, Catherine Polling, Aoife Ruane, Alice Wickersham, Matthew Hotopf, Reza Razavi

Abstract

Introduction

We will use an occupational sample to study the impact of COVID-19 on current staff and postgraduate research students at a large UK university. The cohort study will address some of the key questions about COVID-19 for the international community, while also providing feedback to the employer and educator.

Methods and analysis

Participants were recruited via email to their University email address. Administrative records were available to compare the composition of volunteer participants to underlying staff and postgraduate student populations of the University. The study comprises a baseline survey, longitudinal follow-up surveys and a viral antibody study. Baseline information was collected in April 2020 including demographics, working situation, current stresses and worries, mental health and neurological symptoms. Personal experiences of COVID-19, indirect experiences and attitudes towards the pandemic were queried, as well as satisfaction with communication and support at work. Longitudinal surveys will assess changes in COVID-19 exposure and mental health. A viral antibody detection component is being planned and will also be longitudinal in nature.

Ethics and dissemination

Ethical approval has been gained from KCL's Psychiatry, Nursing and Midwifery Research Ethics Committee (HR-19/20-18247). Participants were provided with information and agreed to a series of consent statements before enrolment. Data are kept on secure servers with access to personally identifiable information limited. Researchers may apply to have access to pseudonymised data. Findings will be disseminated internally to the University and participants, and externally through scientific publications.

Supplementary Material: The KCL CHECK team

- Principal Investigators: Matthew Hotopf & Reza Razavi (check@kcl.ac.uk)
- Research Manager: Gabriella Bergin-Cartwright
- Senior Investigator and Data Controller: Sharon A.M. Stevelink
- Investigators: Ammar Al-Chalabi, Rupa Bhundia, Howard Burdett, Ewan Carr, Katrina A.S. Davis, Grace Lavelle, Daniel Leightley, Candice Middleton, Timothy Nicholson, Lucy O'Neill, Catherine Polling, Aoife Ruane, Alice Wickersham
- Contact: check@kcl.ac.uk



Supplementary Table (ST1): Items in KCL CHECK

Measure	Purpose and Details	Origin
Measures at baseline only		
Gender	Categorise into Male / Female / Other	NA
Age	Categorise into age groups, and as risk-factor for COVID-19	NA
Ethnicity	Categorise in as high granularity as numbers allow. 18+1 categories collected May have to condense to White British+5+1.	ONS
Postcode	Derive Index of Multiple Deprivations, an area-level socioeconomic indicator. Only first part of postcode will be used to protect identifiable information	NA
House ownership	An individual-level socioeconomic indicator	SELCoH ¹
Relationship status	Social variable	SELCoH ¹
Children and carer status	Number, number living with, age of youngest	Novel
Previous remote working	Occupational variable	Novel
Disability	(i) Any workplace adjustments for occupational health variable (ii) Any condition lasting over 12 months as health variable	(i) Novel (ii) ONS
Physical health history (self-report diagnosis)	Identify conditions that place in high-risk group for COVID-19, and whose health is vulnerable to other factors: heart, immune, asthma, kidney, liver, movement disorder, diabetes, HIV, morbid obesity and cancer	Novel, based on self-administered comorbidity questionnaire (SACQ) ²
Mental health history (self-report diagnosis)	Identify people whose mental health is vulnerable: lifetime history of depression, bipolar, psychotic illness, GAD, agoraphobia, OCD, PTSD, personality disorder, eating disorder, addiction	Health Survey England 2014
Mental health history (self-report diagnosis)	Identify people with lifetime history of suicidal ideation, suicide attempts and non-suicidal self-harm.	Wording designed by the Adult Psychiatric Morbidity Survey (APMS) team ³
Staff questions	Characterise the participants: role, full-time/part-time, contract type, pay spine	Novel
Student questions	Characterise the participants: course, year of study, full-time/part-time	Novel
Measures detailed baseline survey and two monthly follow-up surveys		
Social / Occupational		
Key worker status, individual and household	Identify people in frontline roles	COVID-19 Social Study ⁴
Return to work (retirement or other)	Identify people who have returned to former roles to help the COVID-19 response	Novel
Travelling to work	Measure and monitor	Novel
Remote working and challenges	Identify people who have moved to remote working, broadly characterise problems they have encountered, measure and monitor the impact	Novel
Whether social distancing / isolating	Measure and monitor	Novel, definitions from Age UK



Why social distancing / isolating	Measure and monitor	COVID-19 Social Study ⁴
King's College London specific		
Staff questions: Changes required and how communicated	Measure and monitor the satisfaction with KCL over changes required for staff to do their job in the current situation	Novel
Student questions: Communication regarding studies and funding	Measure and monitor the unmet need for communication with students	Novel
Communication at institutional, departmental and line manager level	Measure satisfaction and identify areas for improvement	Novel
Experiences of COVID-19		
Suspect self to have	Identify	Symptom app ⁵
Have been tested	Identify	Symptom app ⁵
Symptoms	A more objective measure of whether likely to have had, and if so, of what severity: fever, cough, fatigue, headache, shortness of breath, sore throat, loss of smell, chest pain, abdominal pain, diarrhoea, confusion, skipping meals	Symptom app ⁵
Further symptoms	These are less conventional symptoms of COVID-19, but common neurological symptoms that may result from an insult to the brain: problems swallowing, nocturia, hallucination, dizziness, double vision, delusions	Selected from Non-Motor Symptoms Questionnaire (NMSQ) for Parkinson's UK ⁶
Hospital and treatment	Measure of severity	Symptom app ⁵
Contact with COVID-19	Identify those with close contact, those caring for people with COVID at work and outside of work	RADAR-CNS ⁷
Other	Know someone who: has symptoms of COVID-19, has been diagnosed with COVID-19 but is fine, has been diagnosed with COVID-19 and is ill, has died with COVID-19	Novel
Indirect impacts of COVID		
Childcare changes	Measure and monitor	Novel
Accommodation changes	Measure and monitor	Novel
Accessing healthcare	Measure and monitor	Novel
Stressful experiences	Identify people for whom one or more of the following had been experienced: Lost job, Another bill-payer lost their job or was unable to earn money, Unable to pay bills, Had difficulties accessing sufficient food, Started using a food bank, Evicted / lost accommodation, Had difficulties accessing required medication, Somebody close to you in hospital, You lost somebody close to you, Difficulties with family or other social relationships, Difficulties with internet access, Difficulties with your pet, Boredom, Having to change or delay major life plans or events	COVID-19 Social Study ⁴



Worry	Measure and monitor worry around: employment, education, finance, accuracy of info, separation from family	Adapted from RAMP ^{8,9}
Positive changes	Identify and describe these	CoRonavirus Health Impact Survey ¹⁰
Attitude to COVID		
Re COVID	Likert scale measuring: Knowledge, threat (self, others, preoccupied), community response	Novel
Adherence to guidance	Likert scale	COVID-19 Social Study ⁴
Confidence	Likert scale of confidence in government, health services, provision of essentials	COVID-19 Social Study ⁴
Psychiatric		
Generalised Anxiety Disorder (GAD-7)	To measure anxiety symptoms in last 2 weeks, plus change since COVID-19	Spitzer et al ¹¹
Patient Health Questionnaire (PHQ-9)	To measure depression symptoms in last 2 weeks, plus change since COVID-19	Kroenke et al ¹²
3-item Suicidal ideation/self-harm	To identify recent instances of suicidal ideation, suicide attempts and non-suicidal self harm	adapted from Adult Psychiatric Morbidity Survey (APMS) ¹³
Alcohol Use Disorder Identification Test (AUDIT)	To detect recent hazardous / harmful alcohol use, plus change since COVID-19	Babor et al ¹⁴ (slightly adapted)
Psychological		
Pandemic PTSD Check List PPCL	To explore potentially maladaptive thought patterns around the pandemic	RAMP ^{8,9} , adapted from Lang et al ¹⁵
Brief Resilience Scale (BRS)	Measure resilience, 6 item	Smith et al ¹⁶
Locus of Control (novel)	Measure and monitor, 4 item	Novel
Wellbeing	2-items from World Health Organisation Quality of Life. 1-item self-reported health from the SF-36	WHOQOL ¹⁷ , Ware et al ¹⁸
UCLA Loneliness Scale (4 item)	4-items to measure feelings of isolation and loneliness	Hays et al ¹⁹
Neurological		
Fatigue	3 questions on severity, unusualness and impairment	Novel
Epworth Sleepiness Scale (ESS)	1 screening question and 7 follow-up items	Johns et al ²⁰
Limb weakness	1 screening question and 3 follow-up items derived from the Revised Amyotrophic Lateral Sclerosis Functional Rating Scale (ALSFRS-R)	Cedarbaum et al ²¹
Face weakness	1 screening and 3 follow-up items derived from the Facial Cinematic Examination (FaCE) Scale	Kahn et al ²²
Tinnitus	1 screening and 1 follow-up item	UK Biobank ²³
Headache via the Migraine Disability Assessment Scale (MIDAS)	1 screening and 5 follow-up items	Stewart et al ²⁴



Smell and Taste	1 screening and 1 follow-up item (novel)	Novel
-----------------	--	-------

Footnotes – NA = Not applicable; Novel = devised by the research team; ONS = office of national statistics recommended wording / categories; RADAR-CNS = Remote Assessment of Disease and Relapse – Central Nervous System; SELCoH = taken from the South East London Community Health study. Where we have used questions from other studies who have not yet published, references are to the study website.

1. National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust (SLaM), Institute of Psychiatry Psychology & Neuroscience (IoPPN), The Guy's and St Thomas' Charity, et al. South East London Community Health Study (SELCoH) 2020 [Available from: <https://www.kcl.ac.uk/ioppn/depts/pm/research/selcoh>].
2. Sangha O, Stucki G, Liang MH, et al. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Care & Research: Official Journal of the American College of Rheumatology* 2003;49(2):156-63.
3. McManus S, Bebbington P, Jenkins R, et al. Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014. Leeds: NHS Digital, 2016.
4. MARCH Network. COVID-19 Social Study University College London2020 [Available from: <https://www.marchnetwork.org/research>].
5. ZOE Global Limited, King's College London, Spector T. COVID Symptom Tracker (covid.joinzoe.com) 2020 [
6. Romenets SR, Wolfson C, Galatas C, et al. Validation of the non-motor symptoms questionnaire (NMS-Quest). *Parkinsonism & related disorders* 2012;18(1):54-58.
7. King's College London. RADAR-CNS: Remote Assessment of Disease and Relapse - Central Nervous System 2020 [Available from: <https://www.radar-cns.org/>].
8. King's College London, MQ Transforming Mental Health, South London and Maudsley NHS Foundation Trust, et al. RAMP: Repeated Assessment of Mental health in Pandemics 2020 [Available from: <https://rampstudy.co.uk/>].
9. Katherine Seaton Young, Kirstin L. Purves, Shannon Bristow, et al. Repeated Assessment of Mental health in Pandemics @OSF. Open Science Repository. <https://osf.io/7p2ek/>, 2020.
10. Kathleen Merikangas, Argyris Stringaris, Michael P. Milham, et al. V0.3 Adult Self-Report. The CoRonaviruS Health Impact Survey (CRISIS). https://www.nlm.nih.gov/dr2/CRISIS_Adult_Self-Report_Follow_Up_Current_Form_V0.3.pdf: National Institute of Mental Health Intramural Research Program Mood Spectrum Collaboration, Child Mind Institute, Kline Institute for Psychiatric Research,, 2020.
11. Spitzer RL, Kroenke K, Williams JB, et al. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of internal medicine* 2006;166(10):1092-97.
12. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *Journal of general internal medicine* 2001;16(9):606-13.
13. McManus S, Bebbington P, Jenkins R, et al. Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014. A survey carried out for NHS Digital by NatCen Social Research and the Department of Health Sciences, University of Leicester. 2016
14. Babor TF, de la Fuente JR, Saunders J, et al. The Alcohol Use Disorders Identification Test: Guidelines for use in. *Primary care* 2001
15. Lang AJ, Stein MB. An abbreviated PTSD checklist for use as a screening instrument in primary care. *Behaviour research and therapy* 2005;43(5):585-94.
16. Smith BW, Dalen J, Wiggins K, et al. The brief resilience scale: assessing the ability to bounce back. *International journal of behavioral medicine* 2008;15(3):194-200.
17. WHOQOL Group. Assessment Instrument (WHOQOL). *Quality of life Research* 1993;2:153-59.
18. Ware J, Kosinski M, Keller S. SF-36 physical and mental health summary scales. *a user's manual* 2001:1994.
19. Hays RD, DiMatteo MR. A short-form measure of loneliness. *Journal of personality assessment* 1987;51(1):69-81.
20. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *sleep* 1991;14(6):540-45.



21. Cedarbaum JM, Stambler N, Malta E, et al. The ALSFRS-R: a revised ALS functional rating scale that incorporates assessments of respiratory function. *Journal of the neurological sciences* 1999;169(1-2):13-21.
22. Kahn JB, Gliklich RE, Boyev KP, et al. Validation of a patient-graded instrument for facial nerve paralysis: the FaCE scale. *The Laryngoscope* 2001;111(3):387-98.
23. McCormack A, Edmondson-Jones M, Fortnum H, et al. The prevalence of tinnitus and the relationship with neuroticism in a middle-aged UK population. *Journal of Psychosomatic Research* 2014;76(1):56-60.
24. Stewart WF, Lipton RB, Dowson AJ, et al. Development and testing of the Migraine Disability Assessment (MIDAS) Questionnaire to assess headache-related disability. *Neurology* 2001;56(suppl 1):S20-S28.