

SUPPLEMENTARY MATERIALS

Modelling the impact of lockdown easing measures on cumulative COVID-19 cases and deaths in England

Ziauddeen H, PhD ^{1,2,3}, Subramaniam N, BSc ^{1,2}, Gurdasani D, PhD† ⁴.

Supplementary Table 1: Parameters estimated by Bayesian model

Variable	Parameter	No.	Priors
c_t , where $t=1\dots6$	Number of initial cases on first six days	6	exponential(1.0/tau)
R_0	Baseline Reproduction Number	1	Normal(2.4,0.5)
R_t	Time varying effective reproduction number	9	Normal(0.8,0.25)
ϕ	variance parameter for negative binomial distribution of deaths	1	normal(0,5)
τ	parameter in prior of c_t	1	exponential(0.03)

Supplementary Table 2: Comparison of Bayesian models with different constraints on changes in R_t

Model	RMSE	EPLD	SE	diff EPLD model3	diff SE model3
model 1	39.8	-446.7	24.8	-1.3	0.5
model 2	29.9	-446.7	24.3	-1.3	0.9
model 3	29.5	-445.4	24.6	NA	NA

Supp. Table 2 represents model comparisons between Models 1-3, as specified in the text. RMSE represents the Root mean squared error between estimated and observed deaths for each model. EPLD represents the expected log pointwise predictive density which approximates leave-one-out (LOO) cross-validation. Less negative scores suggest better fit. SE is the standard error of EPLD. We assess the difference in EPLD between all models and the best performing model (Model 3 in this case), comparing the difference in EPLD (diff_EPLD_model3) with the standard error of the difference (diff_SE_model3). Although all three models appear comparable in performance, Model 3 appears to show the best fit with the lowest RMSE, and the least negative EPLD.

Supplementary Table 3: Comparison of models excluding specific change points for R_t

Change points removed	RMSE	EPLD	SE	diff EPLD	diff SE
23rd March	49.4	-445.4	24.6	-1.3	0.5
16th March	123.6	-508.0	36.2	-61.3	5.1
13th May	40.5	-447.0	24.8	-0.3	0.1
None (all included)	25.9	-446.7	24.8	-	-

Supp. Table 3 represents model comparisons between Models that constrain R_t at each of the 3 hypothesised change points at which point social distancing or lockdown measures were introduced (16th March and 23rd March), or when lockdown measures were eased (13th May). The first column represents the change point left out in each model, with the last model with all three change points being the comparator, as specified in the text. RMSE represents the Root mean squared error between estimated and observed deaths for each model. EPLD represents the expected log pointwise predictive density which approximates leave-one-out (LOO) cross-validation. Less negative scores suggest better fit. SE is the standard error of EPLD. We assess the difference in EPLD between all models and the model with all three change points, comparing the difference in EPLD (diff_EPLD) with the standard error of the difference (diff_SE). The model leaving out 16th March as a change point, i.e. constraining R_t to remain constant at this point appears to adversely impact fit the most. The model with three change points predictably shows the best fit, although the change point on the 13th May appears to be least important to goodness of fit.

Supplementary Table 4: Cumulative cases and deaths in lockdown easing scenarios in primary model

Rt 1st June	Rt 15th June	Rt 1st July	Cumulative cases	Cumulative deaths	Excess deaths	Excess cases
0.81	0.81	0.81	5632361(5173510-6915570)	49458(45501-60367)	0	0
0.6	0.6	0.6	5509476(5140350-6302650)	48481(45234-55452)	-1026(-5096,-87)	-127932(-635317,-10887)
0.65	0.65	0.65	5525601(5144795-6381955)	48617(45270-56123)	-878(-4360,-74)	-110621(-549336,-9414)
0.7	0.7	0.7	5550714(5153275-6491537)	48827(45341-57050)	-686(-3406,-58)	-87720(-435596,-7465)
0.75	0.75	0.75	5579324(5159964,6646394)	49058(45403-49058)	-431(-2140,-36)	-56286(-279485,-4790)
0.8	0.8	0.8	5620480(5170718-6861630)	49393(45484-59965)	-84(-417,-7)	-11301(-56109,-961)
0.85	0.85	0.85	5689416(5184792-7183460)	49866(45589-62258)	400(34,1988)	56019(4768,278083)
0.85	0.85	0.9	5711579(5188792-7308127)	49978(45607-62846)	518(44,2574)	82022(6979,407641)
0.85	0.9	0.9	5743414(5196366-7476020)	50196(45647-50196)	748(63,3722)	115426(9815,574095)
0.85	0.9	0.95	5787395(5201797-7692210)	50357(45676-64793)	924(78,4597)	159073(13525,791551)
0.9	0.9	0.9	5790145(5202141-7701506)	50519(45726-65640)	1094(93,5427)	161374(13737,800867)
0.9	0.9	0.95	5835939(5215932-7948661)	50721(45752-66642)	1298(110,6441)	211937(18038,1052576)
0.9	0.95	0.95	5890916(5231290-8222697)	51059(45837-68393)	1642(139,8157)	266335(22657,1323354)
0.9	0.95	1	5977038(5251087-8651501)	51339(45898-69842)	1946(165,9667)	351460(29894,1747026)
0.95	0.95	0.95	5960687(5247436-8556522)	51482(45914-70654)	2108(179,10453)	333209(28370,1653092)
0.95	0.95	1	6054039(5266297-9050019)	51852(46040-72397)	2458(209,12192)	431313(36717,2140954)
0.95	1	1	6144873(5278363-9505422)	52397(46165-75013)	2977(253,14775)	521833(44410,2591144)
0.95	1	1.05	6305312(5294301-10332764)	52883(46250-77649)	3496(297,17356)	687119(58467,3412953)
1	1	1	6244213(5286410-10011897)	53022(46265-78249)	3619(308,17937)	623524(53104,3092007)
1	1	1.05	6424833(5313936-10950161)	53600(46385-81203)	4215(359,20895)	813129(69240,4033784)
1	1.05	1.05	6576506(5331499-11709305)	54370(46473-85159)	5002(426,24807)	966759(82304,4796947)
1	1.05	1.1	6886939(5369059-13285488)	55235(46541-89574)	5883(501,29183)	1285783(109452,6381636)
1.05	1.05	1.05	6735809(5355535-12497036)	55257(46546-89643)	5906(503,29254)	1127709(96082,5589085)
1.05	1.05	1.1	7094008(5392553-14295062)	56209(46724-94609)	6912(589,34247)	1492080(127108,7397005)
1.05	1.1	1.1	7360787(5413045-15602425)	57402(46866-100532)	8112(691,40201)	1757003(149653,8711591)
1.05	1.1	1.15	7978843(5468689-18619481)	58845(47082-107855)	9596(817,47563)	2368417(201711,11745319)
1.1	1.1	1.1	7627033(5440955-16891454)	58640(47050-106873)	9410(802,46581)	2020669(172250,10008055)
1.1	1.1	1.15	8329204(5503120-20324645)	60320(47209-115191)	11097(946,54942)	2716047(231501,2716047)
1.1	1.15	1.15	8793270(5537174-22616449)	62138(47434-124243)	12933(1102,64044)	3178393(270877,15746561)
1.1	1.15	1.2	9940071(5632587-28376149)	64625(47624-136460)	15412(1313,76329)	4340411(369882,215062610)


Supplementary Table 4 represents the estimated cumulative deaths, cumulative cases, and excess deaths and cases in different scenarios of changing Rt at points of easing lockdown in comparison with the baseline scenario of Rt remaining constant at 0.81.

Supplementary Table 5: Cumulative cases and deaths in lockdown easing scenarios in model with long serial interval

Rt 1st June	Rt 15th June	Rt 1st July	Cumulative cases	Cumulative deaths	Excess deaths	Excess cases
0.81	0.81	0.81	5770514(5211348-6995778)	50512(45795-60673)	0(0,0)	0(0,0)
0.6	0.6	0.6	5565836(5147626-6306260)	48962(45295-55433)	-1530(-5353,-262)	-202140(-707944,-34538)
0.65	0.65	0.65	5595060(5158988-6404095)	49199(45388-56244)	-1292(-4519,-221)	-173129(-606278,-29587)
0.7	0.7	0.7	5634043(5171297-6532308)	49507(45495-57272)	-992(-3470,-170)	-135443(-474240,-23153)
0.75	0.75	0.75	5683427(5187987-6706775)	49882(45615-58553)	-610(-2133,-105)	-85248(-298431,-14578)
0.8	0.8	0.8	5755680(5207104-6941334)	50403(45769-60249)	-116(-404,-20)	-16655(-58290,-2849)
0.85	0.85	0.85	5847345(5234625-7274127)	51026(45947-62450)	532(91,1860)	79473(13603,278057)
0.85	0.85	0.9	5886578(5246680-7415032)	51201(45996-63081)	709(121,2481)	120159(20473,421431)
0.85	0.9	0.9	5930630(5260107-7571204)	51491(46076-64128)	999(170,3506)	163569(27798,574412)
0.85	0.9	0.95	5993581(5273667-7798411)	51727(46140-64953)	1237(210,4344)	223704(37943,786326)
0.9	0.9	0.9	5986872(5271574-7770844)	51874(46198-65466)	1393(239,4863)	217420(37235,760383)
0.9	0.9	0.95	6052601(5290635-8011751)	52150(46271-66414)	1658(284,5796)	284409(48600,996016)
0.9	0.95	0.95	6116912(5306660-8226648)	52555(46349-67901)	2052(351,7182)	347318(59258,1217297)
0.9	0.95	1	6219910(5332062-8589480)	52918(46475-69247)	2409(412,8439)	446404(76045,1565830)
0.95	0.95	0.95	6191412(5327786-8495735)	53054(46502-69733)	2548(437,8892)	419621(71913,1466787)
0.95	0.95	1	6298436(5354125-8877413)	53438(46613-71062)	2944(505,10285)	529664(90659,1853182)
0.95	1	1	6386181(5371562-9195832)	53988(46728-72946)	3480(596,12167)	621934(106359,2177174)
0.95	1	1.05	6543790(5408114-9776279)	54538(46899-74962)	4014(687,14042)	784701(134030,2748698)
1	1	1	6483640(5391724-9545832)	54642(46919-75263)	4113(707,14346)	721424(123736,2520171)
1	1	1.05	6660710(5439589-10176991)	55215(47060-77315)	4704(808,16420)	901676(154510,3152041)
1	1.05	1.05	6799712(5467759-10630673)	55908(47228-79807)	5436(933,18983)	1038508(177872,3631710)
1	1.05	1.1	7065572(5508833-11578976)	56692(47341-82629)	6229(1068,21766)	1304932(223358,4565657)
1.05	1.05	1.05	6941664(5487865-11119455)	56714(47350-82674)	6253(1075,21795)	1178571(202337,4114119)
1.05	1.05	1.1	7230966(5541652-12163037)	57544(47592-85792)	7130(1225,24865)	1472838(252683,5144062)
1.05	1.1	1.1	7435811(5578879-12856271)	58544(47855-89271)	8131(1397,28369)	1677711(287734,5861114)
1.05	1.1	1.15	7868151(5646112-14377496)	59732(48097-93156)	9305(1598,32481)	2112074(362048,7381365)
1.1	1.1	1.1	7635385(5610959-13548308)	59622(48088-92750)	9199(1584,32039)	1878935(322926,6552933)
1.1	1.1	1.15	8114551(5697587-15225799)	60948(48285-97278)	10494(1806,36562)	2357510(404967,8225599)
1.1	1.15	1.15	8418330(5746149-16305138)	62281(48509-102158)	11868(2041,41365)	2666716(457973,9306311)
1.1	1.15	1.2	9109676(5863863-18746630)	63984(48843-108216)	13598(2338,47413)	3371793(578843,11770623)


Supplementary Table 5 represents the estimated cumulative deaths, cumulative cases, and excess deaths and cases in different scenarios of changing Rt at points of easing lockdown in comparison with the baseline scenario of Rt remaining constant at 0.81.

Supplementary Figure 1: Distribution of R-hat for parameters from final model




Supplementary Figure 1 represents the estimated R-hat for parameters of the final model. An R-hat near 1 suggests that between-chain variance for a given parameter is equal to the within-chain variance, suggesting convergence of the model. Mean R-hat for parameters was 0.996, with all values well below 1.05.

Supplementary Figure 2: Pareto shape parameter k distribution for final model




The estimated shape parameter k of the generalized Pareto distribution can be used to assess the reliability of the estimate from approximations of Leave-one-out cross-validation (LOO). The k shape values are all below 0.5, suggesting our estimates are reliable.

Supplementary Figure 3: R_t estimates with uninformative or broad priors




Supplementary Figures 3A. and 3B. represent estimates of R_t when uninformative priors (A.) or broad priors ($SD=1$) are used for estimation. We find that although median estimates, and patterns of changes are similar as for the original model for most time intervals, substantial uncertainty with higher estimates for R_t are observed in the final time interval from the 13th May up to the 29th May.

Supplementary Figure 4: Estimated reproduction number in model with longer serial interval



The figure shows the R_t estimated by a model with a serial interval of mean 6.5 and coefficient of variation of 0.72. While estimates of R_0 are higher in this model, estimates during other time intervals following lockdown are very similar to our primary model.

Supplementary Figure 5: Predicted and observed deaths in model with longer serial interval



Daily deaths predicted by a model specifying longer serial intervals (blue) with 90% credible intervals (grey) show a good fit to the observed deaths from the ONS (red)