

The far side of the COVID-19 epidemic curve:
local re-openings and re-closings based on
globally coordinated triggers may work best

Supplementary Information Appendix

Vadim A. Karatayev^{1,*}, Madhur Anand¹, Chris T. Bauch²

20 June 2020

1: School of Environmental Sciences, University of Guelph, Guelph, Ontario,
Canada

2: Department of Applied Mathematics, University of Waterloo, Waterloo, On-
tario, Canada

*: vkaratay@uoguelph.ca

Table 1: Parameter definitions, baseline values and literature sources

Parameter	Meaning	Baseline Value	Source
$\tau_I^{t_0}$	symptomatic testing probability for $t < t_{n=325}$	0.045/day	[9, 13, 14], calibrated
$\tau_I^{t_f}$	symptomatic testing probability for $t > 60 + t_{n=325}$	0.3/day	[9, 13, 14], calibrated
α	transition probability, $E \rightarrow A$	0.4/day	[12, 16]
σ	transition probability, $A \rightarrow I$	0.4/day	[12, 16]
ρ	transition probability, $I \rightarrow R$	0.67/day	[12, 16]
w	proportion of contacts in schools and workplaces	0.45	[3]
ω	risk perception proportionality constant	4.4×10^4	[9, 13, 14], calibrated
ϵ	physical distancing efficacy	0.62	[9, 13, 14], calibrated
β_0^A	transmission probability, asymptomatic	3.2×10^{-6} /day	[7, 17], one-half β_0^I
β_0^I	transmission probability, symptomatic	6.4×10^{-6} /day	[7, 17], calibrated
ξ	proportion of mass-action incidence	0.27	[2, 6, 8], calibrated
c	transmission probability constant	4.8×10^{-6}	calibrated
s	superspreading parameter	0.2	[10]
π	proportion of individuals asymptomatic	0.2	[11]
η	adherence to isolation	0.8	[5, 15]
r	reduced travel rate if symptomatic	0.19	[4]
m_{jk}	connectivity matrix	see Methods	[1]
γ_l	Threshold to close county	varied	
γ_G	Threshold to close province	varied	
δ_C	Minimum closure duration	30 days	




Figure 1: **Decreasing coordination in testing of symptomatic individuals across counties increases total cases and person days closed under the local strategy.** $\tau_{I,j}$ blue expresses uncorrelated variation among counties according to a uniform distribution with mean $\tau_I = 0.3$.




Figure 2: **Effects of $\pm 20\%$ variation in each parameter around its default value (Table 1) on the importance of coordination and the optimal trigger prevalence for the local strategy.** Note that disease progression parameters can have counter-intuitive effects because they also regulate the number of positive, active cases on which closures and distancing depend.




Figure 3: Comparison of province-wide (A, C) versus county-by-county (B, D) closure strategies as the effect of county population on transmission probability ξ increases from 0.1 (A, B) to 0.5 (C, D). For each ξ value we re-calibrated the model, with $\beta_0^I = 1.37 \times 10^{-5}$, $= 0.67$, $\omega = 1.9 \times 10^4$, $\tau_I^{t_0} = 0.05$, $\tau_I^{t_f} = 0.31$, and $c = 4.5 \times 10^{-6}$ for $\xi = 0.1$ and $\beta_0^I = 2.58 \times 10^{-6}$, $= 0.68$, $\omega = 1.9 \times 10^4$, $\tau_I^{t_0} = 0.02$, $\tau_I^{t_f} = 0.34$, and $c = 4.0 \times 10^{-6}$ for $\xi = 0.5$. Other settings are as in Fig. 3.




Figure 4: Comparison of province-wide (a) versus county-by-county (b) closure strategies under 100% higher travel rates compared to Fig. 3 of the main text. Other settings are as in Fig. 3.




Figure 5: The fit of 20 stochastic model implementations (black lines) to province-wide data on daily reported positive cases (red dots).

References

- [1] Statistics Canada, 2016 census, catalogue no. 98-400-x2016391., Accessed April 7, 2020.
- [2] Janis Antonovics, Yoh Iwasa, and Michael P Hassell. A generalized model of parasitoid, venereal, and vector-based transmission processes. *The American Naturalist*, 145(5):661–675, 1995.
- [3] Bureau of Labor Statistics. American time use survey — 2018 results. <https://www.bls.gov/news.release/pdf/atus.pdf>, 2018. accessed 4 April 2020.
- [4] CDC. Severe outcomes among patients with coronavirus disease 2019 (COVID-19)—United States, February 12–March 16, 2020. 2020.
- [5] Joel Hellewell, Sam Abbott, Amy Gimma, Nikos I Bosse, Christopher I Jarvis, Timothy W Russell, James D Munday, Adam J Kucharski, W John Edmunds, Fiona Sun, et al. Feasibility of controlling covid-19 outbreaks by isolation of cases and contacts. *The Lancet Global Health*, 2020.
- [6] Samuel Heroy. Metropolitan-scale COVID-19 outbreaks: how similar are they? *arXiv preprint arXiv:2004.01248*, 2020.
- [7] Joe Hilton and Matt J Keeling. Estimation of country-level basic reproductive ratios for novel coronavirus (covid-19) using synthetic contact matrices. *medRxiv*, 2020.
- [8] Hao Hu, Karima Nigmatulina, and Philip Eckhoff. The scaling of contact rates with population density for the infectious disease models. *Mathematical biosciences*, 244(2):125–134, 2013.
- [9] Alexander Lachmann. Correcting under-reported covid-19 case numbers. *medRxiv*, 2020.
- [10] James O Lloyd-Smith, Sebastian J Schreiber, P Ekkehard Kopp, and Wayne M Getz. Superspreading and the effect of individual variation on disease emergence. *Nature*, 438(7066):355–359, 2005.
- [11] Kenji Mizumoto, Katsushi Kagaya, Alexander Zarebski, and Gerardo Chowell. Estimating the asymptomatic proportion of coronavirus disease 2019 (covid-19) cases on board the diamond princess cruise ship, Yokohama, Japan, 2020. *Eurosurveillance*, 25(10):2000180, 2020.
- [12] Hiroshi Nishiura, Natalie M Linton, and Andrei R Akhmetzhanov. Serial interval of novel coronavirus (2019-ncov) infections. *medRxiv*, 2020.
- [13] Ontario. Epidemiological summary. covid-19 cases by reported date (figure 1) <https://files.ontario.ca/moh-covid-19-report-en-2020-04-06.pdf>, 2020; accessed March 31, 2020.

- [14] Lydia Saad. Americans step up their social distancing even further. <https://news.gallup.com/opinion/gallup/298310/americans-step-social-distancing-even-further.aspx>, 2020; accessed March 31, 2020.
- [15] FA Soud, MM Cortese, AT Curns, PJ Edelson, RH Bitsko, HT Jordan, AS Huang, JM Villalon-Gomez, and GH Dayan. Isolation compliance among university students during a mumps outbreak, kansas 2006. *Epidemiology & Infection*, 137(1):30–37, 2009.
- [16] Lauren Tindale, Michelle Coombe, Jessica E Stockdale, Emma Garlock, Wing Yin Venus Lau, Manu Saraswat, Yen-Hsiang Brian Lee, Louxin Zhang, Dongxuan Chen, Jacco Wallinga, et al. Transmission interval estimates suggest pre-symptomatic spread of COVID-19. *medRxiv*, 2020.
- [17] Ashleigh R Tuite, David N Fisman, and Amy L Greer. Mathematical modelling of covid-19 transmission and mitigation strategies in the population of ontario, canada. *CMAJ*, 2020.