

Table S3: Key epidemiological parameters used in the SEIR model, with their respective value (when fixed) or the search intervals used for parameter estimations.

Parameter	Description	Search interval	Fixed	Reference
β_0	Pre-intervention transmission rate	[0,2]	-	-
β_1	Post-intervention transmission rate	[0,2]	-	-
β_2	Second post-intervention transmission rate	[0,2]	-	-
t_1	Time of the first transmission rate change	[Day of community transmission, May 15]	-	-
t_2	Time of the second transmission rate change	[April 25, May 22]	-	-
δ	Factor that reduces the asymptomatic infectivity	[0,0.7]	-	[1, 2, 3]
p	Proportion of latent (E) that proceed to symptomatic infective	-	0.2	[1]
κ	Mean exposed period, or incubation time	-	1/4	[1, 4]
γ_a	Mean asymptomatic period	-	1/3.5	[1]
γ_s	Mean symptomatic period	-	1/4	[1]

References

- [1] Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science*. 2020;368(6490):489–493.
- [2] Ferretti L, Wymant C, Kendall M, Zhao L, Nurtay A, Abeler-Dörner L, et al. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science*. 2020;368(6491).
- [3] Ferguson N, Laydon D, Nedjati Gilani G, Imai N, Ainslie K, Baguelin M, et al. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand; 2020.
- [4] Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. High Contagiousness and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2. *Emerging Infectious Diseases*. 2020;26(7).