

# The impact of contact tracing and household bubbles on deconfinement strategies for COVID-19: an individual-based modelling study (SUPPLEMENTARY MATERIAL)

Lander Willem<sup>1</sup>, Steven Abrams<sup>2,3</sup>, Oana Petrof<sup>2</sup>, Pietro Coletti<sup>2</sup>, Elise Kuylen<sup>1,2</sup>, Pieter Libin<sup>2</sup>, Signe Møgelmo<sup>1,2</sup>, James Wambua<sup>2</sup>, Sereina A. Herzog<sup>1,2</sup>, Christel Faes<sup>2</sup>, SIMID COVID19 team<sup>4</sup>, Philippe Beutels<sup>1,5</sup>, Niel Hens<sup>1,2</sup>

<sup>1</sup> Centre for Health Economic Research and Modelling Infectious Diseases, Vaccine & Infectious Disease Institute, University of Antwerp, Antwerp, Belgium. <sup>2</sup> Data Science Institute, Interuniversity Institute of Biostatistics and statistical Bioinformatics, UHasselt, Hasselt, Belgium. <sup>3</sup> Global Health Institute, Department of Epidemiology and Social Medicine, University of Antwerp, Antwerp, Belgium. <sup>4</sup> The members of the SIMID-COVID-19 team are listed at the end of the main text. <sup>5</sup> School of Public Health and Community Medicine, The University of New South Wales, Sydney, Australia

## Contents

<b>S1 Model population</b>	2
<b>S2 Social contact patterns</b>	4
<b>S3 Age-specific proportion symptomatic cases</b>	6
<b>S4 Parameter estimation</b>	7
<b>S5 Age-specific susceptibility</b>	8
<b>S6 Scenario definitions</b>	10
<b>S7 Platform and technical details</b>	11
<b>S8 References (Supplementary Material)</b>	12