

Table 1. References: SARS-CoV-2 and the Role of Orofecal Transmission:

In: Analysis of the Transmission Dynamics of COVID-19: An Open Evidence Review.
<http://www.cebm.net/evidence-synthesis/transmission-dynamics-of-covid-19/>

References: Included Studies (n=59)

- W1. Ahmed W, Angel N, Edson J et al. First confirmed detection of SARS-CoV-2 in untreated wastewater in Australia: A proof of concept for the wastewater surveillance of COVID-19 in the community. *Science of The Total Environment*. 2020;728:138764. <https://doi.org/10.1016/j.scitotenv.2020.138764>
- W2. Amirian ES. Potential fecal transmission of SARS-CoV-2: Current evidence and implications for public health. *Int J Infect Dis*. 2020;95(1878-3511):363-70. <https://doi.org/10.1016/j.ijid.2020.04.057>
- W3. Cahill N, Morris D. Recreational waters - A potential transmission route for SARS-CoV-2 to humans? [published online ahead of print, 2020 Jun 11]. *Sci Total Environ*. 2020;740:140122. <https://doi.org/10.1016/j.scitotenv.2020.140122>
- W4. Cai J, Xu J, Lin D, et al. A Case Series of children with 2019 novel coronavirus infection: clinical and epidemiological features [published online ahead of print, 2020 Feb 28]. *Clin Infect Dis*. 2020;ciaa198. doi:10.1093/cid/ciaa198
- W5. Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. 2020;395(10223):514-523. doi:10.1016/S0140-6736(20)30154-9
- W6. Chavarria-Miró G, Anfruns-Estrada E, Guix S et al. Sentinel surveillance of SARS-CoV-2 in wastewater anticipates the occurrence of COVID-19 cases. [medRxiv. 2020:2020.06.13.20129627](https://doi.org/10.1101/2020.06.13.20129627)
- W7. Chen C, Gao G, Xu Y, et al. SARS-CoV-2-positive sputum and feces after conversion of pharyngeal samples in patients with COVID-19. *Ann Intern Med*. 2020;172(12):832-834. doi:10.7326/M20-0991 1900
- W8. Chen W, Lan Y, Yuan X, et al. Detectable 2019-nCoV viral RNA in blood is a strong indicator for the further clinical severity. *Emerg Microbes Infect*.

2020;9(1):469-473. Published 2020 Feb 26.
[doi:10.1080/22221751.2020.1732837](https://doi.org/10.1080/22221751.2020.1732837)

- W9. Chen Y, Chen L, Deng Q, et al. The presence of SARS-CoV-2 RNA in the feces of COVID-19 patients. *J Med Virol.* 2020;92(7):833-840. [doi:10.1002/jmv.25825](https://doi.org/10.1002/jmv.25825)
- W10. Cheung KS, Hung IFN, Chan PPY, et al. Gastrointestinal manifestations of SARS-CoV-2 infection and virus load in fecal samples from a Hong Kong cohort: systematic review and meta-analysis. *Gastroenterology.* 2020;S0016-5085(20)30448-0. [doi:10.1053/j.gastro.2020.03.065](https://doi.org/10.1053/j.gastro.2020.03.065)
- W11. [COVID-19 Investigation Team](#). Clinical and virologic characteristics of the first 12 patients with coronavirus disease 2019 (COVID-19) in the United States. *Nat Med.* 2020;26(6):861-868. doi:10.1038/s41591-020-0877-5. Also Kujawski SA, Wong KK, Collins JP et al First 12 patients with coronavirus disease 2019 (COVID-19) in the United States. [medRxiv 2020. 10.1101/2020.03.09.20032896](https://medrxiv.org/abs/2020.10.1101/2020.03.09.20032896)
- W12. Ding S, Liang TJ. Is SARS-CoV-2 also an enteric pathogen with potential fecal-oral transmission? A COVID-19 virological and clinical review [published online ahead of print, 2020 Apr 27]. *Gastroenterology.* 2020;S0016-5085(20)30571-0. [doi:10.1053/j.gastro.2020.04.052](https://doi.org/10.1053/j.gastro.2020.04.052)
- W13. Ding Z, Qian H, Xu B et al. Toilets dominate environmental detection of SARS-CoV-2 virus in a hospital.2020. [medRxiv 2020.04.03.20052175](https://medrxiv.org/abs/2020.04.03.20052175); doi: <https://doi.org/10.1101/2020.04.03.20052175>
- W14. Donà D, Minotti C, Costenaro P, Da Dalt L, Giaquinto C. Fecal-Oral Transmission of SARS-CoV-2 In Children: is it Time to Change Our Approach?. *Pediatr Infect Dis J.* 2020;39(7):e133-e134. [doi:10.1097/INF.0000000000002704](https://doi.org/10.1097/INF.0000000000002704)
- W15. Gupta S, Parker J, Smits S, Underwood J, Dolwani S. Persistent viral shedding of SARS-CoV-2 in faeces - a rapid review. *Colorectal Dis.* 2020;22(6):611-620. [doi:10.1111/codi.15138](https://doi.org/10.1111/codi.15138)
- W16. Han C, Duan C, Zhang S, et al. Digestive symptoms in COVID-19 patients with mild disease severity: clinical presentation, stool viral RNA testing, and outcomes. *Am J Gastroenterol.* 2020. [doi:10.14309/ajg.0000000000000664](https://doi.org/10.14309/ajg.0000000000000664)

- W17. Holshue ML, DeBolt C, Lindquist S *et al* First case of 2019 novel coronavirus in the United States. *N Engl J Med* 2019; 382: 929–936. [10.1056/NEJMoa2001191](https://doi.org/10.1056/NEJMoa2001191)
- W18. Jeong HW, Kim SM, Kim HS, Kim YI, Kim JH, Cho JY, Kim SH, Kang H, Kim SG, Park SJ, Kim EH, Choi YK. *Clin Microbiol Infect*. 2020 Jul 22:S1198-743X(20)30427-4. doi: [10.1016/j.cmi.2020.07.020](https://doi.org/10.1016/j.cmi.2020.07.020). Viable SARS-CoV-2 in various specimens from COVID-19 patients.
- W19. Kim J-M, Kim HM, Lee EJ *et al*. *Osong Public Health Res Perspect*. 2020 Jun; 11(3): 112–117. doi: [10.24171/j.phrp.2020.11.3.02](https://doi.org/10.24171/j.phrp.2020.11.3.02)
- W20. Kingsbury J, Lake R. Potential for foodborne transmission of COVID-19: literature review update. New Zealand Food Safety & Science Research Centre. 2020. <https://www.unitedfresh.co.nz/assets/COVID-19/United-Fresh---Potential-for-Foodborne-Transmission-of-Covid-19---Literature-Review-Update-19-May-20.pdf>
- W21. Lamers MM, Beumer J, van der Vaart J, *et al*. SARS-CoV-2 productively infects human gut enterocytes *Science*. [2020;369\(6499\):50-54](https://doi.org/10.1126/science.abc1669). doi:[10.1126/science.abc1669](https://doi.org/10.1126/science.abc1669)
- W22. La Rosa G, Bonadonna L, Lucentini L, Kenmoe S, Suffredini E. Coronavirus in water environments: Occurrence, persistence and concentration methods - A scoping review. *Water Res*. 2020;179:115899. doi:[10.1016/j.watres.2020.115899](https://doi.org/10.1016/j.watres.2020.115899)
- W23. La Rosa G, Iaconelli M, Mancini P, Bonanno Ferraro G, Veneri C, Bonadonna L, *et al*. First detection of SARS-COV-2 in untreated wastewaters in Italy. [medRxiv. 2020:2020.04.25.20079830](https://doi.org/10.1101/2020.04.25.20079830).
- W24. Lescure FX, Bouadma L, Nguyen D *et al* Clinical and virological data of the first cases of COVID-19 in Europe: a case series. *Lancet Infect Dis* 2020; 20: 697–706. [10.1016/S1473-3099\(20\)30200-0](https://doi.org/10.1016/S1473-3099(20)30200-0)
- W25. Ling Y, Xu SB, Lin YX, *et al*. Persistence and clearance of viral RNA in 2019 novel coronavirus disease rehabilitation patients. *Chin Med J (Engl)*. 2020;133(9):1039-1043. doi:[10.1097/CM9.0000000000000774](https://doi.org/10.1097/CM9.0000000000000774)
- W26. Lo IL, Lio CF, Cheong HH *et al* Evaluation of SARS-CoV-2 RNA shedding in clinical specimens and clinical characteristics of 10 patients with COVID-19 in Macau. *Int J Biol Sci* 2020;16: 1698–707.
- W27. Medema G, Heijnen L, Elsinga G, Italiaander R, Brouwer A. Presence of SARS-Coronavirus-2 in sewage. [medRxiv. 2020:2020.03.29.20045880](https://doi.org/10.1101/2020.03.29.20045880).

- W28. Nicastrì E, D'Abramo A, Faggioni G et al Coronavirus disease (COVID-19) in a paucisymptomatic patient: epidemiological and clinical challenge in settings with limited community transmission, [Italy, February 2020. Euro Surveill 2020; 25: 2000230.](#)
- W29. Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, surface environmental, and personal protective equipment contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) from a symptomatic patient. [JAMA. 2020;323\(16\):1610-2. 2020](#)
- W30. Pan Y, Zhang D, Yang P, Poon LL, Wang Q. Viral load of SARS-CoV-2 in clinical samples. [Lancet Infect Dis 2020; 20: 411–2.](#)
- W31. Parasa S, Desai M, Thoguluva Chandrasekar V, et al. Prevalence of gastrointestinal symptoms and fecal viral shedding in patients with coronavirus disease 2019: a systematic review and meta-analysis. *JAMA Netw Open.* 2020;3(6):e2011335. Published 2020 Jun 1. [doi:10.1001/jamanetworkopen.2020.11335](#)
- W32. Peng L, Liu J, Xu W, Luo Q, Deng K, Lin B, Gao Z. 2019 Novel coronavirus can be detected in urine, blood, anal swabs and oropharyngeal swabs samples. Preprint. Posted online February 25, 2020. medRxiv 2020.02.21.20026179. doi:[10.1101/2020.02.21.20026179](#)
- W33. Qian Q, Fan L, Liu W, et al. Direct evidence of active SARS-CoV-2 replication in the intestine. *Clin Infect Dis.* 2020;ciaa925. <https://doi.org/10.1093/cid/ciaa925>
- W34. Santos VS, Gurgel RQ, Cuevas LE, Martins-Filho PR. Prolonged fecal shedding of SARS-CoV-2 in pediatric patients. A quantitative evidence synthesis [published online ahead of print, 2020 May 22]. *J Pediatr Gastroenterol Nutr.* 2020;10.1097/MPG.0000000000002798. [doi:10.1097/MPG.0000000000002798](#)
- W35. Senapati S, Kshatri JS, Prasad P et al. A pilot study to investigate the fecal dissemination of SARS-CoV-2 virus genome in COVID-19 patients in Odisha, India. medRxiv 2020.05.26.20113167; doi: <https://doi.org/10.1101/2020.05.26.20113167>
- W36. Shutler J, Zaraska K, Holding TM, Machnik M, Uppuluri K, Ashton I, et al. Risk of SARS-CoV-2 infection from contaminated water systems. medRxiv. <https://doi.org/10.1101/2020.06.17.20133504>
- W37. Tan LV, Ngoc NM, That BTT et al Duration of viral detection in throat and rectum of a patient with COVID-19. [medRxiv 2020. 10.1101/2020.03.07.20032052](#)

- W38. Tang A, Tong Z-d, Wang H-l et al Detection of novel coronavirus by RT-PCR in stool specimen from asymptomatic child, China. [Emerg Infect Dis J 2020; 26: 1337–1339.](#)
10.3201/eid2606.200301
- W39. Tian Y, Rong L, Nian W, He Y. Review article: gastrointestinal features in COVID-19 and the possibility of faecal transmission. *Aliment Pharmacol Ther.* 2020;51(9):843-851.
[doi:10.1111/apt.15731](#)
- W40. Wang J, Feng H, Zhang S, et al. SARS-CoV-2 RNA detection of hospital isolation wards hygiene monitoring during the Coronavirus Disease 2019 outbreak in a Chinese hospital. [Int J Infect Dis. 2020;94:103-106.](#) 2020
- W41. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA.* 2020;323(18):1843–1844. [doi:10.1001/jama.2020.3786](#)
- W42. Wang XW, Li J, Guo T, et al. Concentration and detection of SARS coronavirus in sewage from Xiao Tang Shan Hospital and the 309th Hospital of the Chinese People's Liberation Army. [Water Sci Technol. 2005;52\(8\):213-221.](#)
- W43. Xiaorong Wang, Yaya Zhou, Nanchuan Jiang, Qiong Zhou, Wan-Li Ma, Persistence of intestinal SARS-CoV-2 infection in patients with COVID-19 leads to re-admission after pneumonia resolved, *International Journal of Infectious Diseases*, Volume 95, 2020, Pages 433-435, <https://doi.org/10.1016/j.ijid.2020.04.063>.
- W44. Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-19. <https://doi.org/10.1038/s41586-020-2196-x>
- W45. Wu Y, Guo C, Tang L, et al. Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol.* 2020. [10.1016/S2468-1253\(20\)30083-2](https://doi.org/10.1016/S2468-1253(20)30083-2)
- W46. Wurtzer S, Marechal V, Mouchel J-M, Maday Y, Teyssou R, Richard E, et al. Evaluation of lockdown impact on SARS-CoV-2 dynamics through viral genome quantification in Paris wastewaters. *MedRxiv.*
<https://doi.org/10.1101/2020.04.12.20062679>
- W47. Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterology.* 2020. [10.1053/j.gastro.2020.02.055](https://doi.org/10.1053/j.gastro.2020.02.055)

- W48. Xiao F, Sun J, Xu Y, et al. Infectious SARS-CoV-2 in Feces of Patient with Severe COVID-19 [published online ahead of print, 2020 May 18]. *Emerg Infect Dis*. 2020;26(8):10.3201/eid2608.200681. [doi:10.3201/eid2608.200681](https://doi.org/10.3201/eid2608.200681)
- W49. Xing Y, Wei N, Qin W et al Prolonged presence of SARS-CoV-2 in feces of pediatric patients during the convalescent phase. [medRxiv 2020. 10.1101/2020.03.11.20033159](https://doi.org/10.1101/2020.03.11.20033159)
- W50. Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med*. 2020;26(4):502-505. [doi:10.1038/s41591-020-0817-4](https://doi.org/10.1038/s41591-020-0817-4)
- W51. Yao H, Xiangyun Lu, Qiong Chen et al. Patient-derived mutations impact pathogenicity of SARS-CoV-2
<https://www.medrxiv.org/content/10.1101/2020.04.14.20060160v2.full.pdf>
- W52. Young BE, Ong SWX, Kalimuddin S, et al; Singapore 2019 Novel Coronavirus Outbreak Research Team. Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore. *JAMA*. 2020;323(15):1488-1494. [doi:10.1001/jama.2020.3204](https://doi.org/10.1001/jama.2020.3204)
- W53. Yuan C, Zhu H, Yang Y, et al. Viral loads in throat and anal swabs in children infected with SARS-CoV-2. *Emerg Microbes Infect*. 2020;9(1):1233-1237. [doi:10.1080/22221751.2020.1771219](https://doi.org/10.1080/22221751.2020.1771219)
- W54. Zang R, Castro MFG, McCune BT et al. The SARS-CoV-2 virus infects cultured ACE2-expressing human enterocytes aided by the TMPRSS2 and TMPRSS4 serine proteases. *Science Immunology* 13 May 2020: Vol. 5, Issue 47, eabc3582 [DOI: 10.1126/sciimmunol.abc3582](https://doi.org/10.1126/sciimmunol.abc3582)
- W55. Zhang J, Wang S, Xue Y. Fecal specimen diagnosis 2019 novel coronavirus-infected pneumonia. *J Med Virol*. 2020. [10.1002/jmv.25742](https://doi.org/10.1002/jmv.25742)
- W56. Zhang T, Cui X, Zhao X, et al. Detectable SARS-CoV-2 viral RNA in feces of three children during recovery period of COVID-19 pneumonia. *J Med Virol*. 2020;92(7):909-914. [doi:10.1002/jmv.25795](https://doi.org/10.1002/jmv.25795)
- W57. Zhang W, Du RH, Li B, et al. Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes. *Emerg Microbes Infect*. 2020;9(1):386-389. <https://doi.org/10.1080/22221751.2020.1729071>

W58. Zhang Y , Cao C, Shuangli Z et al. Isolation of 2019-nCoV from a Stool Specimen of a Laboratory-Confirmed Case of the Coronavirus Disease 2019 (COVID-19)[J]. China CDC Weekly, 2020, 2(8): 123-124. [doi: 10.46234/ccdcw2020.033](https://doi.org/10.46234/ccdcw2020.033)

W59. Zhou J, Li C, Liu X, et al. Infection of bat and human intestinal organoids by SARS-CoV-2 [published online ahead of print, 2020 May 13]. Nat Med. 2020;10.1038/s41591-020-0912-6. [doi:10.1038/s41591-020-0912-6](https://doi.org/10.1038/s41591-020-0912-6)

References: Other

1. Transmission of SARS-CoV-2: implications for infection prevention precautions <https://www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-for-infection-prevention-precautions>
2. Patient-derived mutations impact pathogenicity of SARS-CoV-2 Hangping Yao et al <https://www.medrxiv.org/content/10.1101/2020.04.14.20060160v2.full.pdf>
3. Holmes K.V. Enteric infections with coronaviruses and toroviruses. *Novartis Found Symp.* 2001;238:258–269. discussion 269–275.
4. Ye G, Lin H, Chen S, et al. Environmental contamination of SARS-CoV-2 in healthcare
5. Ong SWX, Tan YK, Chia PY, et al. Air, Surface Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient. *JAMA.* 2020;323(16):1610–1612. doi:10.1001/jama.2020.3227
6. Guo ZD, Wang ZY, Zhang SF, et al. Aerosol and Surface Distribution of Severe Acute
7. Wang J, Feng H, Zhang S, et al. SARS-CoV-2 RNA detection of hospital isolation wards
8. Zhou J et al. Human intestinal tract serves as an alternative infection route for Middle East respiratory syndrome coronavirus. *Sci Adv.* 2017;3:eaa04966. <https://advances.sciencemag.org/content/3/11/eaa04966>
9. Lee SS, Wong NS. Probable transmission chains of Middle East respiratory syndrome coronavirus and the multiple generations of secondary infection in South Korea. *Int J Infect Dis.* 2015 Sep; 38():65-7.
10. [Further studies on human enteric coronaviruses.](#) Caul EO, Egglestone SI.
11. Booth CM et al. Clinical features and short-term outcomes of 144 patients with SARS in the greater Toronto area. *JAMA.* 2003;289:2801–2809.
12. Leung WK, To K-F, Chan PKS, Chan HLY, Wu AKL, Lee N, Yuen KY, Sung JJY. Enteric involvement of severe acute respiratory syndrome-associated coronavirus infection. *Gastroenterology* 125, 1011–1017 (2003).
13. Hung IFN et al. Viral loads in clinical specimens and SARS manifestations. *Emerg. Infect. Dis.* 10, 1550–1557 (2004).
14. Peiris JSM et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet.* 2003;361:1767–1772.
15. WHO environmental health team reports on Amoy gardens. Available at: <http://www.info.gov.hk/gia/general/200305/16/0516114.htm>.
16. Ding Y, He L, Zhang Q, et al. Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus transmission pathways. *J Pathol.* 2004;203(2):622-630. doi:10.1002/path.1560

17. Wang XW, Li JS, Guo TK, et al. Concentration and detection of SARS coronavirus in sewage from Xiao Tang Shan Hospital and the 309th Hospital [published correction appears in *J Virol Methods*. 2005 Dec;130(1-2):210]. *J Virol Methods*. 2005;128(1-2):156-161. doi:10.1016/j.jviromet.2005.03.022
18. Analysis of SARS-CoV-2 transmission clusters and superspreading events Gwen Knight, Quentin Leclerc and Adam Kucharski, on behalf of CMMID working group Report for SPI-M, 3rd June 2020. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/897562/S0473_SPI-M_Superspreading_and_clusters.pdf
19. Hartmann, Katrin (2005). "Feline infectious peritonitis". *Veterinary Clinics of North America: Small Animal Practice*. **35** (1): 39–79. doi:10.1016/j.cvsm.2004.10.011
20. Immune evasion of porcine enteric coronaviruses and viral modulation of antiviral innate signaling.
21. Fan Y, Zhao K, Shi ZL, Zhou P. Bat Coronaviruses in China. *Viruses*. 2019;11(3):210. Published 2019 Mar 2. doi:10.3390/v11030210
22. Killerby ME, Biggs HM, Midgley CM, Gerber SI, Watson JT. Middle East Respiratory Syndrome Coronavirus Transmission. *Emerg Infect Dis*. 2020;26(2):191-198. <https://dx.doi.org/10.3201/eid2602.190697>