

1 Supplementary Table. Sensitivity and specificity of each symptom for identifying  
2 individuals with antibodies against SARS-CoV-2. The EPICOV19 study, third wave.  
3

Symptom	Sensitivity	95% CI	Specificity	95% CI
Headaches	58.0	54.5 - 61.3	64.5	64.0 - 65.1
Change in smell or taste	56.5	53.1 - 59.9	90.9	90.6 - 91.2
Fever	52.1	48.6 - 55.5	87.8	87.4 - 88.1
Cough	47.7	44.3 - 51.1	77.8	77.3 - 78.3
Myalgia	44.1	40.7 - 47.5	84.3	83.9 - 84.7
Sore throat	33.8	30.6 - 37.1	83.4	83.0 - 83.8
Diarrhea	25.6	22.7 - 28.6	88.3	88.0 - 88.7
Difficulty breathing	23.1	20.3 - 26.1	90.6	90.3 - 90.9
Shivering	20.5	17.8 - 23.4	93.9	93.6 - 94.1
Palpitation	20.0	17.4 - 22.9	89.5	89.1 - 89.8
Vomiting	9.5	7.7 - 11.7	96.0	95.8 - 96.2


4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

35 Supplementary Figure 1. Mean number of symptoms and percent asymptomatic in  
 36 subjects negative for antibodies against SARS-CoV-2, according to sociodemographic  
 37 characteristics. The EPICOV19 study, third wave.

38

39 *Notes: diamonds represent the main Y axis for the mean number of symptoms (with*  
 40 *their respective 95% CI). The bars represent the secondary Y axis for the prevalence of*  
 41 *asymptomatic subjects.*


42



44

45

46 Supplementary Figure 2. Scatter diagram between prevalence of asymptomatic  
47 individuals with antibodies and seroprevalence for SARS-CoV-2, among the  
48 municipalities included in the study. The EPICOV19 study, third wave.  
49



50

51