

Supplement 1. True concordance probability under several scenarios.

prevalence	Estimated sensitivity and specificity of nasopharyngeal and saliva test			If the sensitivity of saliva test is 10% less than the sensitivity of nasopharyngeal test			If the sensitivity of saliva test is 20% less than the sensitivity of nasopharyngeal test		
	Point estimates	90%CI		Point estimates	90%CI		Point estimates	90%CI	
0	0.999	0.997	1.000	0.999	0.997	1.000	0.999	0.997	1.000
0.001	0.999	0.997	0.999	0.999	0.997	0.999	0.999	0.997	0.999
0.005	0.997	0.995	0.999	0.998	0.996	0.999	0.997	0.995	0.999
0.01	0.996	0.992	0.998	0.997	0.993	0.998	0.996	0.993	0.998
0.02	0.993	0.987	0.997	0.995	0.988	0.998	0.993	0.987	0.996
0.03	0.990	0.981	0.997	0.993	0.983	0.997	0.990	0.981	0.995
0.04	0.988	0.975	0.996	0.991	0.977	0.996	0.988	0.976	0.994
0.05	0.985	0.969	0.995	0.989	0.972	0.996	0.985	0.970	0.993
0.06	0.982	0.963	0.995	0.987	0.967	0.995	0.982	0.964	0.992
0.07	0.979	0.957	0.994	0.985	0.961	0.995	0.979	0.958	0.991
0.08	0.977	0.951	0.993	0.983	0.956	0.995	0.977	0.952	0.990
0.09	0.974	0.944	0.993	0.981	0.951	0.994	0.974	0.946	0.989
0.1	0.971	0.938	0.992	0.979	0.945	0.994	0.971	0.941	0.988
0.11	0.968	0.932	0.991	0.977	0.940	0.993	0.968	0.935	0.987
0.12	0.966	0.926	0.991	0.975	0.935	0.993	0.966	0.929	0.986
0.13	0.963	0.920	0.990	0.973	0.929	0.992	0.963	0.923	0.985
0.14	0.960	0.914	0.989	0.971	0.924	0.992	0.960	0.917	0.985
0.15	0.957	0.908	0.989	0.969	0.919	0.991	0.957	0.912	0.984
0.16	0.955	0.902	0.988	0.967	0.913	0.991	0.955	0.906	0.983
0.17	0.952	0.896	0.987	0.965	0.908	0.991	0.952	0.900	0.982
0.18	0.949	0.890	0.987	0.963	0.903	0.990	0.949	0.894	0.981
0.19	0.946	0.884	0.986	0.961	0.897	0.990	0.946	0.888	0.980
0.2	0.944	0.878	0.985	0.959	0.892	0.989	0.943	0.883	0.979