

1 **Supplementary Section**

2 **Suppl Table 1.** Presentation of demographic and candidate proinflammatory cytokines in the
 3 study participants.

Measures	Disease Controls (n=16)	Healthy Controls (n=8)	p-value/ comments
Age (yrs.)	41.87±11.04	28.38±4.44	0.002
Sex (F/M), in number	F=4/M=12	F=2/M=6	NA
BMI (units)	27.11±6.14	24.65±3.328	NS
Candidate Cytokines observed in immune storm			
IL-6 (pg/ml)	2.478±1.37	0.406±0.20	≤0.001
IL-10 (pg/ml)	0.335±0.17	0.285±0.12	NS
IL-1β (pg/ml)	0.598±0.30	0.219±0.40	0.017

4 Footnote - yrs.: years; F: Females; M: Males; BMI: Body Mass index; IL: Interleukin; β: beta; pg:
 5 picogram; ml: milliliter; NA: Not Applicable; NS: Not significant.

6
 7 **Suppl Table 2.** Area Under the Curve (AUC), and maximum or peak concentration (C_{max}) derived
 8 for the low and high doses of thiamine (Vit B1).

Vit B1 Dose	79 mg	474 mg
Blood AUC	169.06	483.48
Plasma AUC	139.04	516.66
CL	56%	91%
Blood Cmax	31.6	66.36
Plasma Cmax	30.81	85.32

9 Footnote - Unit for AUC: nmol/Liter x Hours for 10 Hrs.; CL: Clearance; Unit for C_{max}: nmol/Liter;
 10 AUC: Area Under the Curve; C_{max}: Peak Concentration of Drug in Specific Body System/ Organ.
 11 Vit B1: Thiamine; mg: milligram.