

Supplementary materials


**The impact of the COVID-19 pandemic on mental health in the general population – a comparison between Germany and the UK**

**Franziska Knolle<sup>1,2</sup>, Lisa Ronan<sup>2</sup>, Graham K. Murray<sup>2,3</sup>**


<sup>1</sup> Department of Neuroradiology, Klinikum rechts der Isar, TUM, München, Germany

<sup>2</sup> Department of Psychiatry, University of Cambridge, Cambridge UK


<sup>3</sup> Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, United Kingdom




Supp. Figure 1. Distribution of highest personal education by country.




Supp. Figure 2. Percentage of suspected infection by country, presented in varying scales.




Supp. Figure 3. Distribution of symptoms by country, with close-up presented in the left panel.




Supp. Figure 4. Distribution of contacts to people with possible infections by country, with close-up presented in the left panel.



Supp. Figure 5. Impact on working situation by country, with close-up presented in the left panel.



Supp. Figure 6. Each boxplot shows the subjective change of one of the sub-dimensions (experiences and beliefs (AEB), social Anhedonia (SAnh), paranoid ideation (PI), social anxiety (SAnx), eccentricity (Ecc), and disorganised speech (DS)) of the SPQ during the pandemic, separately for Germany (yellow-orange) and the UK (grey).



Supp. Figure 7. Pearson correlation between GSI-27 and total SPQ score, separately per country. GSI-27 and SPQ total reveal a strong positive correlation.