

ONLINE SUPPLEMENT

Sleep Disturbances, Anxiety and Burnout during the COVID-19 Pandemic: a nationwide cross-sectional study in Brazilian Healthcare Professionals.

Luciano F. Drager,^{1,2} Daniela V. Pachito,^{3,4} Claudia R.C. Moreno,^{5,6} Almir R. Tavares Jr,⁷ Silvia G. Conway,^{8,9} Márcia Assis,¹⁰ Danilo A. Sguillar,¹¹ Gustavo A. Moreira,¹² Andrea Bacelar,¹³ Pedro R. Genta¹⁴

AFFILIATIONS

1. Hypertension Unit, Renal Division, University of São Paulo Medical School, Brazil;
2. Hypertension Unit, Heart Institute (InCor), University of São Paulo Medical School, Brazil
3. Núcleo de Avaliação de Tecnologias em Saúde, Hospital Sírio-Libanês, São Paulo, Brazil
4. Fundação Getúlio Vargas, Brazil
5. School of Public Health, University of São Paulo, Brazil
6. Stress Research Institute, Department of Psychology, Stockholm University, Sweden
7. Neurosciences Postgraduate Program, Federal University of Minas Gerais, Brazil
8. Akasa - Formação e Conhecimento, São Paulo, Brazil;
9. Psychiatry Department, University of São Paulo Medical School, Brazil;
10. Clínica do Sono de Curitiba, Hospital São Lucas, Curitiba Paraná, Brazil
11. ENT Department of the Federal University of São Paulo, Brazil
12. Department of Pediatrics and Psychobiology, Federal University of São Paulo, Brazil
13. Carlos Bacelar Clinica, Rio de Janeiro, Brazil
14. Laboratório do Sono, LIM 63, Pulmonary Division, Heart Institute (InCor), Hospital das Clínicas HCFMUSP, Universidade de Sao Paulo, Sao Paulo, SP, BR.

ADDRESS FOR CORRESPONDENCE:

Luciano F. Drager, MD, PhD.
Associate Professor of Medicine
University of São Paulo Medical School
Brazil

Table S1: Logistic regression model for the main outcome (new-onset insomnia or worsening previous insomnia).

	Coefficient	Standard Error	Odds Ratio	95% confidence interval		P
				Lower limit	Upper limit	
Age	0.002	0.003	1.002	0.995	1.008	0.601
Females	0.563	0.085	1.756	1.487	2.075	< 0.001
Physicians	-0.293	0.155	0.746	0.550	1.011	0.058
Nurses	-0.237	0.179	0.789	0.555	1.121	0.185
Nurse technicians	-0.149	0.228	0.862	0.551	1.348	0.513
Physiotherapists	-0.544	0.187	0.580	0.402	0.837	0.004
Dentists	-0.394	0.194	0.674	0.461	0.985	0.042
Psychologists	-0.554	0.217	0.575	0.374	0.878	0.011
Speech therapists	0.048	0.262	1.049	0.627	1.754	0.855
Other healthcare	-0.193	0.199	0.824	0.557	1.218	0.334
Workplace - Ward	0.009	0.086	1.009	0.851	1.194	0.921
Workplace – E.R, intensive care units	0.184	0.097	1.202	0.994	1.454	0.057
Workplace - Office	-0.229	0.086	0.795	0.671	0.942	0.008
Workplace – Sleep Laboratory	-0.249	0.227	0.780	0.495	1.209	0.272
Workplace – Operating room	0.032	0.111	1.033	0.831	1.282	0.770
Workplace - Pharmacy	-0.044	0.314	0.957	0.515	1.774	0.889
Workplace – Administrative area	0.087	0.110	1.091	0.879	1.352	0.428
Previous COVID-19 diagnosis	0.036	0.118	1.037	0.822	1.307	0.759
Recent/current care of patients with COVID-19	0.257	0.081	1.293	1.104	1.514	0.001
Regional COVID-19 death incidence	-0.002	0.015	0.998	0.968	1.028	0.871
Increased work hours	-0.005	0.003	0.995	0.990	1.000	0.038
Weight (decreased)	0.616	0.097	1.852	1.531	2.240	< 0.001
Weight (increased)	0.433	0.078	1.542	1.323	1.799	< 0.001
Family Income (increased)	-0.384	0.182	0.681	0.476	0.970	0.034
Family Income (>0 but <30% reduction)	0.071	0.091	1.074	0.899	1.283	0.433
Family Income (>30% reduction)	0.312	0.092	1.366	1.140	1.636	0.001
Anxiety (GAD-2 - 3 or more)	1.166	0.070	3.209	2.796	3.684	< 0.001
New-onset burnout	0.686	0.086	1.986	1.677	2.352	< 0.001

Table S2: Logistic regression model for sleep quality.

	Coefficient	Standard Error	Odds Ratio	95% confidence interval		P
				Lower limit	Upper limit	
Age	-0.008	0.004	0.992	0.985	1.000	0.037
Females	0.141	0.093	1.151	0.959	1.382	0.130
Physicians	-0.360	0.190	0.698	0.480	1.012	0.059
Nurses	-0.360	0.224	0.698	0.450	1.082	0.108
Nurse technicians	-0.184	0.227	0.832	0.532	1.298	0.418
Physiotherapists	-0.093	0.230	0.911	0.580	1.427	0.684
Dentists	-0.544	0.247	0.580	0.357	0.941	0.028
Psychologists	-0.732	0.288	0.481	0.273	0.846	0.011
Speech therapists	-0.097	0.323	0.908	0.482	1.708	0.763
Other healthcare	-0.181	0.247	0.834	0.514	1.354	0.464
Workplace - Ward	0.008	0.104	1.008	0.822	1.236	0.939
Workplace – E.R, intensive care units	0.225	0.117	1.252	0.995	1.576	0.055
Workplace - Office	-0.043	0.103	0.958	0.784	1.172	0.679
Workplace – Sleep Laboratory	-0.075	0.254	0.928	0.562	1.521	0.769
Workplace – Operating room	-0.091	0.127	0.913	0.711	1.171	0.476
Workplace - Pharmacy	0.275	0.395	1.317	0.611	2.893	0.486
Workplace – Administrative area	-0.112	0.130	0.894	0.693	1.152	0.387
Previous COVID-19 diagnosis	0.035	0.143	1.036	0.783	1.373	0.804
Recent/current care of patients with COVID-19	0.195	0.094	1.215	1.010	1.461	0.038
Regional COVID-19 death incidence	0.044	0.018	1.045	1.010	1.082	0.011
Increased work hours	0.010	0.003	1.010	1.004	1.016	0.002
Weight (decreased)	0.264	0.114	1.302	1.042	1.627	0.020
Weight (increased)	0.298	0.089	1.347	1.131	1.604	0.001
Family Income (increased)	-0.419	0.208	0.658	0.437	0.988	0.044
Family Income (>0 but <30% reduction)	0.090	0.105	1.094	0.890	1.344	0.393
Family Income (>30% reduction)	0.121	0.110	1.129	0.910	1.399	0.271
Anxiety (GAD-2 - 3 or more)	0.919	0.087	2.507	2.115	2.972	< 0.001
New-onset burnout	0.659	0.120	1.933	1.531	2.448	< 0.001
New-onset or worsening Insomnia	2.848	0.108	17.253	14.025	21.399	< 0.001

Table S3: Logistic regression model for sleep quantity (at least one-hour reduction).

	Coefficient	Standard Error	Odds Ratio	95% confidence interval		P
				Lower limit	Upper limit	
Age	0.014	0.003	1.014	1.007	1.021	< 0.001
Females	0.102	0.088	1.107	0.932	1.316	0.248
Physicians	-0.619	0.163	0.538	0.391	0.741	< 0.001
Nurses	-0.486	0.189	0.615	0.424	0.890	0.010
Nurse technicians	-0.436	0.196	0.647	0.439	0.949	0.026
Physiotherapists	-0.280	0.203	0.756	0.507	1.124	0.168
Dentists	-0.740	0.224	0.477	0.307	0.738	0.001
Psychologists	-0.402	0.247	0.669	0.412	1.088	0.104
Speech therapists	-0.325	0.280	0.723	0.416	1.250	0.246
Other healthcare	-0.374	0.212	0.688	0.454	1.041	0.077
Workplace - Ward	0.183	0.091	1.201	1.005	1.436	0.044
Workplace – E.R, intensive care units	0.143	0.102	1.154	0.945	1.409	0.160
Workplace - Office	-0.023	0.092	0.977	0.816	1.171	0.806
Workplace – Sleep Laboratory	-0.246	0.242	0.782	0.482	1.249	0.310
Workplace – Operating room	-0.413	0.121	0.662	0.521	0.837	0.001
Workplace - Pharmacy	0.650	0.342	1.916	0.988	3.790	0.057
Workplace – Administrative area	0.131	0.115	1.140	0.909	1.428	0.256
Previous COVID-19 diagnosis	0.139	0.125	1.149	0.899	1.470	0.267
Recent/current care of patients with COVID-19	0.144	0.085	1.155	0.977	1.366	0.091
Regional COVID-19 death incidence	-0.014	0.016	0.986	0.956	1.018	0.395
Increased work hours	0.020	0.003	1.020	1.014	1.025	< 0.001
Weight (decreased)	0.487	0.102	1.627	1.332	1.990	< 0.001
Weight (increased)	0.384	0.083	1.468	1.249	1.727	< 0.001
Family Income (increased)	0.301	0.185	1.351	0.940	1.943	0.104
Family Income (>0 but <30% reduction)	0.014	0.095	1.014	0.842	1.221	0.886
Family Income (>30% reduction)	-0.073	0.098	0.930	0.768	1.125	0.453
Anxiety (GAD-2 - 3 or more)	0.456	0.077	1.578	1.358	1.834	< 0.001
New-onset burnout	0.480	0.092	1.616	1.351	1.935	< 0.001
New-onset or worsening Insomnia	1.64	0.075	5.155	4.45	5.978	< 0.001

Figure S1: Heat map showing the distributions of the Health Professionals participants in Brazil.




Figure S2: Distribution of health professionals according to their specific occupations.




Figure S3: Mean work hours change (per week) according to their specific occupations.




Figure S4: The frequency of healthcare professional (per category) that reported recent or current care of patients with COVID-19.




Figure S5: The frequency health professional (per category) that got a previous COVID-19 infection.




Figure S6: Frequency of prevalent anxiety among the health professional categories.




Figure S7: Frequency of Burnout development among the health professional categories.




Figure S8: Frequency of family income changes among the health professional categories.




Figure S9: Distribution of new-onset insomnia or impairment of previous insomnia according to the health professional category.




Figure S10: Frequency of new pharmacological treatment according to the health professional category.

