




Infliximab use (% of hospitalized patients with COVID-19) by month




Famotidine use (% of hospitalized patients with COVID-19) by month







Colchicine use (% of hospitalized patients with COVID-19) by month




Aspirin use (% of hospitalized patients with COVID-19) by month







Enoxaparin use (% of hospitalized patients with COVID-19) by month




Tranexamic acid use (% of hospitalized patients with COVID-19) by month




Warfarin use (% of hospitalized patients with COVID-19) by month




Clopidogrel use (% of hospitalized patients with COVID-19) by month




Cilostazol use (% of hospitalized patients with COVID-19) by month




Prazosin use (% of hospitalized patients with COVID-19) by month




Losartan use (% of hospitalized patients with COVID-19) by month




Heparin use (% of hospitalized patients with COVID-19) by month




Metformin use (% of hospitalized patients with COVID-19) by month




Dexamethasone use (% of hospitalized patients with COVID-19) by month




Sitagliptin use (% of hospitalized patients with COVID-19) by month




Amoxicillin use (% of hospitalized patients with COVID-19) by month




Azithromycin use (% of hospitalized patients with COVID-19) by month




Lopinavir use (% of hospitalized patients with COVID-19) by month




Ritonavir use (% of hospitalized patients with COVID-19) by month




Ribavirin use (% of hospitalized patients with COVID-19) by month




Hydroxychloroquine use (% of hospitalized patients with COVID-19) by month




Ceftriaxone use (% of hospitalized patients with COVID-19) by month




Ivermectin use (% of hospitalized patients with COVID-19) by month




Oseltamivir use (% of hospitalized patients with COVID-19) by month




Vitamin C use (% of hospitalized patients with COVID-19) by month




H2 receptor antagonist use (% of hospitalized patients with COVID-19) by month



DPP-4 inhibitors use (% of hospitalized patients with COVID-19) by month




Vitamin D use (% of hospitalized patients with COVID-19) by month





Immunoglobulins use (% of hospitalized patients with COVID-19) by month



Alpha-1 blockers use (% of hospitalized patients with COVID-19) by month

